



Evaluation of resistance to fluoroquinolones and determination of mutations in *gyrA* and *parC* genes in *Escherichia coli* isolated from raw milk of dairy cows with coliform mastitis in Khorasan Razavi province, Iran

Mahdie Mahdavi^a, Behrooz Fathi^b, Abdollah Jamshidi^c, Babak Khoramian^d

^a Graduated from Faculty of Veterinary Medicine, Ferdowsi University of Mashhad, Mashhad, Iran.

^b Department of Basic Sciences, Faculty of Veterinary Medicine, Ferdowsi University of Mashhad, Mashhad, Iran.

^c Department of Food Hygiene and Aquaculture, Faculty of Veterinary Medicine, Ferdowsi University of Mashhad, Mashhad, Iran.

^d Department of Clinical Sciences, Faculty of Veterinary Medicine, Ferdowsi University of Mashhad, Mashhad, Iran.

ABSTRACT

The present study was performed to assess the resistance profile to fluoroquinolone and to determine mutations in *gyrA* and *parC* genes of *Escherichia coli* in bovine coliform mastitis. Fluoroquinolones (norfloxacin (NOR), ciprofloxacin (CIP), enrofloxacin (NFX), levofloxacin (LEV), and ofloxacin (OFL)) were tested against *E. coli* isolates, isolated from bovine mastitis (100 milk samples) by disk diffusion method. To determine the extent of *gyrA* and *parC* mutations associated with fluoroquinolone resistance in *E. coli*, two isolates with the highest resistance to each fluoroquinolone were submitted for the amplification and sequencing of the quinolone resistance-determining regions (QRDRs) of *gyrA* and *parC* genes. The disk diffusion method indicated that *E. coli* isolates had the highest intermediate resistance to OFL (16.7%), followed by NFX and NOR (15%), while they had low resistance to CIP and LEV (3.33%). A few silent mutations in *gyrA* (in codons 91, 100, 111, 131, 132) and in *parC* (in codons 91, 157, 159) were detected in QRDRs, and mutations in nucleotides 65, 80, and 83 in *gyrA*, and 195, 209, 212 in *parC* were detected in the other isolate. These results showed an intermediate rate of resistance to fluoroquinolones in *E. coli* isolates from raw milk of cows with coliform mastitis.

Keywords

Escherichia coli; fluoroquinolone resistance; *gyrA* gene; mastitis; *parC* gene

Number of Figures: 7
Number of Tables: 3
Number of References: 37
Number of Pages: 9

Abbreviations

E. coli: *Escherichia coli*

FQ: Fluoroquinolone

QRDR: Quinolone resistance determining region

MC: MacConkey

EMB: Eosin Methylene Blue

Introduction

Mastitis is considered one of the most important diseases in dairy animals that causes severe losses to the dairy industry [1]. The economic losses due to clinical mastitis include production loss, lower milk yield and value, treatment expenses, and loss of animal value [2]. Coliforms such as *Escherichia*, *Klebsiella spp.*, and *Enterobacter spp.* are the most common etiological agents causing clinical mastitis [3]. *Escherichia coli* is the most common species isolated from coliform mastitis which is a Gram-negative, non-spore-forming rod bacterium that belongs to the family *Enterobacteriaceae* [4, 5]. Clinical signs of *E. coli* mastitis include a wide range from a mild disease with only local inflammation changes in the mammary gland to severe with systemic signs, generally including high fever, increased pulse frequency, lack of appetite, decreased milk production, dehydration, rumen stasis, shock, and death [6, 7]. In cases of mild to moderate *E. coli* mastitis, the use of anti-inflammatory drugs and supportive treatments is recommended. In peracute or acute cases of *E. coli* mastitis, due to the potential risk of bacterial growth in the mammary gland, which in turn may lead to bacteremia, administration of broad-spectrum antimicrobials is recommended to reduce the number of bacteria [8].

The fluoroquinolones are broad-spectrum and bactericidal antibiotics. They are used against gram-positive and especially gram-negative bacteria such as members of the *Enterobacteriaceae* family [9, 10]. They block DNA synthesis by targeting bacterial DNA gyrase and topoisomerase IV, both of which are essential for bacterial DNA supercoiling as the replicating strands separate [11]. DNA gyrase and topoisomerase IV are tetrameric structures composed of two pairs of subunits. The four subunits of DNA gyrase include 2 monomers of A and 2 monomers of B, with the names GyrA and GyrB, respectively. The topoisomerase IV also has ParC and ParE subunits, which are encoded by *parC* and *parE* genes, respectively [12].

The major mechanisms of resistance to quinolone antibiotics include mutations that occur at the target drug sites, mutations that reduce drug accumulation, and plasmid-mediated quinolone resistance [13]. The most common mechanism that produces significant levels of clinical resistance to fluoroquinolones is an alteration in the target enzymes. These changes are caused by self-mutations occurring within the responsible genes. Resistance to fluoroquinolones is due to the substitution of amino acids in a certain region of GyrA or ParC subunits [13, 14]. The broad-spectrum activity of quinolones against various infections

and the widespread use of these antibiotics, the abuse and unnecessary use of them, especially in developing countries, has accelerated the development of resistance mechanisms [15].

Fluoroquinolones are used in the treatment of infectious diseases, including coliform mastitis caused by *E. coli*. Since the drug resistance pattern has regional distribution, determination of this pattern of *E. coli* resistance can be used to determine the appropriate treatment regimen for clinical coliform mastitis [16]. This study aimed to determine the resistance pattern of *E. coli* isolated from cows with coliform mastitis to some fluoroquinolones and also to detect the mutations in QRDR of fluoroquinolone-resistance *E. coli* isolates.

Results

Identification of *E. coli*


In this study, 100 milk samples were subjected to isolation of *E. coli* by selective plating followed by streaking on the Eosin Methylene Blue (EMB) agar at 37 °C for 24 h. Typical colonies of *E. coli* were produced from 45 samples. These 45 presumptive *E. coli* isolates on EMB agar were confirmed by biochemical tests (Figure 1).

Susceptibility testing


The results of susceptibility study showed that less than 20% of *E. coli* isolates had intermediate resistance to each antibiotic (Figure 2). Intermediate resistance was 3.33% to ciprofloxacin and levofloxacin, 15% to enrofloxacin and norfloxacin, and 16.7% to ofloxacin (Table 1).

Identification of *gyrA* and *parC* mutations in clinical isolates of *E. coli*

Amplification of the QRDRs of *gyrA* and *parC* genes was performed by PCR (Figure 3). The results of the DNA sequencing of *gyrA* and those of *parC* were consistent and provided information from both standards for a region between nucleotides 247 to 840 (corresponding to codons 82 to 280) of *gyrA* and from nucleotides 167 to 539 (corresponding to codons 55 to 180) of the *parC* gene, respectively (Table 2, Figures 4, 5, 6, and 7). Accession numbers of *E. coli* isolates based on QRDRs of *gyrA* and *parC* genes deposited in the GenBank are as follows: SRX5988183, SRX5982112 for sample number 2968, and SRR17711097, SRR17711096 for sample number 3077 (Accession number to cite for these SRA data: PRJNA547542).

**Figure 1.**

Isolation and identification of *E. coli*. A) Triple Sugar Iron agar, acid/acid reaction with gas production and no H₂S. B) Simmons citrate agar, the medium remained green. This is a negative result for citrate test. C) Sulfur Indole Motility (SIM) medium, *E. coli* is hydrogen sulfide negative, indole positive, and the cloudy appearance of the medium indicates that *E. coli* is motile. D) MacConkey agar, pink colonies. E) Eosin Methylene Blue agar, colonies of purple with black center and green metallic sheen.

**Figure 2.**

The results of evaluation of resistance to fluoroquinolones with disk diffusion method. The scale is in mm.

Table 1.
Antibiotic resistance pattern of 60 *E. coli* isolates.

Antibiotic (µg)	No. of sensitive isolates	% sensitive isoates	% intermediate resistant isolates	% resistant isolates
Ofloxacin (5)	50	83.3	16.7	-
Enrofloxacin (5)	51	85	15	-
Norfloxacin (10)	51	85	15	-
Ciprofloxacin (5)	58	96.6	3.33	-
Levofloxacin (5)	58	96.6	3.33	-

Discussion

The focus of the current study was to assess the resistance of *E. coli* to some fluoroquinolones in bovine coliform mastitis and to generate the fluoroquinolones resistance profile of isolates. The level of resistance of *E. coli* isolates to enrofloxacin and norfloxacin was similar to the level of resistance to ofloxacin. This is due to the development of cross-resistance to one of the fluoroquinolones. In the present study, among all 60 *E. coli* isolates, less than 20% of isolates had intermediate resistance to fluoroquinolones. This is still a

relatively low figure compared with other published studies, in which the proportion of resistant isolates has ranged from 23% to 63% [15, 17-19]. In the results from Su et al. (2016) *E. coli* isolates showed 4% resistance to ciprofloxacin and levofloxacin; similarly, *E. coli* isolates in the present study showed only 3.33% resistance to ciprofloxacin and levofloxacin whereas Metzger and Hogan (2013) found 12% of *E. coli* isolated from bovine milk samples were non-susceptible to ciprofloxacin [20, 21]. Among fluoroquinolones, enrofloxacin and norfloxacin resistance were found in nine (15%) *E. coli* isolates, and all other isolates




Figure 3. PCR amplification of QRDR of gyrA and parC genes for *E. coli*. Lane M: DNA marker; 100 bp plus. Lane 1, 2: test isolates. The expected product size of gyrA is 253 bp and the expected product size of parC is 434 bp.

Score	Expect	Identities	Gaps	Strand
261 bits(141)	2e-73	151/156(97%)	0/156(0%)	Plus/Minus
NC_000913.3 gyrA_2968	2227 168	CAGTTCATGGGCAATTTTCGCCAGACGGATTTC	CGTATAACGCATTGCCGCCGAGAGTC	2286 109
	 TT		
NC_000913.3 gyrA_2968	2287 108	GCCGTCGATAGAACC GAAGTTACCCTGACCGT	CTACCAGCATATAACGCAGCGAGAATGG	2346 49
	 G G	
NC_000913.3 gyrA_2968	2347 48	CTGCGCCATGCGGACGATCGTGTTCATAGACCG	CCGA	2382 13
	 A		

Figure 4. The result of alignment of the gyrA gene of sample number 2968. The red letters indicate nucleotide changes. Reference sequence is from nucleotide 2227 to 2382 from NC_000913.3:2336793-2339420 Escherichia coli str. K-12 substr. MG1655, complete genome.

Score	Expect	Identities	Gaps	Strand
678 bits(367)	0.0	385/393(98%)	3/393(0%)	Plus/Plus
NC_000913.3 parC_2968	156 10	GGGCCTGAATGCCAGCGCCAAATTTAAAAAAT	-CGGCCCGTACCGTCGGTGACGTACTGG	214 67
	 C		
NC_000913.3 parC_2968	215 68	GTAATAACCATCCGCACGGCGATAGCGCCTG	TATGAAGCGATGGTCTGATGGCGCAAC	274 127
	 G		
NC_000913.3 parC_2968	275 128	CGTTCTCTTACCGTTATCCGCTGGTTGATGGT	CAGGGGAACTGGGGCGCGCCGGACGATC	334 187
			
NC_000913.3 parC_2968	335 188	CGAAATCGTTCGCGGCAATGCGTTACACCGAAT	CCCGGTTGTCGAAATATTCGAGCTGC	394 247
			
NC_000913.3 parC_2968	395 248	TATTGAGCGAGCTGGGGCAGGGGACGGCTGACT	GGGTGCCAAACTTCGACGGCACTTTGC	454 307
			
NC_000913.3 parC_2968	455 308	AGGAGCCGAAAATGCTACCTGCCCGTCTGCCAA	ACATTTTGCTTAACGGCACCACCGGTA	514 367
	 G T		
NC_000913.3 parC_2968	515 368	TTGCCGTCGGCATGGCGACCGATATTCCACCGC		547 400
	 T C		

Figure 5. The result of alignment the parC gene of sample number 2968. The red letters indicate nucleotide changes. Reference sequence is from nucleotide 156 to 547 from NC_000913.3:c3165973_3163715 Escherichia coli str. K_12 substr. MG1655, complete genome.

Score	Expect	Identities	Gaps	Strand
313 bits(169)	3e-89	204/221(92%)	1/221(0%)	Plus/Plus
NC_000913.3 gyrA_3077	2204 5	GTCTCTTTTTTCGAGATCGGCCATCAGTTCATGGGCAATTTTCGCCAGACGGATTTCCGTA		2263 64
NC_000913.3 gyrA_3077	2264 65	TAACGCATTGCCGCCGAGAGTCGCCGTCGATAGAACCGAAGTTACCTGACCGTCTACC		2323 124
NC_000913.3 gyrA_3077	2324 125	AGCATATAACGCAGCGAGAATGGCTGCGCCATG-CGGACGATCGTGTTCATAGACCGCCGA		2382 184
NC_000913.3 gyrA_3077	2383 185	GTCACCATGGGGATGGTATTTACCGATTACGTACCCAACGA	2423	
		.A.....GAA.....	225	

Figure 6. The result of alignment the gyrA gene of sample number 3077. The red letters indicate nucleotide changes. Reference sequence is from nucleotide 2204 to 2423 from NC_000913.3:2336793-2339420 Escherichia coli str. K-12 substr. MG1655, complete genome

Score	Expect	Identities	Gaps	Strand
614 bits(332)	3e-180	359/371(97%)	5/371(1%)	Plus/Plus
NC_000913.3 parC_3077	141 26	TGCGATGTCTGAACTGGGCCTGAATGCCAGCGCCAAATTTAAAAAATCGGCCCGTACCCT		200 82
NC_000913.3 parC_3077	201 83	CGGTGACGTACTGGGTAAATACCATCCGCACGGCGATAGCGCCTGTTATGAAGCGATGGT		260 142
NC_000913.3 parC_3077	261 143	CCTGATGGCGCAACCGTTCTCTTACCGTTATCCGCTGGTTGATGGTCAGGGGAACTGGGG		320 202
NC_000913.3 parC_3077	321 203	CGCGCCGGACGATCCGAAATCGTTCGCGGCAATGCGTTACACCGAATCCCGGTTGTCGAA		380 262
NC_000913.3 parC_3077	381 263	ATATTCCGAGCTGCTATTGAGCGAGCTGGGGCAGGGGACGGCTGACTGGGTGCCAAACTT		440 322
NC_000913.3 parC_3077	441 323	CGACGGCACTTTGCAGGAGCCGAAAATGCTACCTGCCCGTCTGCCAAACATTTTGCTTAA		500 381
NC_000913.3 parC_3077	501 382	CGGCACCACCG	511	
		A.-.....	391	

Figure 7. The result of alignment the parC gene of sample number 3077. The red letters indicate nucleotide changes. Reference sequence is from nucleotide 141 to 511 from NC_000913.3:c3165973_3163715 Escherichia coli str. K_12 substr. MG1655, complete genome.

were susceptible to norfloxacin and enrofloxacin.

This is in general agreement with Malinowski et al. (2008) who found that 16.1% and 14.9% mastitis *E. coli* isolates from Poland were resistant to enrofloxacin and norfloxacin, respectively [22]. However, in a study in Bangladesh, no resistance to fluoroquinolones including ofloxacin, ciprofloxacin, and levofloxacin was reported in *E. coli* isolated from milk of mastitis cattle [23]. Persson et al. (2011) reported that there was no fluoroquinolone resistance in *E. coli* isolated from milk samples of cows with mastitis [24]. In another study by Persson and her colleagues in Sweden (2015), they reported that all isolates (n=57) of *E. coli* from dairy cows with acute clinical mastitis were susceptible to

enrofloxacin [25]. Armanullah et al. (2018), studied the antibiotic resistance profile of *E. coli* isolates from bovine clinical mastitis and reported resistance to ciprofloxacin (16.67%), norfloxacin (8.33%), ofloxacin (8.33%), and intermediate resistance to norfloxacin (8.33%) that was somewhat similar to the finding of the present study [26].

Fluoroquinolone resistance of *E. coli* isolates from bovine mastitis has been studied by several authors and the results have varied, which may be due to different methods and breakpoints used to determine susceptibility. Resistance to fluoroquinolones is still uncommon among *E. coli* isolated from bovine mastitis. In comparison to other studies [15, 17-19, 27], the

results of this study showed a low level of resistance to fluoroquinolones, which may be due to the controlled use of these antibiotics. However, in the present study ciprofloxacin and levofloxacin were proved to be the best antibiotics to treat *E. coli* mastitis in cattle since they were highly effective.

In the present study, the *E. coli* isolates did not have resistance to fluoroquinolones and the rate of intermediate resistance to fluoroquinolones was very low. It is generally accepted that *gyrA* mutations play a major role in the development of fluoroquinolone resistance in *E. coli*, while the mutations in the *parC* gene are additionally associated with resistance [28]. To analyze the correlation between genetic characterization and resistance phenotype, two isolates with the most resistance to each fluoroquinolone were submitted to amplification and sequencing of the QRDR in *gyrA* and *parC* genes. There were two silent mutations in the *gyrA* gene at wobble position in codons 91 and 100; similarly, *E. coli* isolates in the Heisig study showed silent mutations in codons 91 and 100 [28]. Mutation at codons 83 and 87 was found to be the most common *gyrA* mutations of *E. coli* in several studies, and in the present study, there was a mutation in codon 83 of *gyrA* in sample number 3077 [29-32]. In addition, we found a silent mutation in codon 91 in the *parC* gene. Similarly, *E. coli* isolates in the Heisig study showed silent mutation only in codon 91, whereas the most common mutations in *parC* were reported at codons 80, 84, and 87 [19, 28, 31, 32].

In conclusion, the current investigation showed that most *E. coli* isolates isolated from raw milk of cows with coliform mastitis in Khorasan Razavi province were sensitive to fluoroquinolones and some *E. coli* isolates had intermediate resistance to fluoro-

quinolones. In *gyrA* and *parC* genes of *E. coli* isolates with the most intermediate resistance to studied fluoroquinolones, there were silent mutations and mutations. There is some evidence that silent mutations can especially affect the regulation of transcription [33-35].

Materials & Methods

Sample collection

A total of one hundred (100) milk samples were examined in this study. Samples were collected from the milk of dairy cattle with clinical mastitis of three dairy farms in Mashhad (Khorasan Razavi province, Iran). Fifteen isolates of *E. coli* were obtained from "Bacterial Collection of the Mastitis Laboratory", Veterinary Teaching Hospital, Faculty of Veterinary Medicine, Ferdowsi University of Mashhad.

Isolation of *E. coli*

Milk samples were cultured on MacConkey agar media (Merk, Darmstadt, Germany) and were incubated at 37 °C for 24 h. Suspected *E. coli* lactose-fermenting colonies (pink colonies) were used for culture on the Eosin methylene blue (EMB) agar (Merk, Darmstadt, Germany). The appearance of the colonies of purple with black center and green metallic sheen were considered positive for *E. coli* on EMB agar and were selected for further studies. The colonies of presumptive *E. coli* on EMB agar were confirmed by standard biochemical tests, including triple sugar iron agar, Simmons citrate agar, and motility assay.

Antibiotic susceptibility study

Antibiotic susceptibility testing was carried out with equivalence of 0.5 McFarland turbidity standard by agar disk diffusion method on Mueller-Hinton agar (Himedia, Mumbai, India) plates following the Clinical and Laboratory Standards Institute [36]. All *E. coli* isolates were subjected to an antibiotic susceptibility test. The antimicrobial disks (Padtan Teb, Tehran, Iran) used in the experiment included 5 µg ciprofloxacin, 10 µg norfloxacin, 5 µg levofloxacin, 5 µg ofloxacin, and 5 µg enrofloxacin. The antibiotic

Table 2.
Mutations in genes *gyrA* and *parC*.

<i>E. coli</i> isolate	<i>gyrA</i> mutation			<i>parC</i> mutation		
	Codon position	Nucleotide exchange	Amino acid exchange	Codon position	Nucleotide exchange	Amino acid exchange
2968	91	CGT → CGC	ginine ^a	91	CAG → CAA	Glutamine ^a
	100	TAC → TAT	Tyrosine ^a	157	CTG → CTA	Leucine ^a
	111	TCC → TCT	Serine ^a	159	GCT → GCC	Alanine ^a
	131	GCA → GCC	Alanine ^a			
	132	AAT → CAT	Asparagine → Histidine			
3077	65	AAT → CAT	Asparagine → Histidine	195	GGT → GTT	Glycine → Valine
	80	GCA → TCA	Alanine → Serine	209	GTG → TTA	Valine → Leucine
	83	AGG → GGG	Arginine → Glycine	212	GGC → CAC	Glycine → Histidine

^a Silent mutation

disks were placed on Mueller-Hinton agar culture plate. The plates were incubated for 18-24 h at 37 °C. The size of the zone of inhibition was recorded and resistance zone diameter breakpoints adopted for these antimicrobials were the following: ≤ 15 mm for ciprofloxacin, ≤ 12 mm for norfloxacin, ≤ 13 mm for levofloxacin, ≤ 12 mm for ofloxacin, and ≤ 14 mm for enrofloxacin.

DNA extraction

E. coli isolates were grown overnight in Nutrient agar (Merk, Darmstadt, Germany) at 37 °C. One colony was suspended in 250 µL of sterile distilled water. After boiling the suspension for 15 min, followed by freezing and subsequent centrifugation at 14000 rpm for 15 min, the cell debris was pelleted and the supernatant was used as a template for the amplification reaction. [37].

Amplification of quinolone resistance determining regions (QRDRs)

Polymerase chain reaction (PCR) was used to amplify QRDR of *gyrA* and *parC* for mutation detection. The list of primers that were used for amplification of *gyrA* and *parC* genes is shown in Table 3. The PCR amplification was performed in a total reaction volume of 25 µL. The reaction mixture contained 12.5 µL of 2x master mixtures (CinnaGen, Tehran, Iran), 1 µL of each forward

and reverse primer (10 pmol/µL), 8.5 µL of deionized water and 2 µL of DNA template. The PCR program included initial denaturation at 94 °C for 5 min followed by 30 cycles of denaturation (94 °C for 1 min), annealing (55 °C for *gyrA* and 56 °C for *parC* for 1 min), and extension (72 °C for 1 min) with a final extension at 72 °C for 5 min. For amplification of DNA, the PCR was performed with a thermocycler (Techne, Chelmsford, UK). The PCR products were run on a 1% agarose gel in TAE buffer at 100 V for 45 min. After electrophoresis (Padideh Nojen Pars, Mashhad, Iran) in the agarose gel and staining with the green viewer (Sinacolor, Tehran, Iran), they were observed and documented under gel documentation system (Kimiagene, Mashhad, Iran). A 100 bp plus DNA ladder was used to determine the molecular size of the PCR products. Primers used in the study were custom synthesized from Macrogen Inc. (South Korea).

Sequencing and Alignment

The PCR product of *gyrA* and *parC* genes with forward and reverse primers sent for sequencing to Microsynth (Switzerland). DNA sequences were analysed using Chromas software. DNA sequence data were compared to data in the GenBank database using the BLAST algorithm available at the National Center for Biotechnology Information (NCBI) website (www.ncbi.nlm.nih.gov).

Table 3.

The primers were used for amplification of *gyrA* and *parC*

Primer name	Primer direction ^a	Sequence (5' to 3')	Product size (bp)	Annealing temperature (°C)	Reference
<i>gyrA4</i>	F	TCGTTGGTGACGTAATCGGT	253	55	31
<i>gyrA5</i>	R	TCCGTGCCGTCATAGTTATC	253	55	31
<i>parC1</i>	F	AACCTGTTTCAGCGCCGCATT	434	56	31
<i>parC2</i>	R	ATGCGGTGGAATATCGGTTCG	434	56	31

^aF, forward; R, reverse

Authors' Contributions

B.F., A.J., and B.KH. conceived and planned the experiments. M.M. carried out the experiments. B.F., A.J. and B.KH. planned and carried out the simulations. M.M., K.L., and B.KH. contributed to sample preparation. B.F., A.J., and B.KH. contributed to the interpretation of the results. B.F. took the lead in writing the manuscript. All authors provided critical feedback and helped shape the research, analysis, and the manuscript.

Acknowledgements

We wish to thank H.K. Rahmani, A. Kargar and N. Shahbazi for their technical assistance. This research was financially supported by Ferdowsi University of Mashhad, Mashhad, Iran.

Conflict of interest

The authors declare that they have no competing interests.

References

- Janzen J. Economic losses resulting from mastitis. A review. *J Dairy Sci.* 1970; 53(9):1151-60.
- Jingar SC, Mahendra S, Roy AK. Economic losses due to clinical mastitis in cross-bred cows. *Dairy Vet Sci J.* 2017; 3(2):555606.
- Hogan J, Smith KL. Coliform mastitis. *Vet Res.* 2003; 34(5):507-19.
- Cullor JS, Smith WL. Endotoxin and disease in food animals. *Compend. Contin. Educ. Vet.* 1996;18(1): 31-38.
- Bradely AJ, Leach KA, Breen JE, Green LE, Green MJ. Survey of the incidence and etiology of mastitis in dairy farms in En-

- gland and Wales. *Vet Rec.* 2007; 160:253-258.
6. Wenz JR, Barrington GM, Garry FB, Dinsmore RP, Callan RJ. Use of systemic disease signs to assess disease severity in dairy cows with acute coliform mastitis. *J Am Vet Med Assoc.* 2001; 218(4):567-72.
 7. Burvenich C, Van Merris V, Mehrzad J, Diez-Fraile A, Duchateau L. Severity of *E. coli* mastitis is mainly determined by cow factors. *Vet Res.* 2003; 34:521-564.
 8. Suojala L, Kaartinen L, Pyörälä S. Treatment for bovine *Escherichia coli* mastitis – an evidence based approach. *J Vet Pharmacol Ther.* 2013; 36(6):521-31.
 9. Sharfaraj Nawaz M, Bodla R, Kant R, Pratab Singh S, Bhtani R, Kapoor G. Fluoroquinolone as antimicrobial agent: A Review. *Int J Pharm Sci Res.* 2017; 2(3):57-63.
 10. Suh B, Lorber B. Quinolones. *Med Clin North Am.* 1995; 79(4):869-894.
 11. Hooper DC. Mechanisms of action of antimicrobials: focus on fluoroquinolones. *Clin. Infect. Dis.* 2001; 32(1): 9-15.
 12. Higgins PG, Fluit AC, Schmitz FJ. Fluoroquinolones: structure and target sites. *Curr. Drug. Targets.* 2003; 4(2): 181-190.
 13. Hooper DC. Mechanisms of quinolone resistance. In: Hooper DC, Rubinstein E (Eds.), *Quinolone antimicrobial agents.* (3rd Edn.), Washington, D.C, USA: American Society of Microbiology Press. 2003; PP:41-67.
 14. Hopkins KL, Davies RH, Threlfall EJ. Mechanisms of quinolone resistance in *Escherichia coli* and *Salmonella*: recent developments. *Int J Antimicrobial Agents.* 2005; 25:358–373.
 15. Saranya K, Pavulraj S, Kalaiselvi L, Amsaveni S, Ramesh S. Antibacterial susceptibility profiles of coliforms isolated from bovine subclinical and clinical mastitis against fluoroquinolones. *Tamilandu Journal. Vet Anim Sci.* 2013; 9 (4):279-284.
 16. Srinivasan V, Gillespie BE, Lewis MJ, Nguyen LT, Headrick SI, Schukken YH, Oliver SP. Phenotypic and genotypic antimicrobial resistance patterns of *Escherichia coli* isolated from dairy cows with mastitis. *Vet. Microbiol.* 2007; 124(3-4): 319-328.
 17. Alekish M, Al-Qudah K, Al-Saleh A. Prevalence of antimicrobial resistance among bacterial pathogens isolated from bovine mastitis in northern Jordan. *Rev Vet Med.* 2013; 164:319-26.
 18. Chandrasekaran D, Venkatesan P, Tirumurugan K, Nambi A, Thirunavukkarasu P, Kumanan K, et al. Pattern of antibiotic resistant mastitis in dairy cows, *Vet World.* 2014; 7(6):389-394.
 19. Balakrishnan S, Antony PX, Mukhopadhyay HK, Pillai RM, Thanisslass J, Padmanaban V, et al. Genetic characterization of fluoroquinolone-resistant *Escherichia coli* associated with bovine mastitis in India. *Vet World.* 2016; 9(7):705-709.
 20. Su Y, Yu C-Y, Tsai Y, Wang S-H, Lee C, Chu C. Fluoroquinolone-resistant and extended-spectrum β -lactamase-producing *Escherichia coli* from the milk of cows with clinical mastitis in Southern Taiwan. *J Microbiol Immunol Infect.* 2016;49(6):892-901.
 21. Metzger SA, Hogan JS. Short communication: Antimicrobial susceptibility and frequency of resistance genes in *Escherichia coli* isolated from bovine mastitis. *J Dairy Sci.* 2013; 96:3044–3049.
 22. Malinowski E, Lassa H, Smulski S, Kłossowska A, Kaczmarowski M. Antimicrobial susceptibility of bacteria isolated from cows with mastitis in 2006-2007. *Bulletin of the Veterinary Institute in Pulawy.* 2008; 52(565):72.
 23. Tanzin T, Nazir KHMNH, Zahan MN, Md. Parvej S, Zesmin K, Rahman MT. Antibiotic resistance profile of bacteria isolated from raw milk samples of cattle and buffaloes. *J Adv Vet Anim Res.* 2016; 3 (1): 62-67.
 24. Persson Y, Nyman AK, Grönlund-Andersson U. Etiology and antimicrobial susceptibility of udder pathogens from cases of subclinical mastitis in dairy cows in Sweden. *Acta Vet Scand.* 2011; 53(1):36.
 25. Persson Y, Katholm J, Landin H, Mörk Mj. Efficacy of enrofloxacin for the treatment of acute clinical mastitis caused by *Escherichia coli* in dairy cows. *Vet Rec.* 2015; 176(26):673-673.
 26. Armanullah MD, Anjay Dr, Kumar P, Kumari S, Kaushik P, Archana S, Arya S. Prevalence of Multi-Drug Resistant (MDR) *Escherichia coli* in bovine clinical samples. *Int. J. Curr. Microbiol. App. Sci.* 2018; 7: 1476-1485.
 27. Firouzi R, Rajaian H, Tabaei I, Saeedzadeh A. In vitro antibacterial effects of marbofloxacin on microorganisms causing mastitis in cows. *J Vet Res.* 2010; 65(1):51-55.
 28. Heisig P. Genetic evidence for a role of *parC* mutations in development of high-level fluoroquinolone resistance in *Escherichia coli*. *Antimicrob. Agents Chemother.* 1996; 40(4): 879-85.
 29. Chen JY, Siu LK, Chen YH, Lu PL, Ho M, Perg CF. Molecular epidemiology and mutations at *gyrA* and *parC* genes of ciprofloxacin-resistant *Escherichia coli* isolates from Taiwan medical center. *Microb Drug Resist.* 2001; 7, 47-53.
 30. Ruiz J, Gomez J, Navia MM, Ribera A, Sierra JM, Marco F. High prevalence of nalidixic acid resistant, ciprofloxacin susceptible phenotype among clinical isolates of *Escherichia coli* and other Enterobacteriaceae. *Diagn Microbiol and infect Dis.* 2002; 42, 257-61.
 31. Ogbolu DO, Daini O, Ogunledun A, Alli AT, Olusoga-Ogbolu F, Webber M. Effects of *gyrA* and *parC* mutations in quinolone-resistant *Escherichia coli* associated with bovine mastitis in India. *Vet World.* 2016; 9(7):705-709.

- lones resistant clinical gram negative bacteria from Nigeria. Afr J Biomed Res. 2012; 15(2):97-104.
32. Sáenz Y, Zarazaga M, Briñas L, Ruiz-Larrea F, Torres C. Mutations in *gyrA* and *parC* genes in nalidixic acid-resistant *Escherichia coli* strains from food products, humans and animals. J Antimicrob Chemother. 2003; 51(4):1001-5.
33. Yadegari H, Biswas A, Akhter MS, Driesen J, Ivaskevicius V, Marquardt N, Oldenburg J. Intron retention resulting from a silent mutation in the VWF gene that structurally influences the 5' splice site. Blood. 2016; 128(17):2144-2152.
34. Agashe D, Sane M, Phalnikar K, Diwan GD, Habibullah A, Martinez-Gomez NC, et al. Large-effect beneficial synonymous mutations mediate rapid and parallel adaptation in a bacterium. Mol Biol Evol. 2016; 33(6):1542-53.
35. Hauber DJ, Grogan DW, DeBry RW. Mutations to less-preferred synonymous codons in a highly expressed gene of *Escherichia coli*: fitness and epistatic interactions. PLoS One 2016; 11(1):e0146375.
36. CLSI. Performance standards for antimicrobial susceptibility testing. 27th ed. CLSI supplement M100. Wayne, PA: Clinical and Laboratory Standards Institute; 2017.
37. Reischl U, Youssef MT, Kilwinski J, Lehn N, Zhang WL, Karch H, et al. Real-time fluorescence PCR assay for detection and characterization of shiga toxin, intimin, and enterohemolysin genes from shiga toxin-producing *Escherichia coli*. J Clin Microbiol. 2007; 40(7):2555-2565.

COPYRIGHTS

©2022 The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited. No permission is required from the authors or the publishers.

**How to cite this article**

Mahdavi M, Fathi B, Jamshidi A, Khoramian B. Evaluation of fluoroquinolones resistance and determination of mutations in *gyrA* and *parC* genes in *Escherichia coli* isolated from raw milk of dairy cows with coliform mastitis in Khorasan Razavi province, Iran. Iran J Vet Sci Technol. 2022; 14(1): 20-28.
DOI: <https://doi.org/10.22067/ijvst.2022.71423.1059>
URL: https://ijvst.um.ac.ir/article_42057.html