



Foodborne Bacteria in Iran: A 23-year Systematic Review of High-risk Foods

Mohammad Hashemi,^{a,b} Fateme Asadi Touranlou,^{a,b} Shiva Adibi,^{a,b} Asma Afshari,^{a,b} Golshan Shakeri^b

^a Medical Toxicology Research Center, Mashhad University of Medical Sciences, Mashhad, Iran.

^b Department of Nutrition, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.

ABSTRACT

Foodborne diseases are a significant global public health concern due to their high prevalence, mortality, and economic losses. The aim of this study was to conduct a systematic review of identified foodborne pathogens and outbreaks in Iran over the past 23 years to provide an overview of the risk assessment and prevention approaches in the country. Using appropriate keywords and searching major databases, such as ScienceDirect, Scopus, PubMed, Google Scholar, and the Iranian Scientific Knowledge Database, we initially identified 4,740 articles. Finally, 328 articles were selected for evaluation. Among these articles, publications on *Salmonella*, *Staphylococcus aureus*, and *Listeria* were the most numerous. Poultry meat was found to be the main source of major foodborne pathogens in Iran, including *Campylobacter* (46.21%), *Listeria monocytogenes* (38.45), *Salmonella* (24.83%), and *Yersinia enterocolitica* (16.81%). Given the high prevalence of foodborne bacteria in Iranian foods, it is crucial to implement effective control measures to reduce the risk and burden of foodborne diseases. In particular, poultry meat, which poses a high risk for the occurrence of foodborne diseases in Iran, should be subjected to further risk assessment and control measures throughout the food chain.

Keywords

Foodborne bacteria, *Salmonella*, *Staphylococcus aureus*,
Listeria, food, prevalence

Abbreviations

EWHO: World Health Organization
CDC: Center for Disease Control and Prevention
RTE: Ready-To-Eat
EFSA: European Food Safety Authority

Number of Figures: 13
Number of Tables: 12
Number of References: 369
Number of Pages: 47

Introduction

Foodborne diseases typically result from the consumption of food or water contaminated with pathogens or their toxins [1]. These illnesses often present as acute health problems with diverse symptoms, such as gastrointestinal distress (e.g., diarrhea, vomiting, nausea, and abdominal cramps) or neurological symptoms (e.g., headache, paralysis, and paresthesia) [2, 3]. The bacterial pathogens most commonly associated with foodborne illness worldwide include *Listeria monocytogenes*, *Escherichia coli* O157:H7, *Staphylococcus aureus*, *Salmonella enterica*, *Bacillus cereus*, *Vibrio* spp., *Campylobacter jejuni*, and *Clostridium perfringens* [2, 4, 5]. The food products most frequently implicated in outbreaks include poultry, ground meat, seafood, dairy products, as well as fruits and vegetables [6].

The food industry faces significant challenges in ensuring the safety and nutritional quality of food products for consumers due to various sources of contamination, such as animals, soil, water, air, and food handlers during production and storage [7, 8]. However, the implementation of proper cold preservation methods (e.g., refrigeration and freezing) and appropriate thermal processing of foods can effectively prevent foodborne diseases [3].

In the contemporary era, regulatory frameworks and directives pertaining to food safety have been fortified and intensified. Nevertheless, foodborne diseases continue to represent a significant threat to global public health and an economic burden, particularly in developing countries [9]. In its inaugural estimation of the global burden of foodborne diseases in 2015, the WHO attributed 600 million cases of foodborne diseases, 420,000 deaths, and a loss of 33 million years of healthy life worldwide to unsafe food consumption [9].

In 2018, the United States documented 25,606 cases of foodborne infections, resulting in 5,893 hospitalizations and 120 deaths [10]. The burden of foodborne diseases is particularly significant in low- and middle-income countries. Identifying the source of contamination and transmission route is of paramount importance for preventing foodborne illnesses and implementing effective interventions in food safety. However, attributing an infection to specific food and identifying foodborne transmission is challenging and requires source attribution methodologies. Consequently, there is a dearth of studies identifying the sources of foodborne infections, particularly in developing countries [11].

In this study, we aimed to conduct a systematic review of the prevalence of foodborne pathogens in different types of foods in Iran. As a result, we can

gain an overview of the role of food in the transmission of infections and emphasize the importance of food safety in controlling foodborne diseases and reducing their health and economic burden on society.

Materials and Methods

Search strategy

A comprehensive and systematic search was conducted in various databases, including ScienceDirect, Scopus, PubMed, Google Scholar, and local Iranian databases, namely the Iranian Scientific Information Database (www.sid.ir). The literature review was limited to studies published during 2000-2023. The keywords used for searching included "prevalence", "detection", and "identification" in conjunction with terms, such as "food", "Iran", "foodborne pathogen", "food infection", "food poisoning", "food illness", "food disease", "foodborne bacteria", "Campylobacter", "Listeria", "Salmonella", "Helicobacter pylori", "Vibrio", "Clostridium botulinum", "Clostridium difficile", "Clostridium perfringens", "Mycobacterium tuberculosis", "Coxiella burnetii", "Staphylococcus aureus", "Shigella", "Pseudomonas", "Bacillus cereus", "Brucella", and "Yersinia enterocolitica".

Eligibility criteria

This systematic review included articles that focused on the prevalence of foodborne pathogens in any type of food in Iran. Duplicate reports and articles without a clear sample size or other essential data were excluded.

Data extraction

Data collection included extracting information, such as the year of publication, types of foods tested for pathogen contamination, sample size, and number of positive samples contaminated with foodborne pathogens.

Results and Discussion

Results and Discussion

Figure 1 illustrates the study selection process presented in the PRISMA diagram. A systematic literature search using Scopus, ScienceDirect, Google Scholar, SID, Magiran, and cross-references yielded an initial total of 4740 articles. After removing duplicates, 1719 articles remained for title/abstract screening. Following this screening, 655 articles were selected for full-text review. Finally, 328 eligible studies were included in the systematic review.




Figure 1. Flow diagram showing the results of search

1) *Salmonella* spp. prevalence in food

Salmonella (*S.*) *enterica enterica* has more than 2300 serotypes, with *S. Enteritidis* and *S. Typhimurium* being the most commonly reported serotypes. Symptoms of salmonellosis include abdominal pain, vomiting, nausea, diarrhea, and fever [12]. Raw meat, particularly poultry, and egg products, are the main sources of foodborne salmonellosis. Other reported foods that transmit *Salmonella* to humans include fish, peanuts, unpasteurized juice, and milk. It is important to cook raw foods thoroughly to a safe minimum internal temperature to prevent foodborne salmonel-

losis, as *Salmonella* is heat-sensitive. However, processed foods, such as RTE meats and salads can become contaminated through cross-contamination during processing [12]. In Europe in 2020, 0.15% of RTE food samples and 2.4% of non-RTE food samples were positive for *Salmonella* [13].

Table 1 presents the prevalence of *Salmonella* in different foods in Iran based on our review. The highest levels of contamination were found in

poultry meat (23.03%), followed by red meat (14.13%), dairy products (11.66%), RTE foods (11.34%), eggs (9.93%), vegetables (7.8%), fish and shrimp (5.93%), raw milk (3%), and water (2.25%) (Figure 2). In a study conducted in China in 2019, out of 1035 different food samples, a total of 147 samples (14.2%) were positive for *Salmonella*. In their study, the highest prevalence of *Salmonella* was found in fresh meat samples (28%), followed by RTE foods (9%), frozen foods (7.1%), and fresh produce (4.5%) [14]. Fresh meat is a common source of *Salmonella* contamination due to the nature of its production and processing [15]. During the slaughter and processing of animals, there is a high risk of cross-contamination with various bacteria, such as *Salmonella* [16]. In addition, fresh meat products consumed raw or undercooked increase the risk of foodborne illness [17]. The handling and storage of fresh meat products can also contribute to *Salmonella* contamination [18]. In contrast, RTE foods and frozen foods undergo processing and packaging that can reduce the risk of *Salmonella* contamination [19]. However, it is still possible for *Salmonella* to be introduced during the processing or packaging of these products [20]. Fresh produce, while less likely to be contaminated with *Salmonella* compared to fresh meat, can still pose a risk if not properly handled and washed before consumption [18].

2) *Staphylococcus aureus* prevalence in food

Although *Staphylococcus* (*S.*) *aureus* is the primary causative agent of hospital and community-acquired infections, it has also been associated with foodborne diseases. *S. aureus* can cause various gastrointestinal illnesses, which are characterized by nausea, vomiting, abdominal cramps, weakness, and diarrhea [21]. Table




Figure 2. Prevalence of *Salmonella* spp. in different foods in Iran.

Table 1. Summary of the studies reporting the prevalence of *Salmonella spp.* in Iran.

Year	Sample type	Sample size	Positive sam- ples (N)	Prevalence (%)	Type of <i>Salmonella spp.</i>	Referenc- es	City	
1	2023	Eggs	40	4	10	<i>Salmonella</i> <i>Enteritidis</i>	[72]	Qazvin
2	2023	poultry meat	440	39	9	<i>Salmonella</i> <i>enteritidis</i> and <i>Salmonella</i> <i>typhimurium</i>	[73]	Shahrekord
3	2022	Eggs	500	405	81	<i>Salmonella spp.</i>	[74]	Lahijan
4	2022	retail raw meat	60	16	26	<i>Salmonella spp.</i>	[75]	Urmia
5	2022	poultry products	80	11	13.75	<i>Salmonella spp.</i>	[76]	Ardabil
6	2022	Eggs duck	130	21	16.6	<i>Salmonella spp.</i>	[77]	Qazvin
7	2022	Chicken meat	150	4	2.7	<i>Salmonella spp.</i>	[78]	Zahedan
8	2022	cattle raw milk	100	2	2	<i>Salmonella spp.</i>	[79]	Mazandaran
9	2022	Red meat	300	35	11	<i>Salmonella spp.</i>	[80]	Shahrekord
10	2021	Chicken meat	100	6	6	<i>Salmonella spp.</i>	[81]	Ardabil
11	2021	poultry and egg	3125	250	8	<i>Salmonella spp.</i>	[82]	six provinces of Iran
12	2021	chicken meat, beef	450	40	8	<i>Salmonella enterica</i>	[83]	Tehran
13	2019	Olivier salad	26	6	23	<i>Salmonella spp.</i>	[84]	Mashhad
14	2019	Raw chicken meat	60	29	48.3	<i>Salmonella spp.</i>	[85]	Karaj
		Egg yolk	30	0	0			
15	2019	Eggs (shell & contents)	525	0	0	<i>Salmonella spp.</i>	[86]	Isfahan
16	2019	Traditional cheeses	100	0	0	<i>Salmonella spp.</i>	[87]	Mohabad
17	2018	Pastry	75	5	6.6	<i>Salmonella spp.</i>	[88]	Mashhad
18	2018	Chicken Meat	100	7	7	<i>Salmonella spp.</i>	[89]	Mashhad
19	2018	Industrial eggs (shell & contents)	60	0	0	<i>Salmonella spp.</i>	[90]	Zanjan
		Local eggs shell	60	0	0			
		Local egg contents	60	2	1.6			

Table 1 cont.

		Industrial eggs (shell & contents)	34	0	0			
20	2018	Local eggs shell	42	6	4.2	<i>Salmonella spp.</i>	[91]	Sanandaj
		Local egg contents	42	2	4.7			
		Bulk eggs shell	40	3	7.5			
		Bulk egg contents	40	0	0			
21	2018	Industrial eggs (shell & contents)	60	0	0	<i>Salmonella spp.</i>	[92]	Qazvin
22	2018	Shrimp	245	33	13.4	<i>Salmonella spp.</i>	[93]	Khuzestan
23	2017	Pizza	90	8	8.8	<i>S. arizonae</i>	[94]	Ilam
		Frankfurter	90	22	24.4			
		Sausages	90	19	21.1			
24	2017	Camel meat	150	0	0	<i>S. Typhimurium</i>	[95]	Kohgiluyeh & Boyerahmad/ Isfahan
25	2017	Hamburgers	100	2	2	<i>Salmonella spp.</i>	[96]	Kashan
26	2016	beaf meat	190	7	3.6	<i>S. Typhimurium</i>	[97]	Gilan
		Poultry meat	190	15	7.8			
27	2016	Poultry meat	183	52	28.4	<i>S. Typhimurium</i>	[98]	Zanjan
28	2016	Industrial Olovier salad	48	0	0	<i>Salmonella spp.</i>	[99]	Isfahan
		Traditional Olovier salad	54	11	20.4			
29	2016	Traditional ice cream	90	62	68.8	<i>Salmonella spp.</i>	[100]	Zabol
30	2015	Pastry cream	120	0	0	<i>Salmonella spp.</i>	[101]	Arak
31	2015	Olivier salad	50	9	18	<i>S. Ttypimurium</i>	[102]	Shahrekord
32	2015	Poultry meat	625	35	5.6	<i>S. Enteritidis</i>	[103]	Tehran
		Eggs	625	25	4			
33	2015	Eggs	50	5	10	<i>S. Enteritidis</i>	[104]	Shiraz
34	2015	Native eggs	64	0	0	<i>Salmonella spp.</i>	[105]	Yasuj
35	2015	Traditional & Industrial cheese	200	34	17	<i>Salmonella spp.</i>	[106]	Shahrekord
				11	5.5			
36	2015	Eggshells	150	2	1.3	<i>S. Enteritidis & S. Typhimurium</i>	[107]	Tabriz
		Egg contents	150	0	0			

Table 1 cont.

37	2014	Meat Products (Burgers, Sau-sages, Kababs, Cutlets)	42	0	0	<i>Salmonella spp</i>	[108]	Urmia
38	2014	Unpasteurized cream	100	2	2	<i>S. Paratyphi B</i>	[109]	Tehran
39	2014	Chicken	190	86	45	<i>Salmonella spp.</i>	[110]	Tehran
		Beef meat	189	38	20.2	<i>Salmonella spp.</i>		
		Beef meat	189	19	10	<i>S. thompson</i>		
		Chicken	190	65	34.2	<i>S. thompson</i>		
40	2014	Chicken meat	200	58	29	<i>Salmonella spp.</i>	[111]	Alborz
		Liver	120	26	21.6			
		Heart	120	17	14.1			
41	2014	Poultry meat	89	28	31.6	<i>Salmonella spp.</i>	[112]	Shahrekord
		Beef meat	98	26	26.1			
42	2013	Olovier salad	50	9	18	<i>S. Typhimurium</i>	[112]	Shahrekord
43	2013	Tap water	144	5	3.4	<i>Salmonella spp</i>	[113]	Isfahan
		304	304	5	1.1			
44	2013	Poultry slaugh-tered	250	7	2.8	<i>S. infantis</i>	[114]	Brijand
45	2013	Local eggs	210	14	66.6	<i>Salmonella spp.</i>	[115]	Kohgiluyeh & Boyerahmad
46	2013	Industrial egg contents	100	0	0	<i>Salmonella spp.</i>	[116]	Talesh
		Industrial egg-shells	100	19	19			
		Local egg con-tents	100	0	0			
		Local eggshells	100	4	4			
		Local chicken meat	100	21	21			
		Industrial chick-en meat	100	5	5			
		Red meat	150	5	3.3			
47	2013	Industrial Olovieh salad	200	0	0	<i>Salmonella spp.</i>	[117]	Yazd
48	2012	Seafood	384	19	5	<i>Salmonella spp.</i>	[118]	Bushehr, Hormozgan, Khuzestan
49	2012	Chicken meat	150	14	9.3	<i>Salmonella spp.</i>	[119]	Isfahan & Shahrekord
		Turkey meat	105	7	6.7			
		Ostrich meat	45	1	2.2			
50	2012	Beef meat	60	7	11.6	<i>Salmonella spp.</i>	[120]	Sanandaj
				4	6.6	<i>S. Typhimurium</i>		
51	2012	Packed chicken meat	96	19	19.7	<i>Salmonella spp.</i>	[121]	Mazandaran
		Unpacked chicken meat	104	24	23			

Table 1 cont.

52	2012	Salt water fish	70	2	2.9	<i>Salmonella spp.</i>	[122]	Ahvaz
		Shrimp	70	3	4.3			
		Shrimp burge	10	1	10			
53	2011	Raw cow's milk	350	14	4	<i>Salmonella spp.</i>	[123]	Shahrekord
54	2010	Egg	100	0	0	<i>Salmonella spp.</i>	[124]	Shahrekord
55	2010	Chicken meat	190	86	45	<i>Salmonella spp.</i>	[125]	Tehran
		Beef meat	189	38	20			
56	2010	Turkey meat	144	14	9.7	<i>Salmonella spp.</i>	[126]	Isfahan
		Ostrich meat	65	3	4.6			
		Partridge meat	40	0	0			
57	2009	Eggshells	250	4	1.6	<i>S. Typhimurium</i>	[127]	Mashhad
		Egg contents	250	0	0	<i>Salmonella spp.</i>		
58	2009	Poultry car-	60	5	8.3	<i>Salmonella spp.</i>	[128]	Mashhad
		casses	60	1	1.6	<i>S. Typhimurium</i>		
59	2009	Egg contents	120	0	0	<i>Salmonella spp.</i>	[129]	Zanjan
		Eggshells	120	68	56.6			
		Chicken meat	120	104	86.6			
60	2009	Chicken meat	67	32	47.7	<i>Salmonella spp.</i>	[130]	Tehran
		Beef meat	66	19	28.7			
61	2008	Local egg con-	500	1	0.2	<i>Salmonella spp.</i>	[131]	Birjand
		tents						
62	2008	Local eggshells	500	2	0.4	<i>Salmonella spp.</i>	[132]	Isfahan
		Raw poultry	134	24	17.9			
		Cooked poultry	56	3	5.3			
		Turkey	3	1	33.3			
		Quail	5	2	40			
		Red meat	101	8	7.9			
		Cooked meat	118	2	1.6			
		Vegetables	38	3	7.8			
		Fish	15	0	0			
		Yogurt	32	0	0			
		Olovieh salad	20	0	0			
Hamburger	5	0	0					
Mayonnaise souse	8	0	0					
63	2007	Poultry car-	132	92	69	<i>Salmonella spp.</i>	[133]	Tehran
casses								
64	2007	Traditional cheeses	200	0	0	<i>Salmonella spp.</i>	[134]	Jahrom
65	2006	Liver	145	12	8.1	<i>Salmonella spp.</i>	[135]	Yazd
		Meat (before chiller)	145	28	18.4			
		Meat (after chiller)	145	50	34.4			
66	2006	Local eggs	500	3	0.6	<i>Salmonella spp.</i>	[133]	Birjand

Table 2.
Summary of the studies reporting the prevalence of *S. aureus* in Iran.

Year	Sample type	Sample size	Positive samples (N)	Prevalence (%)	City	
1	2023	Poultry meat	94	16	17	Shahrekord
2	2022	Sausages and Bologna	100	31	31	Tehran
3	2022	Raw and ready-to-eat green leafy vegetables	366	134	36.6	Tehran
4	2022	Retail raw meat	60	23	39	Urmia
5	2022	Raw Milk And Traditional Dairy	150	23	15.33	Alborz
6	2022	Ready To Eat Food	320	10	3.12	Tehran
7	2022	Raw Milk	380	42	11.05	Alborz
8	2022	Raw Cow Milk	90	35	38.88	Shahrekord
9	2021	Baghlava	112	3	2.67	Qazvin
10	2021	Raw milk	250	46	18.4	Mashhad
11	2021	Cheese, raw and pasteurized milk	100	10	10	-
12	2019	Meat Products	160	26	16.25	Shahrekord
13	2021	Milk and Cheese	200	23	11	Khuzestan
14	2021	Meat retail	90	31	34.5	Zanjan
15	2021	Ready-to-eat food	415	64	15.42	Tehran
16	2021	Chicken meat	24	6	25	Tehran
17	2019	Zoolbia & Bamieh	75	21	28	Mashhad
18	2019	Fowl meat	240	22	9.6	Tehran
19	2019	Traditional cheese	100	21	21	Maragheh
20	2018	Traditional cheese	100	45	45	Mahabad
21	2017	Raw foods with animal origin	84	20	23.8	Isfahan
		Cooked foods with animal origin	132	12	9	
		Cooked foods without animal origin	269	15	5.7	
22	2017	Season salad	18	0	0	Bandar abbas
		Pasta salad	5	2	40	
		Lettuce	16	0	0	
		Shirazi salad	7	0	0	

Table 2 cont.

23	2017	Pizza	90	11	12.2	Ilam
		Frankfurter	90	25	27.7	
		Sausages	90	22	24.4	
24	2016	Cheese	120	18	41.6	Hamedan
25	2016	Meat	380	78	20.5	Gilan
26	2016	Shrimp	300	84	28	Persian Gulf, Caspian Sea
		Fish	300	122	47	
27	2015	Red meat	379	36	9.4	Hamadan
		Dairy products	671	62	7.2	
28	2015	Raw milk	320	88	27.5	Chaharmahal va Bakhtiari
		Dairy products	350	87	24.8	
29	2015	Shrimp	300	74	24.6	Persian Gulf, Tehran
30	2015	Raw milk	1930	248	12.8	Mazandaran
		Dairy products	720	80	11.1	
31	2015	Bovine milk	92	44	47	Maku
		Sheep milk	86	32	37	
32	2015	Industrial Olivier salad	30	15	50	Shahrekord
		Traditional Olivier salad	20	8	40	
33	2015	Cheese	80	80	100	Marand
34	2015	Chicken nuggets	420	24	5.7	Isfahan
35	2015	Different food	606	12	1.9	Gilan
36	2014	Cream pastry	450	194	43.3	Gorgan
37	2014	Milk	100	9	9	Tabriz
		Cheese	100	45	45	
38	2014	Traditional ice cream	30	2	6.7	Yasuj
		Olover salad	4	0	0	
		Cream suit	30	9	30	
39	2014	Raw milk	300	125	41.6	Ahwaz
40	2014	Dairy product	460	127	27.6	Marand
41	2014	Cheese	80	80	100	Tehran, Gilan
42	2014	Doogh	126	86	68	,Mazandaran

Table 2 cont.

43	2014	Raw milk	120	49	40.8		Kurdistan
44	2014	Meat products	150	19	12.6		Tonekabon
45	2014	Traditional cheeses	100	16	16		East- Azer- bajjan
46	2013	Dairy products	347	20	5.8		Isfahan, Chaharmahal va Bakhtyari, Khuzestan,
47	2013	Industrial Olivier salad	200	40	20	[117]	Yazd
48	2013	Milk	200	22	11	[175]	Fars
49	2012	Traditional white cheese	100	26	26	[176]	Tabriz
		butter	150	24	16		
50	2012	Ground-meat kebab	72	72	100	[177]	Shahrekord
		Bakkhtiyari Kebab	72	72	100		
		Fish	72	72	100		
		Salad	72	72	100		
51	2012	Seafood products	245	22	8.9	[122]	Different mar- kets of Iran
52	2012	Packaged hamburger	256	64	25	[178]	Tehran
53	2012	Raw milk	100	50	50	[176]	Urmia
		Pasteurized milk	100	2	2		
		Ice cream	100	26	26		
54	2012	Raw milk	348	46	13.2	[179]	Shahrekord
55	2010	Fruit juice	360	32	8.8	[180]	Shahrekord
56	2008	Different food	216	30	55.6	[181]	Tehran
57	2006	Fresh fish	67	15	22.3	[182]	Gilan, Caspian Sea

2 presents the findings of studies conducted in Iran regarding the prevalence of this pathogen in different food categories, including seafood (38.51%), meat products (35.47%), dairy products (31.70%), red meat (25.85%), RTE foods (23.59%), raw milk (23.32%), and poultry meat (14.32%) (Figure 3). Seafood and fish are conducive to microbial growth due to their abundant protein and water content. *S. aureus* is not typically found in the natural microflora of fish, there-

fore, its presence can indicate poor personal hygiene, new contamination, or potential disease in the fish [22]. Improper conditions in the fishery, storage, and non-standard transportation provide conditions for pathogens to grow [23]. Furthermore, the hot climate in Iran can facilitate the growth and proliferation of *S. aureus* bacteria in food products, such as meat and dairy items, particularly if they are not stored and refrigerated correctly [24].




Figure 3. Prevalence of *S. aureus* in different foods in Iran.

Table 3. Summary of the studies reporting the prevalence of *Listeria* spp. in Iran.

Year	Sample type	Sample size	Positive samples (N)	Prevalence (%)	Type of <i>Listeria</i> spp.	References	Area	
1	2023	Various Food	900	136	15.1	<i>L. monocytogenes</i>	[183]	Mazandaran and Golestan
2	2022	Retail raw meat	60	25	42	<i>L. monocytogenes</i>	[75]	Urmia
3	2022	Curd and cheese	150	14	9.33	<i>L. monocytogenes</i>	[184]	-
4	2021	Non-Pasteurized Milk	50	30	60	<i>L. monocytogenes</i>	[185]	Kerman
5	2021	Traditional cheeses	60	1	1.6	<i>L. monocytogenes</i>	[186]	Tehran
6	2021	Raw Milk	100	10	10	<i>Listeria</i> spp.	[187]	Tehran
7	2021	Seafood	350	40	11.42	<i>Listeria</i> spp.	[188]	Genaveh port
8	2020	Beef and chicken meat	90	45	50	<i>L. monocytogenes</i>	[189]	Zanjan
9	2019	Eggs	525	0	0	<i>Listeria</i> spp.	[86]	Isfahan
10	2019	Chicken meat retailers	811	257	30.5	<i>Listeria</i> spp.	[190]	Mashhad
11	2018	Traditional dairy products	545	64	11.7	<i>Listeria</i> spp.	[191]	Yazd
			22	4.3	<i>L. monocytogenes</i>			
12	2017	Food (sausage, milk, cheese, chicken and meat)	267	8	2.9	<i>Listeria</i> spp.	[192]	Urmia
13	2017	Fresh chicken carcasses	200	80	40	<i>Listeria</i> spp.	[193]	Mashhad
14	2016	Dairy products	107	9	8.4	<i>L. monocytogenes</i>	[194]	Tehran
		Processed meat	210	11	5.2	Karaj &		
15	2016	Seafood	237	7	2.9	<i>L. monocytogenes</i>	[195]	Tehran

Table 3 cont.

16	2016	Argyrosomus	240	30	12.5	<i>Listeria spp.</i>	[196]	Isfahan & Bandaranzali	
		hololepidotus		5	16.6	<i>Bandaranzali</i>			
17	2015	Koozeh cheeses	100	3	3	<i>L. monocytogenes</i>	[197]	Urmia	
18	2015	Minced beef	150	4	2.7	<i>Listeria spp.</i>	[198]	Ahvaz	
				1	0.6	<i>Mazandaran</i>			
19	2015	Raw fish	488	104	21.3	<i>Listeria spp.</i>	[199]	Mazandaran	
20	2015	Raw milk	60	0	0	<i>L. monocytogene</i>	[200]	Zanjan	
21	2015	Traditional dairy products	292	21	19.7	<i>Listeria spp.</i>	[201]	Isfahan	
22	2015	Raw milk	100	5	5	<i>L. monocytogene</i>	[202]	Kerman	
23	2014	Ready to eat food (olovier salad, Yogurt stew, macaroni salad and meat salad)	235	20	8.5	<i>Listeria spp.</i>	[203]	Shahrekord	
24	2014	Meat products	98	12	32.4	<i>L. monocytogene</i>	[204]	Qazvin	
		Milk products		84	25	29.7			<i>Fars & Khuzestan</i>
25	2014	Bulk milk	260	27	10.4	<i>Listeria spp.</i>	[205]	Fars & Khuzestan	
				7	2.7	<i>Bandar anzali</i>			
26	2014	Smoked fish	80	7	8.8	<i>Listeria spp.</i>	[206]	Isfahan & Bandar anzali	
				2	2.5	<i>L. monocytogene</i>			
				6	15	<i>Listeria spp.</i>			
27	2013	Meat and meat products	60	8	13.3	<i>Listeria spp.</i>	[207]	Khoramabad & Tehran	
				2	6.6	<i>L. monocytogene</i>			
28	2013	Crayfish meat	40	3	7.5	<i>L. monocytogene</i>	[208]	Aras	
29	2013	Raw cow milk	986	25	2.5	<i>Listeria spp.</i>	[209]	Isfahan	
				20	2	<i>L. monocytogene</i>			
30	2013	Vegetables and ready mayonnaise salads	300	26	8.7	<i>Listeria spp.</i>	[210]	Tehran	
				21	7	<i>L. monocytogene</i>			
31	2013	Raw seafood products	331	16	4.8	<i>L. monocytogene</i>	[211]	Shahrekord	
		RTE seafoods		321	46	14.5			<i>L. monocytogene</i>
32	2013	Raw milk	466	83	18.6	<i>Listeria spp.</i>	[212]	Tehran	
33	2013	Dairy products	185	7	3.8	<i>Listeria spp.</i>	[213]	Kermanshah	
		Meat products		187	51				27.2
		Ready-to-eat foods		158	8				5.1
34	2013	Seafood	300	24	8	<i>Listeria spp.</i>	[214]	Isfahan & Shahrekord	
				18	6	<i>L. monocytogene</i>			
35	2013	Quail products	150	10	6.6	<i>Listeria spp.</i>	[215]	Isfahan	
				1	0.6	<i>L. monocytogene</i>			
36	2013	Lamb	200	5	2.5	<i>L. ivanovii</i>	[216]	Shahrekord	
37	2012	Different types of raw meat	1107	141	12.7	<i>Listeria spp.</i>	[217]	Shahrekord, Isfahan, Ahvaz, Shiraz, Yazd,	
				27	2.4				
38	2012	Poultry product	402	134	33.3	<i>Listeria spp.</i>	[218]	Shahrekord	
39	2012	Seafood	264	20	7.6	<i>Listeria spp.</i>	[219]	Isfahan & Shahrekord	

Table 3 cont.

40	2012	Various seafood products	245	2	0.8	<i>L. monocytogene</i>	[122]	Different markets of Iran				
41	2011	Eggs	100	0	0	<i>L. monocytogene</i>	[124]	Shahrekord				
42	2011	Fish	194	24	12.3	<i>Listeria spp.</i>	[220]	Urmia				
				5	2.5	<i>L. monocytogene</i>						
43		Raw cow milk	45	5	1.1	<i>Listeria spp.</i>	[221]	Shiraz				
				2	4.4	<i>L. monocytogene</i>						
		Raw goat milk		1	3.1	<i>Listeria spp.</i>						
				1	3.1	<i>L. monocytogene</i>						
		Traditional cheese		41	10	24.4			<i>Listeria spp.</i>			
					4	9.7			<i>L. monocytogene</i>			
Traditional ice-cream	60	8	11.7	<i>Listeria spp.</i>								
			2	3.3	<i>L. monocytogene</i>							
44		Raw milk	100	4	4	<i>L. monocytogene</i>	[222]					
45		Dairy products	360	6	1.6	<i>L. monocytogene</i>	[223]					
46		Chilled ready to eat foods	41	3	7.3	<i>L. monocytogene</i>	[224]					
		Meat, meat products							332	4	1.2	<i>L. monocytogene</i>
		Milk and dairy products							88	0	0	<i>L. monocytogene</i>
47		Cattle carcasses	203	6	3	<i>L. monocytogene</i>	[225]					

3) *Listeria monocytogenes* prevalence in food

Listeria (L.) *monocytogenes* represents a significant public health concern due to its ability to be transmitted from the environment to food, which can lead to foodborne listeriosis in humans [25]. In 2020, the EFSA reported a total of 1876 cases of listeriosis, with 97.1% of these cases necessitating hospitalization [26]. Moreover, the EFSA indicated an increase in the case fatality rate and hospitalization rate associated

with *L. monocytogenes* infections in 2020. Among all the reported zoonoses in Europe in 2020, listeriosis had the highest case fatality rate of 13% [26]. Those at the greatest risk of developing listeriosis include pregnant women, the elderly, newborns, and patients with compromised immune systems [27]. Moreover, a multitude of food items were identified as potential sources of listeriosis outbreaks during this period. Specifically, 4.8% of RTE meat products and 0.44% of




Figure 4. Prevalence of *Listeria* in different foods in Iran.

Table 4
Summary of the studies reporting the prevalence of *C. burnetii* in Iran..

Year	Sample type	Sample size	Positive samples (N)	Prevalence (%)	References	Area	
1	2022	Unpasteurized Camel Milk	100	6	6	[226]	Mashhad
2	2022	Kope cheese and cattle milk	800	103	12.87	[227]	West Azerbaijan
3	2021	Raw Milk	162	23	14	[228]	Tehran, Hamadan, and Mazandaran
4	2021	Raw milk	100	27	27	[229]	Mazandaran
5	2020	Raw milk	204	21	10.2	[230]	Gilan
6	2020	Sheep and goats milk	420	51	12.1	[231]	West Azerbaijan
7	2019	Milk	126	44	34.9	[232]	Qom
8	2019	Milk	840	14	16.9	[233]	West Azerbaijan
		Cattle milk			14.4		
		Buffalo milk			19.3		
9	2019	Cream & butter	200	6	3	[234]	Shahrekord
		Traditional bovine cream,	69	4	5.7		
		Traditional sheep butter,	20	1	5		
		Traditional bovine butter	39	1	2.5		
10	2018	Raw milk	500	9	1.8	[235]	Khorramabad
		Sheep milk		3			
		Goat milk		6			
11	2018	Bulk milk	100	3	3	[236]	Shiraz
12	2018	Non-pasteurized dairy products	238	20	8.4	[29]	Shiraz
13	2018	Raw milk	100	10	10	[237]	Tehran
		Traditional unpacked cheese	40	3	7.5		
14	2016	Raw sheep milk	72	15	20.8	[238]	Khorramabad
15	2015	Bovine bulk milk	70	12	17.4	[239]	Jahrom
				7	10		
				7	10		
16	2015	Individual raw milk	60	7	11.6	[200]	Zanjan
		Bovine	38	5	8.3		
		Ovine	22	2	3.3		
17	2015	Cow milk	150	18	12	[240]	Tehran
18	2015	Goat milk	31	5	16.1	[241]	Kerman
19	2015	Cow milk	80	20	25	[242]	Ajabshir
20	2014	Cheese	28	2	7.1	[243]	Mashhad
		Yoghurt	26	2	7.6		
		Sheep milk	23	8	34.7		
		Cow milk	60	2	3.3		
21	2014	Bovine bulk tank	100	5	5	[244]	Mashhad
22	2014	milk	51	21	41.1	[245]	Khoramabad.
23	2013	Goat milk	100	14	14	[246]	Qom

Table 4 cont.

24	2013	Bovine bulk milk	100	11	11	[247]	Jahrom
		Bovine milk	247	8	3.2		
25	2011	Ovine bulk milk	140	8	5.7	[248]	Kerman
		Caprine bulk milk					
		Camel bulk milk	110	5	4.5		
26	2010	Bulk milk	296	6	2	[249]	Fars, Ghom, Kerman, Yazd Khuzestan
		Cow milk	210	13	6.2		
27	2010	Sheep milk	110	0	0	[250]	Chaharmahal va Bakhtiari
		Goat milk	56	1	1.8		

milk and milk products were found to be contaminated with *L. monocytogenes* [26]. Table 3 and Figure 4 present the findings of studies conducted in Iran regarding the prevalence of *L. monocytogenes* in various food types. As illustrated in Figure 4, poultry meat exhibited the highest contamination rate of 38.45%, followed by meat products (14.94%), red meat (13.45%), raw milk (9.77%), dairy products (9.48%), seafood (8.75%), and RTE foods (7.32%) (Figure 4). A previous review study conducted in Iran until 2015 yielded comparable results regarding the contamination of food with *Listeria*. The highest prevalence of *L. monocytogenes* was approximately 9.2%, which was observed in RTE foods [25]. Therefore, RTE foods should be considered a potential hazard to consumers [25]. Similarly, other developing countries have also yielded comparable results. For example, a study conducted in Ethiopia revealed that 28.4% of raw milk and milk products were contaminated with *Listeria* spp., with 5.6% of these samples testing positive for *L. monocytogenes* [23].

4) *Coxiella burnetii* prevalence in food

Coxiella burnetii is a zoonotic pathogen that causes Q fever in humans and coxiellosis in livestock. Cattle, goats, and sheep serve as the primary reservoirs for the pathogen, facilitating its transmission to humans [28]. The primary routes of human infection are through the inhalation of contaminated aerosols or the consumption of unpasteurized milk and dairy products [29]. In Europe, 523 cases of Q fever were identified in 2020, resulting in a case fatality rate of 2.1% [30]. Table 4 presents the results of studies conducted in Iran concerning the prevalence of *C. burnetii* in different food items. As illustrated in Figure 5, the foods with the highest contamination rates were raw milk (12.36%) and dairy products (6.40%). *C. burnetii* is a bacterium that causes Q fever, a zoonotic disease that can be transmitted from animals to humans. In numerous rural regions of Iran, milk is still produced and processed using traditional methods that fail to meet the requisite modern hygiene standards [31]. The absence of adequate hygiene protocols in milk




Figure 5. Prevalence of *C. burnetii* in different foods in Iran.

production and processing facilities may result in the contamination of milk with *C. burnetii*. Moreover, the proximity of animals to humans in the rural areas of Iran contributes to the high levels of contamination of raw milk and dairy products with *C. burnetii* [32]. Animals, such as cows and goats, can carry the bacterium and shed it in their milk, which can then be transmitted to humans through consuming contaminated dairy products [33].

A study conducted in Italy in 2017 reported that 15% of milk samples were contaminated with *C. burnetii*, with a higher prevalence of contamination in bovine milk (41%) compared to sheep milk (12%) [34]. In Brazil, in 2020, 9.43% of cheese samples (out of 53 samples) were positive for *C. burnetii* DNA [35]. Another research in the United States reported that 94% of bulk milk samples from dairy herds were contaminated with *C. burnetii* [36]. Our review indicates that the data from Iran align with the reports from other countries. However, it should be noted that the prevalence of *C. burnetii* contamination varies depending on the type of dairy products, including specific variations within milk.

5) *Bacillus cereus* prevalence in food

Bacillus cereus spores are a well-documented contaminant of food that can survive high temperatures during cooking and pasteurization [37]. This bacterium is associated with two distinct types of gastrointestinal diseases: the emetic (vomiting) syndrome and the diarrheal syndrome [38]. In Europe, 835 cases of foodborne illness caused by *B. cereus* were reported

in 2020, with a hospitalization rate of 1.2% and a mortality rate of 0.1% [30]. The diarrheal syndrome is typically attributed to the consumption of contaminated foods, including raw and cooked beef, meat products, fish, poultry, soups, sauces, stews, milk, and vegetables. In contrast, the emetic syndrome is associated with the consumption of a toxic dose of the pre-formed emetic (cereulide) toxin produced by *B. cereus* in starchy foods, such as rice, pasta, noodles, potatoes, bread, pastries, and sesame products [39]. Table 5 presents the results of studies conducted in Iran regarding the prevalence of *B. cereus* in different food items. As illustrated in Figure 6, the highest prevalence of *B. cereus* contamination was observed in rice (100%), followed by raw milk (48.8%), poultry meat (42.17%), spices (42%), infant food (32.62%), dried vegetables (31.42%), meat products (11.16%), red meat (9.33%), and dairy products (8.9%) (Figure 6). In Australia, *B. cereus* contamination was identified in a variety of food samples, including uncooked pizza bases (1.58%), cooked pizzas (4.57%), processed meats (0.28%), cooked meat pies (4.45%), cooked sausage rolls (3.26%), and raw diced chicken (5.45%) out of 1,263 retail food samples [40]. In China, *B. cereus* contamination was observed in 50% of rice and noodle samples, 34% of cooked meat samples, and 22% of cold vegetable dishes [41]. In Poland, the highest prevalence of *B. cereus* contamination was found in herbs and spices, with a rate of 63.3%. Moreover, other food items, including breakfast cereals, pasta, rice, pasteurized milk, infant formulas, as well as fresh and ripening cheeses, were also found to be contaminated with *B. cereus* [37].




Figure 6. Prevalence of *B. cereus* in different foods in Iran.

Table 5.
Summary of the studies reporting the prevalence of *B. cereus* in Iran.

Year	Sample type	Sample size	Positive samples (N)	Prevalence (%)	References	Area	
1	2023	200	84	42	[251]	Isfahan	
	Various Spices						
	Individual meat						
2	2020	60	7	3.5	[252]	Zanjan	
							Raw lamb
							Raw beef
3	2020	150	16	10.6	[253]	Tabriz	
							Traditional dairy products
4	2019	10	10	100	[254]	Zanjan	
5	2018	120	13	10.8	[255]	Tabriz	
6	2018	140	44	31.4	[256]	Tehran	
7	2018	62	0	0	[257]	Zanjan	
8	2017	80	18	22.5	[258]	Tehran	
9	2017	300	9	3	[259]		
10	2017	125	84	67.2	[260]		
11	2017	42	41	97.6	[261]		
12	2016	200	10	5	[262]		
13	2016	380	44	11.8	[97]		
14	2016	230	46		[263]		
15	2015	104	80	76.5	[264]	Tehran	
							Poultry meat foods Frozen
							Semi cooked
16	2014	408	408	100	[265]	Urmia	
							Rice
17	2013	200	84	42	[266]	Isfahan	
18	2012	32	9	28	[267]	Tehran	
							Kefir type drinks
19	2007	60	11	18.3	[268]	Tehran	
							Pasteurized milk

6) *Yersinia enterocolitica* prevalence in food

In Europe, 236 cases of foodborne yersiniosis were reported in 2020, with 4.7% of cases necessitating hospitalization [30]. *Yersinia enterocolitica* contamination has been documented in a variety of foods in Europe, including red meat (beef, pork, and lamb), poultry, seafood, eggs, milk and milk products, bean sprouts, vegetables, tofu, and stewed mushrooms [42]. Table 6 presents the results of studies conducted in Iran regarding the prevalence of *Y. enterocolitica* in different food items. As illustrated in Figure 7, poultry meat exhibited the highest contamination rate of 16.81% in Iran. This was followed by raw milk (11.93%), red

meat (11.63%), and dairy products (10%) (Figure 7). In Europe, 5.2% of RTE meat was found to be positive for *Yersinia* in 2020, which is a relatively high and concerning rate [30]. A study conducted in Argentina in 2019 reported chicken (12.4%) and bovine-originated foods (10.2%) as the most contaminated foods with *Y. enterocolitica* [43], which aligns with the findings in Iran. However, the latter study reported a lower prevalence of contamination in dairy products (0.7%) compared to the findings in Iran [43]. The elevated contamination rates of *Y. enterocolitica* in poultry meat observed in Iran and Argentina can be attributed to several factors, including the hygiene practices employed during the processing, transportation, and

Table 6. Summary of the studies reporting the prevalence of *Y. enterocolitica* in Iran.

Year	Sample type	Sample size	Positive samples (N)	Prevalence (%)	References	Area	
1	2022	Raw Milk	360	3	0.83	[269]	Tehran
2	2021	red meat	200	26	13	[270]	Shiraz
3	2021	Bovine Raw Milk	100	33	33	[271]	Mashhad
4	2021	Traditional Cheeses	200	38	19	[272]	Khorasan Razavi and Golestan
5	2020	Raw milk	360	3	0.8	(Soltan Dallal, 2020)	Tehran
6	2019	Cheeses	200	38	19	[273]	Khorasan Razavi and Golestan
		Raw milk	100	33	33		
7	2018	Chicken meat	100	25	25	[274]	Mashhad
8	2018	Raw milk (sheep & goats)	100	9	9	[275]	Shahrekod
9	2018	Turkey meat	300	55	18.3	[276]	Shahrekord
		Meat	450	56	12.4		
10	2015	Chicken meat	226	35	15.4	[277]	Tehran
		Beef meat	224	21	9.3		
11	2015	Raw milk	446	19	4.3	[278]	Varamin
12	2014	Dairy products	552	28	5	[279]	Isfahan
13	2014	Raw chicken meat	300	65	21.6	[280]	Shahrekod
14	2014	Unpasteurized cream	100	3	3	[281]	Tehran
15	2013	Chicken meat	720	132	18.3	[282]	Shahrekod
16	2012	Bulk raw milk	354	8	2.6	[283]	Eastern Azerbaijan
		Cheeses	200	8	4		
17	2012	Chicken meat	200	18	9	[121]	Mazandaran
18	2012	Broiler meat	120	19	15.8	[284]	Tabriz
19	2011	Beef and chicken meat	379	48	12.6	[285]	Tehran




Figure 7. Prevalence of *Y. enterocolitica* in different foods in Iran.

storage of these products [44]. Poultry meat has been identified as a significant source of *Y. enterocolitica* contamination due to the presence of the bacterium in the intestinal tracts of birds [45]. Inappropriate handling and processing of poultry can result in the cross-contamination of the meat with *Y. enterocolitica*. In addition, raw milk, red meat, and dairy products can serve as reservoirs for *Y. enterocolitica* if not properly pasteurized or handled [46].

7) *Campylobacter* prevalence in food

Campylobacter spp. has been identified as the leading cause of foodborne gastroenteritis in Europe since 2005 [30]. In addition to acute gastroenteritis, *Campylobacter*

infections can also result in chronic manifestations in humans [47]. Among the various species within the genus *Campylobacter*, *C. jejuni* and *C. coli* are the most commonly reported causes of *Campylobacteriosis* in humans [48]. Table 7 presents the results of studies conducted in Iran regarding the prevalence of *Campylobacter* in different food items. As illustrated in Figure 8, the most prevalent occurrence of *Campylobacter* contamination in Iran was observed in poultry meat (46.21%), followed by red meat (40%) and eggs (28.06%). The contamination of dairy products and raw milk was observed in 2.36% and 2.5% of samples, respectively (Figure 8). A study conducted in

Table 7. Summary of the studies reporting the prevalence of *Campylobacter spp.* in Iran.

Year	Sample type	Sample size	Positive samples (N)	Prevalence (%)	<i>Campylobacter spp.</i>	References	Area
1	2023	chicken meat	100	81	81	<i>Campylobacter spp.</i>	[286] Hamedan
2	2023	chicken meat	255	64	25.09	<i>Campylobacter spp.</i>	[287] Shahrekord
3	2023	Mushrooms	740	74	10	<i>Campylobacter spp.</i>	[288] -
4	2022	poultry meat	380	24	6.25	<i>Campylobacter spp.</i>	[289] Shahrekord
5	2022	poultry meat	100	35	35	<i>Campylobacter spp.</i>	[290] Tehran
6	2022	raw meat	200	27	13.5	<i>Campylobacter jejuni</i>	[291] -
7	2022	cattle raw milk	100	7	7	<i>Campylobacter jejuni</i>	[292] Mazandaran
8	2021	Poultry Carcasses	370	203	54.8	<i>Campylobacter spp.</i>	[293] south of Iran
9	2021	Camels meat	40	5	12.5	<i>Campylobacter spp.</i>	[294] Chaharmahal and Bakhtiari
10	2019	poultry meat	328	217	66.7	<i>Campylobacter spp.</i>	[295] Jahrom
11	2019	Industrial chicken meat	50	1	0.6	<i>Campylobacter spp.</i>	[295] Ahvaz
		meat		0	0		
		Traditional chicken meat		8	16	<i>C. jejuni</i>	
		meat		3	37.5	<i>Campylobacter spp.</i>	
		Fresh packed chicken meat		0	0	<i>C. jejuni</i>	
		meat		7	14	<i>Campylobacter spp.</i>	
Beef meat	7	100	<i>C. jejuni</i>				
Mutton meat	12	24					
		12	100				
		Water buffalo meat	50	4	8	<i>Campylobacter spp.</i>	
				4	100	<i>C. jejuni</i>	
12	2019	Packed chicken meat		26	28.9	<i>Campylobacter spp.</i>	[296] Shiraz

Table 7 cont.

		Red meat	90	21 13	23 61.9	<i>Campylobacter spp.</i>		
13	2016	Chicken-meat	120	33 22	27.5 66.6	<i>C. jejuni</i> <i>Campylobacter spp.</i>	[297]	Zanjan
		Eggshells	120	38 20	31.6 52.6	<i>C. jejuni</i> <i>Campylobacter spp.</i>		
14	2015	Chicken wing	96	37	38.5	<i>Campylobacter spp.</i>	[298]	Urmia
15	2015	Meat	360	227 200	63.1 88.1	<i>Campylobacter spp.</i> <i>C. jejuni</i>	[299]	Mashhad
16	2015	Raw ovine milk	38	0	0	<i>C. jejuni</i>	[300]	Zanjan
		Raw bovine milk	22	0	0	<i>C. jejuni</i>		
17	2014	Chicken	250	110 87	44 79	<i>Campylobacter spp.</i>	[300]	Tehran
		Pasteurized milk	30	0	0			
		Camel milk	37	0	0	<i>Campylobacter spp.</i>		
		Commercial dairy	290	0	0			
18	2013	Raw cow milk	80	5	6.2	<i>Campylobacter spp.</i>	[301]	Isfahan & Chaharmahal va Bakhtyari
		Raw sheep milk	60	1	1.6	<i>Campylobacter spp.</i>		
		Raw goat milk	60	2	3.3	<i>Campylobacter spp.</i>		
		Traditional cheese	60	3	5	<i>Campylobacter spp.</i>		
		Traditional ice-cream	35	1	2.8	<i>Campylobacter spp.</i>		
		Traditional butter	25	1	4	<i>Campylobacter spp.</i>		
19	2012	Packed chicken meat	96	22	22.9	<i>Campylobacter spp.</i>	[121]	Mazandaran
		Unpacked chicken meat	104	31	28.8	<i>Campylobacter spp.</i>		
20	2011	Eggs	100	0	0	<i>C. jejuni</i>	[124]	Shahrekord
21	2011	Raw bovine milk	120	3	2.5	<i>Campylobacter spp.</i>	[302]	Isfahan
		Chicken	200	94 91	47 96.8	<i>Campylobacter spp.</i> <i>C. jejuni</i>		
		Turkey	49	49 41	49 83.7	<i>Campylobacter spp.</i> <i>C. jejuni</i>		
22	2011	Quail	33	37 33	43 89.2	<i>Campylobacter spp.</i> <i>C. jejuni</i>	[303]	Shahrekord
		Partridge	6	6 6	35.3 100	<i>Campylobacter spp.</i> <i>C. jejuni</i>		
		Ostrich	21	1 1	4.8 100	<i>Campylobacter spp.</i> <i>C. jejuni</i>		
		Chicken meat	60	37 35	61.7 94.6	<i>Campylobacter spp.</i> <i>C. jejuni</i>		
23	2010	Turkey meat	50	18 15	83.3 36	<i>Campylobacter spp.</i> <i>C. jejuni</i>	[304]	Ahvaz
		Sheep meat	50	3 1	6 33.3	<i>Campylobacter spp.</i> <i>C. jejuni</i>		
		Goat meat	45	17 17	4.4 100	<i>Campylobacter spp.</i> <i>C. jejuni</i>		

Table 7 cont.

24	2010	Raw camel meat	107	1	0.9	<i>Campylobacter spp.</i>	[305]	Isfahan & Yazd
				0	0	<i>C. jejuni</i>		
		Beef meat	190	5	2.4	<i>Campylobacter spp.</i>		
				3	60	<i>C. jejuni</i>		
		Lamb meat	225	27	12	<i>Campylobacter spp.</i>		
				23	92	<i>C. jejuni</i>		
Goat meat	180	17	9.4	<i>Campylobacter spp.</i>				
		16	94.1	<i>C. jejuni</i>				
25	2008	Raw chicken meat	280	157	56.1	<i>Campylobacter spp.</i>	[306]	Isfahan
				140	89.2	<i>C. jejuni</i>		
		Quail meat	248	68	27.4	<i>Campylobacter spp.</i>		
				53	77.9	<i>C. jejuni</i>		
		Turkey meat	212	145	68.4	<i>Campylobacter spp.</i>		
				92	63.4	<i>C. jejuni</i>		
		Ostrich meat	60	7	11.7	<i>Campylobacter spp.</i>		
				3	42.9	<i>C. jejuni</i>		

the United States in 2020 reported that while various broiler products carry the risk of *Campylobacter spp.* contamination, the highest prevalence of contamination was observed in chicken carcasses [49]. Similarly, in the European Union, *C. jejuni* has been identified as the most prevalent species (51%) in broiler meat, followed by *C. coli* (35.5%) [47]. Consequently, poultry meat represents the greatest risk of *Campylobacter* transmission to humans worldwide. The consistent reporting of the highest prevalence of *Campylobacter* contamination in poultry meat in multiple studies, including those conducted in Iran, the United States, and the European Union, underscores the importance

of addressing this issue [47, 49]. This finding highlights the necessity of implementing rigorous food safety measures and regulations in the poultry industry to prevent the transmission of *Campylobacter* to consumers.

8) *Helicobacter pylori* prevalence in food

Helicobacter pylori is associated with several digestive diseases, including peptic ulcer, mucosa-associated lymphoid tissue lymphoma, gastritis, and an increased risk of gastric cancer [50]. It is estimated that approximately 50% of the global population is infected with *H. pylori* [51]. The prevalence of *H. pylori*




Figure 8. Prevalence of *Campylobacter* in different foods in Iran.

infection is observed to be higher in developing countries, with rates ranging from 70% to 90%, compared to developed countries, where rates are reported to be 25%-50%. Iran is considered a high-risk region for *H. pylori* infection due to the high prevalence (60%-90%)

among its population [52]. *H. pylori* can be found in a variety of animal-derived foods, vegetables, and water sources, which contribute to its transmission [50]. Table 8 presents the findings of studies conducted in Iran regarding the prevalence of *H. pylori*

Table 8. Summary of the studies reporting the prevalence of *H. pylori* in Iran.

Year	Sample type	Sample size	Positive samples (N)	Prevalence (%)	References	Area	
1	2023	Raw Poultry Meat	320	20	6.25	[307]	Shahrekord
2	2020	Red meat	600	52	8.6	[308]	Tehran
3	2018	Traditional dairy products	800	31	3.8	[309]	Isfahan
4	2017	Red meat	220	11	5	[310]	Isfahan
5	2017	Meat	150	11	7.3	[311]	Alborz
		Milk	150	24	16		
		Vegetable	40	5	12.5		
		Cream-candy	50	9	18		
		Traditional bread	50	3	6		
		Sausage	50	0	0		
		Salami	50	0	0		
6	2016	Hamburger	50	1	2	[312]	Isfahan
		Soup	50	11	22		
		Restaurant salad	50	15	30		
		Falafel	50	3	6		
		Olivier salad	50	18	36		
		Chicken nugget	50	0	0		
		Fruit salad	50	14	28		
7	2016	Milk	420	92	21.9	[313]	Shahrekord
		Meat	400	105	26.2		
8	2016	Meat products	150	11	7.3	[314]	Isfahan
			120	5	4.2		
			110	2	1.8		
9	2016	Drinking water (total)	100	3	3	[315]	Isfahan
							Shiraz
							Yazd
							Shahrekord
10	2016	Ready to eat fish	70	2	2.8	[316]	Shiraz
		Ham	60	9	15		
		Chicken sandwich	60	5	8.3		
		Vegetable sandwich	40	2			
		Meat sandwich	40	18	5		
		Minced meat	50	10	45		
		Minced meat	50	16	20		
11	2015	Raw milk	210	28	13.3	[317]	Tehran
		Bovine milk	120	20	16.6		
		Traditional cheese	80	10	[318]		
12	2015	Traditional cream	40	3	7.5	[318]	
		Total	240	33	[319]		

Table 8 cont.

13	2014	Vegetable and salad	460	44	9.5	[319]	Shahrekord
14	2014	Vegetable and salad	430	59	13.7	[320]	Isfahan
15	2013	Water	200	14	7.2	[51]	Isfahan
16	2012	Milk	447	56	12.5	[321]	Isfahan, Fars, Chaharmahal & Bakhtiari, Khuzestan

in various food items. As illustrated in Figure 9, the highest prevalence of *H. pylori* in food samples in Iran was observed in RTE foods (25.5%) and vegetables (22.14%), followed by raw milk (16.06%), red meat (15.82%), dairy products (7.93%), meat products (6.26%), and water (3.8%) (Figure 9). In other countries, studies have also identified the presence of *H. pylori* in a variety of food sources. In Japan, the ureA gene of *H. pylori* was found in 72.2% of raw milk samples and 55% of pasteurized milk samples [53]. In Italy, the glmM gene of *H. pylori* was identified in 34.7% of raw milk samples [54]. In the United States, *H. pylori* was detected in 44% of RTE raw tuna meat and 36% of raw chickens using a multiplex PCR assay [55]. These findings underscore the potential presence of *H. pylori* in various food sources and the significance of food as a potential route of transmission.

9) Clostridium prevalence in food

Clostridium botulinum

Clostridium botulinum is a gram-positive, anaerobic bacterium that is capable of producing spores. It is known to cause botulism, a severe illness characterized by the production of a potent neurotoxin. Table 9 presents the findings of research conducted in Iran on the prevalence of *C. botulinum* in various food items. As illustrated in Figure 10, the most prevalent contamina-

tion of *C. botulinum* in Iran was observed in seafood (12.56%), followed by red meat (12.23%), dairy products (9.02%), and honey (2%) (Figure 10). Honey is recognized as a reservoir for *C. botulinum* spores, particularly types B and A, and has been implicated in cases of neonatal botulism [30]. Studies conducted in various countries, including Turkey, Brazil, Denmark, Sweden, and Norway, have demonstrated the presence of *C. botulinum* spores in honey samples, with prevalence rates ranging from 2% to 26% [30]. In Iran, the prevalence of *C. botulinum* contamination in honey samples was reported to be 2% (Figure 10), indicating a relatively lower level of contamination compared to some other regions.

While *C. botulinum* spores may be present in certain foods, the risk of botulism is contingent upon the conditions that facilitate the germination of spores and toxin production, such as inadequate food processing, storage, or handling. Proper food safety practices, including adequate cooking, storage at appropriate temperatures, and hygienic handling, can help prevent the growth and toxin production of *C. botulinum* in food.

Clostridium perfringens

C. perfringens is a significant contributor to foodborne gastrointestinal illnesses in both humans and animals. The spores of *C. perfringens* exhibit remarkable resilience to external influences. In Europe in 2020, there




Figure 9. Prevalence of *H. pylori* in different foods in Iran.




Figure 10.
Prevalence of *Clostridium* in different foods in Iran.

were 682 reported cases of food poisoning caused by *C. perfringens* toxins, with a hospitalization rate of 1.5%. Conversely, there were fewer cases (n = 34) of food poisoning due to *C. botulinum* toxins, yet the hospitalization rate for botulism cases was 100%. It is noteworthy that no fatalities were reported in these cases. Early diagnosis, hospitalization, and treatment are essential for reducing the severity of botulism [30]. Table 9 presents the findings of studies conducted in Iran regarding the prevalence of *C. perfringens* in various food items. *C. perfringens* type A is the most prevalent cause of food poisoning associated with this bacterium. The available data indicate that

C. perfringens was most commonly isolated from red meat in Iran. It is of paramount importance to ensure that meat is cooked and handled properly to minimize the risk of contamination with *C. perfringens* and subsequent foodborne illnesses. In Europe in 2019, two outbreaks were associated with pig meat and products, one caused by toxins produced by *C. perfringens* and the other by *C. botulinum*. Conversely, vegetables, juices, and other related products were linked to a greater number of outbreaks, with two outbreaks reported for each category during the same period [30]. Nevertheless, only one study has been conducted in Iran regarding the presence of *C. perfringens* in vege-

Table 9.
Summary of the studies reporting the prevalence of *Clostridium* spp. in Iran.

Year	Sample type	Sample size	Positive samples (N)	Prevalence (%)	Clostridium spp.	References	Area
1	2023	Meat native birds	300	35	11.6	<i>C. perfringens</i>	[322] Shahrekord
2	2023	Meat Nuggets	600	7	1.17	<i>C. perfringens</i>	[323] Isfahan
3	2023	Ground Beef	133	24	18.04	<i>C. perfringens</i>	[324] Qazvin
			94	3	3.22		
4	2022	Raw Meat	240	7	2.91	<i>C. perfringens</i>	[325] Mazandaran
5	2022	Raw Beef Meats	133	18	13.53	<i>C. perfringens</i>	[326] Qazvin
6	2022	raw and ready-to-eat green leafy vegetables	366	66	18	<i>C. perfringens</i>	[139] Tehran
7	2022	Olivier Salad	26	0	0	<i>C. perfringens</i>	[327] Mashhad
8	2021	Cattle and sheep carcasses	200	61	30.5	<i>C. perfringens</i>	[328] Shiraz

Table 9 cont.

9	2021	Broiler chickens	122	95	77.8	<i>C. perfringens</i>	[329]	Kerman
10	2019	Broiler chickens	400	169	42.2	<i>C. perfringens</i>	[330]	Chaharmahal & Bakhtiari
11	2019	Honey	130	0	0	<i>C. perfringens</i>	[331]	
12	2017	Traditional curds	50	12	25	<i>C. perfringens</i>	[332]	Shahrekord
		Commercial curds	50	5	10			
		Beef meat	20	1	6			
		Lamb meat	23	3	13			
13	2015	Broiler meat	200	31	15.5	<i>C. perfringens</i>	[333]	Mashhad
14	2015	Minced meat	200	25	12.5	<i>C. perfringens</i>	[334]	Mashhad
15	2013	Honey	100	2	2	<i>C. perfringens</i>	[335]	Shiraz
16	2013	Fish	80	4	5	<i>C. perfringens</i>	[336]	Shiraz
		Honey	50	2	4			
		Kashk	80	2	2.5			
		Dough	80	1	1.2			
17	2013	Dairy products	57	12	21	<i>C. perfringens</i>	[337]	Gilan, Tehran, Golestan, Hamedan
		Fish	68	18	26.4			
		Meat	14	1	7.1			
18	2010	Cheese	57	2	3.5	<i>C. perfringens</i>	[338]	Gilan
		Kashk	11	0	0			
		Salted fish	63	4	6.3			

tables and juices, and other related products. Further research and surveillance are necessary to gain a more comprehensive understanding of the prevalence and sources of *C. perfringens* in various food items in Iran.

10) *Brucella* prevalence in food

Brucella spp. are the causative agents of brucellosis [56], an infectious disease of humans that presents with chronic and recurring febrile symptoms that can be life-threatening [57]. The primary etiological agent of the disease is *B. melitensis*, although other species, including *B. abortus*, *B. canis*, and *B. suis*, can also result in human brucellosis [58]. The infection can be transmitted to humans from various animals, including buffalo, cattle, yak, elk, camel, domestic pig, and rodents [58]. Globally, approximately 500,000 cases of human brucellosis are reported annually, with animals and animal-derived foods serving as the primary sources of infection [57]. A global systematic review conducted in 2020 revealed that the Southeast Asia region exhibited the highest prevalence of *Brucella spp.* at 25.55% [57]. The consumption of unpasteurized dairy products plays a significant role in the transmission of *Brucella spp.* to humans [57]. Table 10 presents the results of studies conducted in Iran on the prevalence of *Brucella spp.* in food. As illustrated in Figure 11, the primary sources of reported contam-

ination with *Brucella spp.* are dairy products (34.28%) and raw milk (16.64%). Dairy products, particularly unpasteurized or inadequately pasteurized ones, can serve as reservoirs for *Brucella* contamination [57]. This can occur due to infected dairy animals shedding the bacteria in their milk. Raw milk, in particular, has been identified as a common source of *Brucella* infection in various parts of the world, including Iran. Improper handling and processing of raw milk can contribute to the transmission of *Brucella spp.* to humans [59].

In Iran, where dairy products hold cultural and dietary significance, ensuring the safety of these products from *Brucella* contamination is crucial for public health [60]. Implementing stringent control measures in dairy production, processing, and distribution can help mitigate the risk of *Brucella* transmission through dairy products and raw milk [57, 59, 60].

11) *Vibrio* prevalence in food

Vibrio spp. are halophilic marine bacteria. Some species, including *V. cholerae*, *V. parahaemolyticus*, and *V. vulnificus*, have the potential to cause gastroenteritis or septicemia in humans. The primary mode of transmission for this foodborne illness is the ingestion of raw, undercooked, or mishandled seafood contaminated by bacteria [61]. Table 11 presents the re-




Figure 11. Prevalence of *Brucella* in different foods in Iran.

Table 10. Summary of the studies reporting the prevalence of *Brucella spp.* in Iran.

Year	Sample type	Sample size	Positive samples (N)	Prevalence (%)	<i>Brucella spp.</i>	References	Area		
1	2022	Unpasteurized Milk and Dairy	291	12	4.1	<i>Brucella spp.</i>	[339]	Hamadan	
2	2021	Bovine Milk	240	16	6.66	<i>Brucella spp.</i>	[340]	Kurdistan	
3	2020	Dairy products	227	9	4	<i>Brucella spp.</i>	[341]	Hamadan	
		Non boiling milk	43	1	2.3				
		Fresh cheese Cream	21	2	[341]				
4	2019	Raw camel milk	96	3	3	<i>Brucella spp.</i>	[342]	Isfahan, Semnan	
			51	2	[342]				
5	2018	Dairy Products	208	60	28.8	<i>Brucella spp.</i>	[58]	Tehran	
			Goat raw milk	33	15				45.5
			Non-pasteurized cheese	23	9				39.1
			Sheep raw milk	33	9				27.3
			Cow raw milk	57	15				26.3
			Pasteurized cheese	28	7				25
6	2017	Dairy Products	14	11	78.6	<i>Brucella spp.</i>	[343]	Tehran	
			8	72.7	<i>B. melitensis</i>				
			3	27.3	<i>B. abortus</i>				
7	2017	Sheep raw milk Goat raw milk	530	41	8.1	<i>Brucella spp.</i>	[344]	Kerman	
8	2017	Raw milk	700	9	1.28	<i>Brucella spp.</i>	[345]	Kerman	
			Sheep's raw milk	300	3				1
9		Unpasteurized milk	400	6	1.5	<i>Brucella spp.</i>	[346]	Isfahan	
			Dairy products	132	4				3
10	2016	Cow's raw milk	48	4	8.3	<i>Brucella spp.</i>	[347]	Kerman	

Table 10 cont.

11	2016	Milk	225	20	8.9	<i>Brucella spp.</i>	[348]	Shahrekord & Isfahan
		Sheep milk	125	12	9.6			
		Goat milk	100	18	18			
12	2016	Raw goat milk	470	51	10.8	<i>Brucella spp.</i>	[344]	Southeast region of Iran
		Raw sheep milk	330	18	5.4			
13	2015	Raw milk	60	32	53.3	<i>Brucella spp.</i>	[200]	Zanjan
		Raw cow milk	57	19	33			
		Pasteurized cow milk	34	10	29			
14	2014	Pasteurized cheese	28	8	28	<i>Brucella spp.</i>	[349]	Tehran
		Traditional cheese	23	14	60			
		Raw goat milk	33	21	63			
		Raw sheep milk	33	19	57			
15	2013	Cattle milk	1117	18	1.6	<i>Brucella spp.</i>	[350]	Urmia
		Sheep milk	598	99	16.5			

sults of studies conducted in Iran on the prevalence of *Vibrio spp.* in different types of food. *Vibrio spp.* were predominantly detected in seafood, including lobster, fish products, crayfish, fish, and shrimp, as well as drinking water. As illustrated in Figure 12, the prevalence of *Vibrio spp.* was highest in seafood, with fish exhibiting the greatest incidence (49.33%), followed by lobster (21.53%), crayfish (8.63%), shrimp (8.12%), fish products (7.8%), and drinking water (1.3%) (Figure 12). The findings from Iran are in alignment with those from other countries. For instance, a comprehensive systematic review conducted in 2016 revealed that *V. parahaemolyticus* contamination was observed in 63.4% of oysters, 52.9% of clams, 51% of fish, and 48.3% of shrimps [62]. A similar study in China in

2020 reported that 15.34% of shrimp samples, 14.17% of fish samples, and 3.67% of RTE food were contaminated with *V. parahaemolyticus* [63]. However, there are no reports available from Iran regarding the prevalence of *V. parahaemolyticus* in RTE foods.

12) *Shigella* prevalence in food

The *Shigella* genus encompasses four known species: *S. dysenteriae*, *S. boydii*, *S. flexneri*, and *S. sonnei*, which have also been classified as subgroups A to D, respectively [64]. While *S. flexneri* has traditionally been reported as the main cause of shigellosis in developing countries, recent studies have shown that *S. sonnei* has become the predominant species of *Shigella* in Iran [64]. According to the WHO, *Shigella*




Figure 12. Prevalence of *Vibrio* in different foods in Iran.

Table 11.
Summary of the studies reporting the prevalence of *Vibrio spp.* in Iran.

Year	Sample type	Sample size	Positive sam- ples (N)	Prevalence (%)	Type of <i>Vibrio spp.</i>	References	Area	
1	2021	Fish	64	61	95	<i>Vibrio species</i>	[351]	Isfahan
2	2020	Frozen fish	200	0	0	<i>V. parahaemolyticus</i>	[352]	Mazandaran
3		Shrimp	70	12	17.1	<i>V. parahaemolyticus</i>	[353]	Zanjan
4	2018	Cold Smoked Salt- ed Fishes	200	46	23	<i>Vibrio spp.</i>	[354]	Mazandaran
5	2016	Fish	58	18	31	<i>V. parahaemolyticus</i>	[355]	Persian Gulf
		Shrimps	55	7	12.7			
6	2015	Fresh shrimps	30	2	6.6	<i>V. parahaemolyticus</i>	[356]	Genaveh
		Salted shrimps	30	2	6.6			seaport
7	2014	Shrimps	36	7	19.4	<i>Vibrio spp.</i>	[357]	South coast of Iran
8	2014	Fish	100	22	22	<i>V. parahaemolyticus</i>	[358]	Bushehr, Persian Gulf
		Lobster	60	13	21.6			
		Crab caught	40	7	17.5			
9	2014	Crayfish	97	11	11.3	<i>V. vulnificus</i>	[357]	Aras
				7	7.2	<i>V. harveyi</i>		
				2	2	<i>V. alginolyticus</i>		
				1	1	<i>V. mimicus</i>		
10	2013	Tap-water	144	3	2	<i>V. cholerae</i>	[360]	Isfahan
		Bottled mineral water	304	3	0.6			
11	2012	Fresh shrimp	70	5	7.1	<i>V. parahaemolyticus</i>	[122]	
		Salted fishes	70	2	2.9			
		Fish nugget	10	0	0			
		Shrimp burger	10	0	0			
12	2012	Lobsters	100	40	40	<i>Vibrio spp.</i>	[361]	Persian Gulf
			100	3	3	<i>V. parahaemolyticus</i>		
		Crab	32	4	12.5	<i>Vibrio spp.</i>		
			32	1	3.1	<i>V. parahaemolyticus</i>		
13	2010	Fresh shrimp	300	29	9.6	<i>V. parahaemolyticus</i>	[361]	Bohsheer. Hor- mozgan, Khoozc- stan
14	2004	Fresh shrimp	770	16	2.1	<i>Vibrio spp.</i>	[361]	Bohsheer. Hor- mozgan, Khooz- estan

spp. cause approximately 165 million cases of *bacillary* dysentery and 1 million deaths worldwide each year [64]. In general, *Shigella spp.* are among the most prevalent causes of acute diarrhea in Iran, with a particularly high incidence among children and young adults. A diverse array of foods, encompassing meat, dairy products, and vegetables, have been identified as potential sources of shigellosis outbreaks worldwide [64]. Table 12 presents the results of studies conducted in Iran on the prevalence of *Shigella spp.* in different

types of food. As illustrated in Figure 13, contamination with *Shigella spp.* is most commonly reported in RTE foods (1.72%) and vegetables (1.05%), followed by red meat (0.4%). In contrast to the data from Iran, a high prevalence of *Shigella spp.* contamination has been reported in vegetables (25.25%) in India [65], and in beef, chicken, and dairy products in Egypt [66]. According to our review, poultry meat should be considered a high-risk food with the potential to spread foodborne zoonoses in Iran. In general, poultry meat

Table 12.
Summary of the studies reporting the prevalence of *Shigella spp.* in Iran.

Year	Sample type	Sample size	Positive sam- ples (N)	Prevalence (%)	Type of <i>Shigella spp.</i>	References	Area	
1	2022	raw milk, ground meat, and raw vegetable	580	13	2.24	<i>Shigella sonnei</i>	[364]	Tehran and Qazvin
2	2021	Vegetable salad, ground meat, and raw cow's milk	405	18	4.44	<i>Shigella spp.</i>	[365]	Qazvin
3	2021	meat, vegetable salad and raw milk	165	8	4.84	<i>Shigella spp.</i>	[366]	Qazvin
4	2019	Ready to eat food	250	2	0.8	<i>S. sonnei</i>	[64]	Isfahan, Fars, Hormozgan, Kohkiluyeh va Boyer Ahmad
			250	0	0	<i>S. flexneri</i>		
			250	0	0	<i>S. dysenteriae</i>		
			250	0	0	<i>S. boydii</i>		
		Fresh meat	150	1	0.7	<i>S. sonnei</i>		
			150	2	1.3	<i>S. flexneri</i>		
			150	0	0	<i>S. dysenteriae</i>		
		Frozen meat	150	0	0	<i>S. boydii</i>		
			100	0	0	<i>Shigella spp.</i>		
			100	0	0	<i>Shigella spp.</i>		
Vegetables	650	650	8	1.2	<i>S. sonnei</i>			
		650	6	0.9	<i>S. flexneri</i>			
		650	0	0	<i>S. dysenteriae</i>			
		650	0	0	<i>S. boydii</i>			
1400	19	1.4	<i>Shigella spp.</i>					
5	2018	Ready-to-Eat Salad	90	7	7.8	<i>Shigella spp.</i>	[367]	Tehran
6	2018	Food (vegetables, chicken, minced meat, fish)	100	6	6	<i>Shigella spp.</i>	[368]	Shiraz
7	2014	Camel milk	18	0	0	<i>Shigella spp.</i>	[369]	Golestan

is more susceptible to contamination during processing and handling due to its higher water content and pH levels, which provide an optimal environment for the proliferation of bacteria [67]. Moreover, poultry meat is frequently sold and consumed in its raw state, thereby increasing the probability of contamination if the requisite hygiene standards are not observed during slaughter, processing, and storage. In contrast, red meat and seafood have lower contamination rates compared to poultry meat, likely due to differences in processing and handling practices [68]. These findings underscore the necessity of developing strategies to reduce the contamination levels of poultry meat to effectively control and prevent foodborne illnesses in Iran.

The risk of food contamination, particularly in meat products, is significant. However, to effectively

underscore the importance of foodborne diseases, it is imperative to document the consequences of infection with these pathogens and generalize this information to the population in Iran. Currently, foodborne diseases in Iran are not generally reported, leading to a likely gross underestimation of their burden. This underestimation is attributable to the fact that many foodborne illnesses do not exhibit sufficient severity, duration, or specific diagnostic criteria for accurate identification and intervention. Similar circumstances exist in developed countries, such as the United States. For instance, the CDC estimates that foodborne pathogens cause approximately 48 million illnesses, 128,000 hospitalizations, and 3,000 deaths annually in the US [70].

Therefore, it is crucial to emphasize the necessity of establishing robust monitoring systems in Iran.

Such a surveillance network would require the collaboration of multidisciplinary teams comprising medical doctors, veterinarians, microbiologists, public health specialists, and other relevant experts, in alignment with the One Health concept. By adopting a methodology similar to that employed by the CDC's Foodborne Diseases Active Surveillance Network (FoodNet), which monitors the incidence of nine foodborne pathogens in ten US cities, representing approximately 15% of the American population [71], Iran can enhance the awareness of foodborne disease events and trends. These practices enable the implementation of effective intervention and prevention strategies.

Authors' Contributions

MH suggested the topic and supervised the conduction of the systematic review. SA wrote the first draft of the manuscript. FA and SA performed the literature review. GS was the major contributor in writing the manuscript. AA gave advice for conducting and writing the manuscript. All authors read and approved the final manuscript.

Competing Interests

The authors declare that there is no conflict of interest.

Acknowledgements

It is not applicable

References

1. Touranlou FA, Noori SMA, Salari A, Afshari A, Hashemi M. Application of kefir for reduction of contaminants in the food industry: A systematic review. *International Dairy Journal*. 2023;105748. Doi:10.1016/j.idairyj.2023.105748.
2. Law JW-F, Ab Mutalib N-S, Chan K-G, Lee L-H. Rapid methods for the detection of foodborne bacterial pathogens: principles, applications, advantages and limitations. *Frontiers in microbiology*. 2015;5:122692. Doi:10.3389/fmicb.2014.00770.
3. Imre K, Herman V, Morar A. Scientific achievements in the study of the occurrence and antimicrobial susceptibility profile of major foodborne pathogenic bacteria in foods and food processing environments in Romania: Review of the last decade. *BioMed Research International*. 2020;2020. Doi:10.1108/NFS-01-2023-0023.
4. Shakeri G, Jamshidi A, Ghazvini K. Isolation of broad-host-range bacteriophages against food-and patient-derived Shiga toxin-producing *Escherichia coli*. *Iranian Journal of Veterinary Science and Technology*. 2020;12(1):11-20. Doi:10.22067/veterinary.v12i1.85298.
5. Moghimani M, Noori SMA, Afshari A, Hashemi M. D-tryptophan, an eco-friendly natural, safe, and healthy compound with antimicrobial activity against food-borne pathogens: A systematic review. *Food Science & Nutrition*. 2024;12(5):3068-79. Doi:10.1002/fsn.3.3987.
6. Gourama H. *Foodborne pathogens*. Food safety engineering: Springer; 2020. p. 25-49. Doi:10.1007/978-3-030-42660-6_2.
7. Tîrziu E, Bărbălan G, Morar A, Herman V, Cristina RT, Imre K. Occurrence and antimicrobial susceptibility profile of *Salmonella* spp. in raw and ready-to-eat foods and *Campylobacter* spp. in retail raw chicken meat in Transylvania, Romania. *Foodborne Pathogens and Disease*. 2020;17(8):479-84. Doi:10.1089/fpd.2019.2738.
8. Hashemi M, Erfani A, Dadgar Moghadam M, Dousti Nouri M, Asadi Touranlou F, Azadi N, et al. Assessment of knowledge, Attitudes, and Practices of Food Service Staff Regarding Food Safety at Mashhad University Hospitals. *Journal of Nutrition, Fasting and Health*. 2024. Doi: 10.22038/JNFH.2024.78388.1505.
9. WHO. WHO estimates of the global burden of foodborne diseases: foodborne disease burden epidemiology reference group 2007-2015: World Health Organization; 2015.
10. Tack DM, Marder EP, Griffin PM, Cieslak PR, Dunn J, Hurd S, et al. Preliminary incidence and trends of infections with pathogens transmitted commonly through food—Foodborne Diseases Active Surveillance Network, 10 US Sites, 2015–2018. *Morbidity and Mortality Weekly Report*. 2019;68(16):369. Doi:10.15585/mmwr.mm6816a2.
11. Pires SM, Evers EG, Van Pelt W, Ayers T, Scallan E, Angulo FJ, et al. Attributing the human disease burden of foodborne infections to specific sources. *Foodborne pathogens and disease*. 2009;6(4):417-24. Doi:10.1089/fpd.2008.0208.
12. Mukhopadhyay S, Ramaswamy R. Application of emerging technologies to control *Salmonella* in foods: A review. *Food Research International*. 2012;45(2):666-77. Doi:10.1016/j.foodres.2011.05.016.
13. WHO. Estimating the burden of foodborne diseases: a practical handbook for countries: a guide for planning, implementing and reporting country-level burden of foodborne disease. 2021.
14. Yang J, Zhang Z, Zhou X, Cui Y, Shi C, Shi X. Prevalence and characterization of antimicrobial resistance in *Salmonella enterica* isolates from retail foods in Shanghai, China. *Foodborne Pathogens and Disease*. 2020;17(1):35-43. Doi:10.1089/fpd.2019.2671.

15. Nair DV, Kollanoor Johny A. Salmonella in poultry meat production. Food safety in poultry meat production. 2019:1-24. Doi:10.1007/978-3-030-05011-5_1.
16. Shafini A, Son R, Mahyudin N, Rukayadi Y, Zainazor TT. Prevalence of Salmonella spp. in chicken and beef from retail outlets in Malaysia. International Food Research Journal. 2017;24(1):437.
17. Carrasco E, Morales-Rueda A, García-Gimeno RM. Cross-contamination and recontamination by Salmonella in foods: A review. Food Research International. 2012;45(2):545-56. Doi:10.1016/j.foodres.2011.11.004.
18. Ehuwa O, Jaiswal AK, Jaiswal S. Salmonella, food safety and food handling practices. Foods. 2021;10(5):907. Doi:10.3390/foods10050907.
19. Ehuwa O, Jaiswal A, Jaiswal S. Salmonella, Food Safety and Food Handling Practices. Foods 2021, 10, 907. Food-borne Pathogens and Food Safety. 2021:175. Doi:10.3390/foods10050907.
20. Kowalska B. Fresh vegetables and fruit as a source of Salmonella bacteria. Annals of Agricultural and Environmental Medicine. 2023;30(1). Doi: 10.26444/aaem/156765.
21. Kadariya J, Smith TC, Thapaliya D. Staphylococcus aureus and staphylococcal food-borne disease: an ongoing challenge in public health. BioMed research international. 2014;2014. Doi:10.1155/2014/827965.
22. Naas HT, Edarhoby RA, Garbaj AM, Azwai SM, Abolghait SK, Gammoudi FT, et al. Occurrence, characterization, and antibiogram of Staphylococcus aureus in meat, meat products, and some seafood from Libyan retail markets. Veterinary World. 2019;12(6):925. Doi: 10.14202/vet-world.2019.925-931.
23. Dallal MMS, Foroushani AR, Sharifi-Yazdi S, Sharifi-Yazdi MK, Arfatahery N. Prevalence of Staphylococcus aureus in Shrimps in Tehran during 2013. Journal of Medical Bacteriology. 2015;4(5-6):42-6.
24. Pourbabaee M, Hadadi MR, Hooshyar H, Pourbabaee P, Nazari-Alam A. Prevalence of Staphylococcus aureus in Raw Hamburgers from Kashan in 2017. International Archives of Health Sciences. 2020;7(1):47-50. Doi: 10.4103/iahs.iahs_39_19.
25. Hamidiyan N, Salehi-Abargouei A, Rezaei Z, Dehghani-Tafti R, Akrami-Mohajeri F. The prevalence of Listeria spp. food contamination in Iran: A systematic review and meta-analysis. Food Research International. 2018;107:437-50. Doi:10.1016/j.foodres.2018.02.038.
26. Authority EFS, Prevention ECfD, Control. The European Union one health 2020 zoonoses report. EFSA Journal. 2021;19(12):e06971. Doi:10.2903/j.efsa.2021.6971.
27. Khademi F, Sahebkar A. A systematic review and meta-analysis on the prevalence of antibiotic-resistant Listeria species in food, animal and human specimens in Iran. Journal of food science and technology. 2019:1-17. Doi:10.1007/s13197-019-04040-w.
28. Dalton HR, Dreier J, Rink G, Hecker A, Janetzko K, Juhl D, et al. Coxiella burnetii-pathogenic agent of Q (query) fever. Transfusion medicine and hemotherapy. 2014;41(1):60-72. Doi:10.1159/000357107.
29. Abdali F, Hosseinzadeh S, Berizi E, Shams S. Prevalence of Coxiella burnetii in unpasteurized dairy products using nested PCR assay. Iranian Journal of Microbiology. 2018;10(4):220.
30. EFSA. The European Union one health 2018 zoonoses report. EFSA Journal. 2019;17(12):e05926. Doi:10.2903/j.efsa.2019.5926.
31. Rafiee S, Khoshnevisan B, Mohammadi I, Aghbashlo M, Clark S. Sustainability evaluation of pasteurized milk production with a life cycle assessment approach: an Iranian case study. Science of the total environment. 2016;562:614-27. Doi:10.1016/j.scitotenv.2016.04.070.
32. Moradnejad P, Esmaili S, Maleki M, Sadeghpour A, Kamali M, Rohani M, et al. Q fever endocarditis in Iran. Scientific reports. 2019;9(1):15276. Doi:10.1038/s41598-019-51600-3.
33. Van den Brom R, Van Engelen E, Roest H, Van der Hoek W, Vellema P. Coxiella burnetii infections in sheep or goats: an opinionated review. Veterinary microbiology. 2015;181(1-2):119-29. Doi:10.1016/j.vetmic.2015.07.011.
34. Guidi F, Petruzzelli A, Ciarrocchi F, Duranti A, Valiani A, Amagliani G, et al. Prevalence of Coxiella burnetii in cows' and ewes' bulk tank milk samples from selected dairy farms of Central Italy. Italian Journal of Animal Science. 2017;16(4):673-6. Doi:10.1080/1828051X.2017.1321474.
35. Rozental T, Faria LSD, Forneas D, Guterres A, Ribeiro JB, Araújo FR, et al. First molecular detection of Coxiella burnetii in Brazilian artisanal cheese: a neglected food safety hazard in ready-to-eat raw-milk product. Brazilian Journal of Infectious Diseases. 2020;24(3):208-12. Doi:10.1016/j.bjid.2020.05.003.
36. Kim SG, Kim EH, Lafferty CJ, Dubovi E. Coxiella burnetii in bulk tank milk samples, United States. Emerging infectious diseases. 2005;11(4):619. Doi: 10.3201/eid1104.041036.
37. Berthold-Pluta A, Pluta A, Garbowska M, Stefańska I. Prevalence and toxicity characterization of Bacillus cereus in food products from Poland. Foods. 2019;8(7):269. Doi:10.3390/foods8070269.
38. Tewari A, Abdullah S. Bacillus cereus food poisoning: international and Indian perspective. Journal of food science and technology. 2015;52(5):2500-11. Doi:10.1007/s13197-014-1344-4.

39. Webb MD, Barker GC, Goodburn KE, Peck MW. Risk presented to minimally processed chilled foods by psychrotrophic *Bacillus cereus*. *Trends in food science & technology*. 2019;93:94-105. Doi:10.1016/j.tifs.2019.08.024.
40. Eglezos S, Huang B, Dykes GA, Fegan N. The prevalence and concentration of *Bacillus cereus* in retail food products in Brisbane, Australia. *Foodborne pathogens and disease*. 2010;7(7):867-70. Doi:10.1089/fpd.2009.0469.
41. Yu S, Yu P, Wang J, Li C, Guo H, Liu C, et al. A study on prevalence and characterization of *Bacillus cereus* in ready-to-eat foods in China. *Frontiers in microbiology*. 2020;10:3043. Doi:10.3389/fmicb.2019.03043.
42. Gupta V, Gulati P, Bhagat N, Dhar M, Virdi J. Detection of *Yersinia enterocolitica* in food: an overview. *European Journal of Clinical Microbiology & Infectious Diseases*. 2015;34(4):641-50. Doi:10.1007/s10096-014-2276-7.
43. Lucero-Estrada C, Favier GI, Escudero ME. An overview of *Yersinia enterocolitica* and related species in samples of different origin from San Luis, Argentina. *Food microbiology*. 2020;86:103345. Doi:10.1016/j.fm.2019.103345.
44. Mahdavi S, Farshchian MR, Amini K, Abbasi M, Rad MG, Ebadi AR. Survey of *Yersinia enterocolitica* contamination in distributed broiler meats in Tabriz City, Iran. *African J Microbiol Res*. 2012;6(12):3019-23. Doi: 10.5897/AJMR12.094.
45. Zadernowska A, Chajęcka-Wierzchowska W, Łaniewska-Trokenheim Ł. *Yersinia enterocolitica*: a dangerous, but often ignored, foodborne pathogen. *Food Reviews International*. 2014;30(1):53-70. Doi:10.1080/87559129.2013.853775.
46. Chlebicz A, Ślizewska K. *Campylobacteriosis, salmonellosis, yersiniosis, and listeriosis as zoonotic foodborne diseases: a review*. *International journal of environmental research and public health*. 2018;15(5):863. Doi:10.3390/ijerph15050863
47. Soro AB, Whyte P, Bolton DJ, Tiwari BK. Strategies and novel technologies to control *Campylobacter* in the poultry chain: A review. *Comprehensive Reviews in Food Science and Food Safety*. 2020. Doi:10.1111/1541-4337.12544.
48. Rahimi E, Ameri M, Alimoradi M, Chakeri A, Bahrami AR. Prevalence and antimicrobial resistance of *Campylobacter jejuni* and *Campylobacter coli* isolated from raw camel, beef, and water buffalo meat in Iran. *Comparative Clinical Pathology*. 2013;22(3):467-73. Doi:10.1007/s00580-012-1434-5.
49. Thames H, Sukumaran A. A Review of *Salmonella* and *Campylobacter* in Broiler Meat: Emerging Challenges and Food Safety Measures. *Foods*. 2020;9(6):776. Doi:10.3390/foods9060776.
50. Quaglia NC, Dambrosio A. *Helicobacter pylori*: A food-borne pathogen? *World journal of gastroenterology*. 2018;24(31):3472. Doi: 10.3748/wjg.v24.i31.3472.
51. Bahrami AR, Rahimi E, Ghasemian Safaei H. Detection of *Helicobacter pylori* in city water, dental units' water, and bottled mineral water in Isfahan, Iran. *The Scientific World Journal*. 2013;2013. Doi:10.1155/2013/280510.
52. Souod N, Kargar M, Doosti A, Ranjbar R, Sarshar M. Genetic analysis of *cagA* and *vacA* genes in *Helicobacter pylori* isolates and their relationship with gastroduodenal diseases in the west of Iran. *Iranian Red Crescent Medical Journal*. 2013;15(5):371. Doi: 10.5812/ircmj.3732.
53. Fujimura S, Kawamura T, Kato S, Tateno H, Watanabe A. Detection of *Helicobacter pylori* in cow's milk. *Letters in Applied Microbiology*. 2002;35(6):504-7. Doi:10.1046/j.1472-765X.2002.01229.x.
54. Quaglia N, Dambrosio A, Normanno G, Parisi A, Patrono R, Ranieri G, et al. High occurrence of *Helicobacter pylori* in raw goat, sheep and cow milk inferred by *glmM* gene: a risk of food-borne infection? *International journal of food microbiology*. 2008;124(1):43-7. Doi:10.1016/j.ijfoodmicro.2008.02.011.
55. Meng X, Zhang H, Law J, Tsang R, Tsang T. Detection of *Helicobacter pylori* from food sources by a novel multiplex PCR assay. *Journal of food safety*. 2008;28(4):609-19. Doi:10.1111/j.1745-4565.2008.00135.x.
56. Mirnejad R, Jazi FM, Mostafaei S, Sedighi M. Epidemiology of brucellosis in Iran: A comprehensive systematic review and meta-analysis study. *Microbial pathogenesis*. 2017;109:239-47. Doi:10.1016/j.micpath.2017.06.005.
57. Dadar M, Fakhri Y, Shahali Y, Khaneghah AM. Contamination of milk and dairy products by *Brucella* species: A global systematic review and meta-analysis. *Food Research International*. 2020;128:108775. Doi:10.1016/j.foodres.2019.108775.
58. Moslemi E, Soltandadal MM, Beheshtizadeh MR, Taghavi A, Kheiri Manjili H, Mahmoudi Lamouki R, et al. Detection of *Brucella* spp. in dairy products by real-time PCR. *Archives of Clinical Infectious Diseases*. 2018;13(1). Doi:10.5812/archcid.12673.
59. Esmaeili S, Naddaf SR, Pourhossein B, Hashemi Shahraki A, Bagheri Amiri F, Gouya MM, et al. Seroprevalence of brucellosis, leptospirosis, and Q fever among butchers and slaughterhouse workers in south-eastern Iran. *PloS one*. 2016;11(1):e0144953. Doi:10.1371/journal.pone.0144953.
60. Islam MS, Islam MA, Rahman MM, Islam K, Islam MM, Kamal MM, et al. Presence of *Brucella* spp. in milk and dairy products: a comprehensive review and its perspectives. *Journal of Food Quality*. 2023;2023(1):2932883. Doi:10.1371/journal.pone.0144953.
61. Letchumanan V, Chan K-G, Lee L-H. *Vibrio parahaemolyticus*: a review on the pathogenesis, prevalence, and advance

- molecular identification techniques. *Frontiers in microbiology*. 2014;5:705. Doi:10.3389/fmicb.2014.00705.
62. Odeyemi OA. Incidence and prevalence of *Vibrio parahaemolyticus* in seafood: a systematic review and meta-analysis. *Springerplus*. 2016;5(1):464. Doi:10.1186/s40064-016-2115-7.
 63. Li Y, Xie T, Pang R, Wu Q, Zhang J, Lei T, et al. Food-Borne *Vibrio parahaemolyticus* in China: Prevalence, Antibiotic Susceptibility, and Genetic Characterization. *Frontiers in microbiology*. 2020;11:1670. Doi:10.3389/fmicb.2020.01670.
 64. Shahin K, Bouzari M, Wang R, Yazdi M. Prevalence and molecular characterization of multidrug-resistant *Shigella* species of food origins and their inactivation by specific lytic bacteriophages. *International journal of food microbiology*. 2019;305:108252. Doi:10.1016/j.ijfoodmicro.2019.108252.
 65. Gundappa M, Gaddad SM. Prevalence of *Salmonella*, *Shigella* and *E. coli* in Vegetables of Various Markets in Kalaburagi (India). *Indian J Nat Sci*. 2016;6(1):1-5. Doi:10.1007/s13197-017-2964-2.
 66. Shahin K, Bouzari M. Bacteriophage application for biocontrolling *Shigella flexneri* in contaminated foods. *Journal of food science and technology*. 2018;55(2):550-9.
 67. Rahman S, Park J, Song KB, Al-Harbi NA, Oh DH. Effects of slightly acidic low concentration electrolyzed water on microbiological, physicochemical, and sensory quality of fresh chicken breast meat. *Journal of food science*. 2012;77(1):M35-M41. Doi:10.1111/j.1750-3841.2011.02454.x.
 68. Sofos JN. Meat and meat products. *Food safety management: Elsevier*; 2014. p. 119-62. Doi:10.1016/B978-0-12-381504-0.00006-8.
 69. Pesavento G, Ducci B, Nieri D, Comodo N, Nostro AL. Prevalence and antibiotic susceptibility of *Listeria* spp. isolated from raw meat and retail foods. *Food control*. 2010;21(5):708-13. Doi:10.1016/j.foodcont.2009.10.012.
 70. Centers for Disease Control and Prevention [Internet]. 2011. Available from: <https://www.cdc.gov/foodborneburden/2011-foodborne-estimates.html>.
 71. Tack DM. Preliminary incidence and trends of infections with pathogens transmitted commonly through food—Foodborne Diseases Active Surveillance Network, 10 US sites, 2015–2018. *MMWR Morbidity and mortality weekly report*. 2019;68.
 72. Rahimi Z, Mahmoudi R, Ghajarbeyg P, Mosavi S, Mehrabi A. The Prevalence of *Salmonella* Enteritidis in Packaged and Tray Eggs Samples, Qazvin, Iran. *Journal of Chemical Health Risks*. 2021. Doi:10.22034/jchr.2021.1922123.1250.
 73. Nazari Moghadam M, Rahimi E, Shakerian A, Momtaz H. Prevalence of *Salmonella* Typhimurium and *Salmonella* Enteritidis isolated from poultry meat: virulence and antimicrobial-resistant genes. *BMC microbiology*. 2023;23(1):1-8. Doi:10.1186/s12866-023-02908-8.
 74. Saram M, Nosrati AC, Modiri L, Issazadeh K. Investigation of Genetic-molecular Sources of Multidrug Resistance of *Salmonella* Isolated from Eggs. *Social Determinants of Health*.8:1-11. Doi:10.22037/sdh.v8i1.39325.
 75. Hassanzadeh P, Mohammadzadeh S. Antimicrobial Effects of Different Synbiotic Compounds against Pathogenic Bacteria Isolated from Beef, Mutton, and Chicken. *Archives of Razi Institute*. 2022;77(6):2105. Doi: 10.22092/ARI.2022.357834.2107.
 76. Shayegh J, Shahbazi G, Ghazaei S, Movassagh Ghazani MH, Hanifian S. Prevalence, virulence factor and antibiotics susceptibility patterns of *Salmonella* spp. from poultry products in Ardabil province. 2022.
 77. Rahimi Z, Ghajarbeygi P, Mahmoudi R, Mosavi S, Mehrabi A. PREVALENCE OF *SALMONELLA* STRAINS ISOLATED FROM INDUSTRIAL QUAIL EGGS AND LOCAL DUCK EGGS, IRAN. *Carpathian Journal of Food Science & Technology*. 2022;14(2). Doi:10.34302/crpfjst/2022.14.2.14.
 78. Mir R, Salari S, Najimi M, Rashki A. Determination of frequency, multiple antibiotic resistance index and resistotype of *Salmonella* spp. in chicken meat collected from southeast of Iran. *Veterinary Medicine and Science*. 2022;8(1):229-36. Doi:10.1002/vms3.647.
 79. Kazemeini H. Molecular detection of *Campylobacter* species and *Salmonella* spp. In cattle raw milk specimens in Mazandaran province. *Journal of food science and technology (Iran)*. 2022;19(125):101-8. Doi:10.22034/FSCT.19.125.101.
 80. Norbakhsh F, Behshood P. Determination of molecular properties of *Salmonella* isolates isolated from red meat in Shahrekord, Iran. *Journal of Zoonosis*. 2022;2(1):18-25.
 81. Azizpour A. Prevalence and antibiotic resistance of *Salmonella* serotypes in chicken meat of Ardabil, Northwestern Iran. *Iranian Journal of Medical Microbiology*. 2021;15(2):232-46. Doi: 10.30699/ijmm.15.2.232.
 82. Bahramianfard H, Derakhshandeh A, Naziri Z, Khaltabadi Farahani R. Prevalence, virulence factor and antimicrobial resistance analysis of *Salmonella* Enteritidis from poultry and egg samples in Iran. *BMC veterinary research*. 2021;17(1):196. Doi:10.1186/s12917-021-02900-2.
 83. Sabzali S, Bouzari M. Isolation, identification and some characteristics of two lytic bacteriophages against *Salmonella* enterica serovar Paratyphi B and *S. enterica* serovar Typhimurium from various food sources. *FEMS Microbiology Letters*. 2021;368(7):fnab037. Doi:10.1093/femsle/fnab037.

84. Afshari A, Tavassoli M, Ram M, Ranjbar G. Assessment of Microbial and Chemical Quality of a Ready to Eat Food, Olivier Salad, in Mashhad City. *Journal of Nutrition, Fasting and Health*. 2019;7(4 (Spe)):175-81.
85. Forouhar M, Harzandi N. Molecular Survey of Quinolone Resistance in Salmonella spp. Isolated From Poultry Products in Karaj, Iran. *Int J Enteric Pathog*. 2019;7(2):55-9. Doi: 10.15171/ijep.2019.13.
86. Mahdavi M, Jalali M, Safaei HG, Shamloo E. Microbial quality and prevalence of Salmonella and Listeria in eggs. *International Journal of Environmental Health Engineering*. 2012;1(1):48.
87. Hamzeh Pour S, Vaziri S, Molaee Aghae E. Survey on the contamination rate and determination of antibiotic resistance of Staphylococcus aureus, Escherichia coli and Salmonella strains isolated from traditional cheeses distributed in Mahabad, Iran. *Iranian Journal of Health and Environment*. 2019;11(4):465-76.
88. Mahdavi S, Azizi Dehbokri M, Isazadeh A. Contamination of chicken meat with salmonella spp distributed in mahabad city, iran. *Int J Enteric Pathog*. 2018;6(3):65-8. Doi: 10.15171/ijep.2018.18
89. Hashemi M, Baygan A, Balouchzahi Z, Dousti Nouri M, Afshari A. Evaluation of microbial quality of traditional sweets, Zoolbia and Bamieh, during Ramadan in Mashhad, Iran. *Journal of Nutrition, Fasting and Health*. 2018;6(3):145-9. Doi:10.22038/JNFH.2018.34900.1147.
90. Karimiazar F, Soltanpour MS, Aminzare M, Hassanzadazar H. Prevalence, genotyping, serotyping, and antibiotic resistance of isolated Salmonella strains from industrial and local eggs in Iran. *Journal of Food Safety*. 2019;39(1):e12585. Doi:10.1111/jfs.12585.
91. Rezaee R, Mohammadi S, Faizee S. A study of salmonella spp. contamination of eggs in Sanandaj in 2017. *Zanko Journal of Medical Sciences*. 2018;19(61):43-9.
92. Khedmati Morasa H, Mahmoudi R, Ghajarbeygi P, Abasi N, Mousavy S. Effects of cold-water egg shell washing on Salmonella contamination in the shell and its contents. *The Journal of Qazvin University of Medical Sciences*. 2018;22(2):83-9.
93. Sana M, Hosseini Siah Z. The determination of total microbial count and prevalence of Salmonella in the shrimp supply in Khuzestan province. *Iranian Journal of Health and Environment*. 2018;11(2):149-56.
94. Eslami A, Gholami Z, Nargesi S, Rostami B, Avazpour M. Evaluation of microbial contamination of ready-to-eat foods (pizza, frankfurters, sausages) in the city of Ilam. *Environmental Health Engineering and Management Journal*. 2017;4(2):117-22. Doi:10.15171/EHEM.2017.16.
95. Rahimi E, Shakeriyan A. Salmonella contamination of camel meat in various stages of destruction in Isfahan and Chaharmahal va Bakhtiari. *Journal of Food Microbiology*. 2017;4(2):29-36.
96. Sharafati Chaleshtori R, Mazroii Arani N, Taghizadeh M, Sharafati Chaleshtori F. Antibiotic resistance pattern of salmonella isolated from hamburgers and detection of their sensitivity to some essential oils. *Journal of Mazandaran University of Medical Sciences*. 2017;27(148):136-42.
97. Mojaddar Langroodi A, Ehsani A, Ebadi A. Comparison of bacteria in beef meat and poultry in terms of packaging and without packaging in Gilan province. *Journal of Food Microbiology*. 2016;3(1):33-42.
98. Dousandeh S, Asaadi Tehrani G, Amini B, Maziri P. Simultaneous detection of invA, STM4497, and fliC183 genes in Salmonella typhimurium by multiplex PCR method in poultry meat samples in Zanjan, Iran. *Infection, Epidemiology and Microbiology*. 2016;2(4):20-3.
99. Vahabi AN, Abbasvali M. Determination of microbial contamination of olovie salads consumed in Isfahan. 2016.
100. Khammar F, Alipour Eskandari M, Saadati D. Study of Salmonella contamination of traditional ice creams in Zabol City, Iran. *Iranian Journal of Medical Microbiology*. 2017;11(1):83-9.
101. Asadi S, Maram ZR, Kooshk F. Evaluation of microbial contamination of pastry cream in Arak city of Iran. *Journal of Food Safety and Hygiene*. 2015;1(1):26-9.
102. Tajbakhsh F, Tajbakhsh E, Momeni M. Detection of Staphylococcus aureus and Salmonella typhimurium in traditional and industrial Olivier salads in shahrekord city. 2015.
103. Khodadadipour T, Amini K, Mahmoudi R. Evaluation of virulence and enterotoxin genes in Salmonella enteritidis strains isolated from Meat and Egg samples by Multiplex-PCR. *Journal of Food Microbiology*. 2016;3(2):25-33.
104. Ghorbani Ranjbary A. Study of Drug Resistance in Salmonella spp. Isolated from Native Eggs of Iran's Southern Region. *Journal of pure and applied microbiology*. 2015;9(2):175-9.
105. Khoramrooz S, Sarikhani M, Khosravani S, Farhang Falah M, Mahmoudi Y, Sharifi A. Microbial contamination determination of cream suit, traditional ice cream and Olovia in Yasuj City. *Armaghane danesh*. 2015;20(6):526-37.
106. Doosti A, Doosti E, Rahimi E, Ghasemi-Dehkordi P. Frequency of antimicrobial-resistant genes in Salmonella enteritidis isolated from traditional and industrial iranian white cheeses. *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences*. 2017;87(1):73-80. Doi:10.1007/s40011-015-0572-3.

107. Moosavy M-H, Esmaeili S, Amiri FB, Mostafavi E, Salehi TZ. Detection of *Salmonella* spp in commercial eggs in Iran. *Iranian journal of microbiology*. 2015;7(1):50.
108. Kheyri A, Fakhernia M, Haghghat-Afshar N, Hassanzadazar H, Kazemi-Ghoshchi B, Zeynali F, et al. Microbial contamination of meat products produced in the factories of West Azerbaijan Province, North West of Iran. *Global Veterinaria*. 2014;12(6):796-802. Doi: 10.5829/idosi.gv.2014.12.06.83145.
109. Soltan Dallal MM, Emadi Koochak H, Sharifi Yazdi MK, Taheri Mirghaed A, Choobineh H. Determination of *Yersinia* spp. and *Salmonella paratyphi B* isolated from possibly contaminated cream samples in the city of Tehran. *Journal of Payavard Salamat*. 2014;8(1):34-43.
110. Dallal MMS. Prevalence of *Salmonella* spp. in packed and unpacked red meat and chicken in south of Tehran. *Jundishapur Journal of Microbiology*. 2014;7(4).
111. Sodagari HR, Mashak Z, Ghadimianazar A. Prevalence and antimicrobial resistance of *Salmonella* serotypes isolated from retail chicken meat and giblets in Iran. *The Journal of Infection in Developing Countries*. 2015;9(05):463-9. Doi: 10.3855/jidc.5945.
112. Zahedi M, Rahimi E, Zahedi M, Momtaz H. Prevalence of *Salmonella enteritidis* and *S. typhimurium* in marketed meat in Shahrekord in 2014. *Journal of Shahrekord University of Medical Sciences*. 2017;19.
113. Momtaz H, Dehkordi FS, Rahimi E, Asgarifar A. Detection of *Escherichia coli*, *Salmonella* species, and *Vibrio cholerae* in tap water and bottled drinking water in Isfahan, Iran. *BMC public health*. 2013;13(1):556. Doi:10.1186/1471-2458-13-556.
114. Bidaki MZ, Pour AT, Dadpour S, Gholizadeh H. Prevalence of *Salmonella* in poultry carcasses serotypes in Birjand industrial slaughterhouses. *Journal of Birjand University of Medical Sciences*. 2013;20(02).
115. Monadi M, Kargar M, Naghiha A, Mohammadi R. *Salmonella* contamination of eggs of native Kohgiluyeh va Boyer-Ahmad using PCR1 techniques and the evaluation of drug resistance. *Armaghane danesh*. 2014;19(2):179-87.
116. Amirmozaffari N, Rahmani Z, Iesazadeh K. Evaluation of the level of contamination with *Salmonella* spp. in red meat, chicken, and domestic and industrial eggs produced in Talesh city and assessment of their antibiotic resistance pattern, Iran. *Qom University of Medical Sciences Journal*. 2013;7(5):60-5.
117. Kaseb F, Shiranian M, Abdar M, Aminalroayaei H, Fallahzadeh H. The prevalence of *salmonella* and *Staphylococcus aureus* in industrial Olivier salad in Yazd in 2013. *Tolooebehdasht*. 2015;14(3):51-9.
118. Rahimi E, Shakerian A, Falavarjani AG. Prevalence and antimicrobial resistance of *Salmonella* isolated from fish, shrimp, lobster, and crab in Iran. *Comparative Clinical Pathology*. 2013;22(1):59-62. Doi:10.1007/s00580-011-1368-3.
119. Rahimi E. Prevalence and antimicrobial resistance of *Salmonella* spp isolated from retail chicken, turkey and ostrich by-products in Iran. *Revue de MédecineVétérinaire*. 2012;163(52):271-5. Doi:10.1007/s00580-011-1368-3.
120. Karimi Darehabi H, Esmailneshad F, Ebrahimi Mohammadi K. Contamination of Fresh Beef to *Salmonella typhimurium* and *Salmonella enteritidis* in Sanandaj during 2012. *Food Hygiene*. 2012;2(3 (7)):41-8.
121. Vaskas AT, Dahesht AE, Seifi S, Rahmani M, Motaghifar A, Safanavaee R. Study and comparison of the bacterial contamination outbreak of chicken meat consumed in some cities of Mazandaran Province (Iran). *African Journal of Microbiology Research*. 2012;6(33):6286-90. Doi: 10.5897/AJMR12.984.
122. Zarei M, Maktabi S, Ghorbanpour M. Prevalence of *Listeria monocytogenes*, *Vibrio parahaemolyticus*, *Staphylococcus aureus*, and *Salmonella* spp. in seafood products using multiplex polymerase chain reaction. *Foodborne Pathogens and Disease*. 2012;9(2):108-12. Doi:10.1089/fpd.2011.0989.
123. Tajbakhsh F, Tajbakhsh E, Momeni M, Rahimi E, Sohrabi R. Occurrence and Antibiotic Resistance of *Salmonella* spp isolated from raw Cow's Milk from Shahrekord. *Iran Inter J Microbiol Res*. 2012;3:242-5. Doi: 10.5829/idosi.ijmr.2012.3.3.61302.
124. Safaei HG, Jalali M, Hosseini A, Narimani T, Sharifzadeh A, Raheimi E. The prevalence of bacterial contamination of table eggs from retail markets by *Salmonella* spp., *Listeria monocytogenes*, *Campylobacter jejuni* and *Escherichia coli* in Shahrekord, Iran. *Jundishapur Journal of Microbiology*. 2011;4(4).
125. Dallal MMS, Doyle MP, Rezadehbashi M, Dabiri H, Sanaei M, Modarresi S, et al. Prevalence and antimicrobial resistance profiles of *Salmonella* serotypes, *Campylobacter* and *Yersinia* spp. isolated from retail chicken and beef, Tehran, Iran. *Food Control*. 2010;21(4):388-92. Doi:10.1016/j.foodcont.2009.06.001.
126. Rahimi E, Ameri M, Kazemeini H, Elbagi M. PREVALENCE AND ANTIMICROBIAL RESISTANCE OF *SALMONELLA* ISOLATED FROM RETAIL RAW TURKEY, OSTRICH, AND PARTRIDGE MEAT IN IRAN. *Bulgarian Journal of Veterinary Medicine*. 2010;13(1).
127. Jamshidi A, Kalidari GA, Hedayati M. Isolation and identification of *Salmonella Enteritidis* and *Salmonella Typhimurium* from the eggs of retail stores in Mashhad, Iran using conventional culture method and multiplex PCR assay. *Journal of food safety*. 2010;30(3):558-68. Doi:10.1111/j.1745-4565.2010.00225.x.
128. Jamshidi Ae, basami m, afshari ns. Identification of *Salmo-*

- nella spp. and Salmonella typhimurium by a multiplex PCR-based assay from poultry carcasses in Mashhad-Iran. 2009.
129. Shapouri R, Rahnama M, Eghbalzadeh S. PREVALENCE OF SALMONELLA SEROTYPES IN POULTRY MEAT AND EGG AND DETERMINE THEIR ANTIBIOTIC SENSIVITY IN ZANJAN CITY. 2009.
 130. Soltan Dmm, Taremi M, Gachkar L, MODARESI S, Sanaei M, Bakhtiari R, et al. Characterization of antibiotic resistant patterns of Salmonella serotypes isolated from beef and chicken samples in Tehran. 2009.
 131. Namaei M, Ziaee M, Kafi MG. Prevalence of salmonella contamination in locally (non-industrially) produced eggs in Birjand (2006). Journal of Birjand University of medical sciences. 2009;16(02).
 132. Jalali M, Abedi D, Pourbakhsh SA, Ghokasin K. Prevalence of salmonella spp. in raw and cooked foods in Isfahan-Iran. Journal of food safety. 2008;28(3):442-52. Doi:10.1111/j.1745-4565.2008.00122.x.
 133. AR B. Qualitative and quantitative assessment of poultry carcasses contaminated with salmonella in Tehran industrial slaughter houses. 2008.
 134. Daneshmand A, Rasooli M, Kargar M, Kyani S. A survey on bacterial contamination of fresh traditional cheeses with Salmonella spp. and Staphylococcus aureus in Jahrom, Fars Province. ISMJ. 2007;10(1):19-26.
 135. Bonyadian M, Ale AS, MOTAHARI FA. Isolation and identification of Salmonellae from chicken carcasses in processing plants in Yazd province, central Iran. 2007.
 136. Namaei M, ZIAEI M, GHANAD KM. Prevalence of salmonella contamination in locally (non-industrially) produced eggs in Birjand (2006). 2009.
 137. Nourbakhsh SA, Rahimi E. The occurrence of some food-borne pathogens recovered from poultry meat in Shahrekord, Iran. Journal of Advanced Veterinary and Animal Research. 2023;10(2):205. Doi: 10.5455/javar.2023.j670.
 138. Azad A, Shakerian A. Evaluation of Staphylococcus Aureus, Escherichia Coli, and Salmonella Contamination in Sausages and Bologna Sold in Tehran City, Iran. Journal of Alternative Veterinary Medicine| Spring. 2023;6(16).
 139. Azimirad M, Nadalian B, Alavifard H, Panirani SN, Bonab SMV, Azimirad F, et al. Microbiological survey and occurrence of bacterial foodborne pathogens in raw and ready-to-eat green leafy vegetables marketed in Tehran, Iran. International Journal of Hygiene and Environmental Health. 2021;237:113824. Doi:10.1016/j.ijheh.2021.113824.
 140. Doudaran HO, Mahsk Z, Kohdar V. Distribution Of Antibiotic Resistance Genes Amongst The Staphylococcus Aureus Strains Isolated From Raw Milk And Traditional Dairy Products. Journal of Pharmaceutical Negative Results. 2022;9735-41.
 141. Feizolahnezhad A, Mashak Z. Study The Prevalence And Antibiotic Resistance Of Staphylococcus Aureus Bacteria Isolated From Street Ready To Eat Food Samples. Journal of Pharmaceutical Negative Results. 2022;9714-20. Doi:10.47750/pnr.2022.13.S09.1135.
 142. Kabiri E, Mashak Z. Antimicrobial resistance properties and enterotoxigenic gene profile of methicillin-resistant and methicillin-susceptible Staphylococcus aureus isolates from raw milk. Journal of Positive School Psychology. 2022;6(8):1244-52.
 143. Pajohesh R, Tajbakhsh E, Momtaz H, Rahimi E. Relationship between Biofilm Formation and Antibiotic Resistance and Adherence Genes in Staphylococcus aureus Strains Isolated from Raw Cow Milk in Shahrekord, Iran. International Journal of Microbiology. 2022;2022. Doi:10.1155/2022/6435774.
 144. Hassani S, Pakbin B, Allahyari S, Mahmoudi R, Mousavi S, Ghajarbeygi P. PREVALENCE AND ANTIMICROBIAL SUSCEPTIBILITY OF FOODBORNE BACTERIAL PATHOGENS ISOLATED FROM BAGHLAVA AN IRANIAN EXPORTING PASTRY SWEET. Carpathian Journal of Food Science & Technology. 2022;14(3). Doi:10.34302/crp-jfst/2022.14.3.5.
 145. Shariatifar M, Salavati-Hamedani M, Rezaeian-Doloei R, Rezaeigolestani M, Mohsenzadeh M. Antibiotic resistance and enterotoxin gene profiles of Staphylococcus aureus isolated from raw milk in Iran. Journal of the Hellenic Veterinary Medical Society. 2023;74(1):5433-40. Doi:10.12681/jhvms.29671.
 146. Bastam MM, Jalili M, Pakzad I, Maleki A, Ghafourian S. Pathogenic bacteria in cheese, raw and pasteurised milk. Veterinary Medicine and Science. 2021;7(6):2445-9. Doi:10.1002/vms3.604.
 147. Abbasi K, Tajbakhsh E, Momtaz H. Antimicrobial resistance and biofilm encoding genes amongst the Staphylococcus aureus bacteria isolated from meat and meat products. Egyptian Journal of Veterinary Sciences. 2021;52(1):55-62. Doi:10.21608/EJVS.2020.39385.1186.
 148. Maktabi S, Gharibi D, Hormozi T. Detection of Enterotoxin A Gene (sea) and Determination of Different Biotypes of Staphylococcus Aureus Isolates From Milk and Cheese in Khuzestan Province, Iran. Int J Enteric Pathog. 2021;9(3):117-23.
 149. Haghi F, Zeighami H, Hajiloo Z, Torabi N, Derakhshan S. High frequency of enterotoxin encoding genes of Staphylococcus aureus isolated from food and clinical samples. Journal of Health, Population and Nutrition. 2021;40(1):1-6. Doi:10.1186/s41043-021-00246-x.

150. Mesbah A, Mashak Z, Abdolmaleki Z. A survey of prevalence and phenotypic and genotypic assessment of antibiotic resistance in *Staphylococcus aureus* bacteria isolated from ready-to-eat food samples collected from Tehran Province, Iran. *Tropical Medicine and Health*. 2021;49:1-12. Doi:10.1186/s41182-021-00366-4.
151. Dadmehr A, Sadighara P, Zeinali T. A study on microbial and chemical characterization of mechanically deboned chicken in Tehran, Iran. *International Journal of Environmental Health Research*. 2022;32(11):2396-405. Doi.org/10.1080/09603123.2021.1967889.
152. Saadati A, Mashak Z, Yarmand MS. Prevalence of Staphylococcal Cassette Chromosome mec and Panton-Valentine Leukocidin Gene Amongst the Methicillin-resistant *Staphylococcus aureus* Strains Isolated From Fowl Meat. *Int J Enteric Pathog*. 2019;7(3):93-8. Doi:10.15171/ijep.2019.20.
153. Mahdavi S, Isazadeh A. Investigation of contamination rate and determination of pattern of antibiotic resistance in coagulase positive staphylococcus aureus isolated from domestic cheeses in Maragheh, Iran. *Pathobiology Research*. 2019;22(2):85-9.
154. Dehkordi FS, Gandomi H, Basti AA, Misaghi A, Rahimi E. Phenotypic and genotypic characterization of antibiotic resistance of methicillin-resistant *Staphylococcus aureus* isolated from hospital food. *Antimicrobial Resistance & Infection Control*. 2017;6(1):104. Doi:10.1186/s13756-017-0257-1.
155. Goudarzi B, Alipour V, Rezaei L, Dindarlu K, Heidari M, Rahmaniyan O. Bacteriological quality of ready to use salads at restaurants in Bandar Abbas Abstract. *Journal of Preventive Medicine*. 2016;3(3):38-1.
156. Varmazyar-najafi M, Pajohi-alamoti M, Mohammadzadeh A, Mahmoodi P. Detection of methicillin-resistance gene in *Staphylococcus aureus* isolated from traditional white cheese in Iran. *Archives of Hygiene Sciences*. 2016;5(4):302-9.
157. Arfatahery N, Davoodabadi A, Abedimohtasab T. Characterization of toxin genes and antimicrobial susceptibility of *Staphylococcus aureus* isolates in fishery products in Iran. *Scientific reports*. 2016;6(1):1-7. Doi:10.1038/srep34216.
158. Arabestani MR, Kazemian H, Tabar ZK, Hosseini SM. Prevalence of virulence genes, agr and antimicrobial resistance of *Staphylococcus aureus* isolated from food and dairy products in Hamadan, Iran. *Der Pharmacia Lettre*. 2016;8(8):62-7.
159. Dehkordi AA, Tajbakhsh E, Tajbakhsh F, Khamesipour F, Shahraki MM, Momeni H. Molecular typing of staphylococcus aureus strains from Iranian raw milk and dairy products by coagulase gene polymorphisms. *Adv Stud Biol*. 2015;7(4):169-77. Doi:10.12988/asb.2015.516.
160. Arfatahery N, Mirshafiey A, Abedimohtasab T, Zeinolabedinizamani M. Study of the prevalence of *Staphylococcus aureus* in marine and farmed shrimps in Iran aiming the future development of a prophylactic vaccine. *Procedia in Vaccinology*. 2015;9:44-9. Doi:10.1016/j.provac.2015.05.008.
161. Jamali H, Paydar M, Radmehr B, Ismail S, Dadrasnia A. Prevalence and antimicrobial resistance of *Staphylococcus aureus* isolated from raw milk and dairy products. *Food Control*. 2015;54:383-8. Doi:10.1016/j.foodcont.2015.02.013.
162. Sadeghi M. The prevalence of Enterotoxigenic Isolates of *Staphylococcus aureus* in Bovine and Sheep Bulk Tank Milk Samples of Maku. 2015. Doi:10.1016/j.foodcont.2015.02.013.
163. Eslami M, Koohi M, Zadehashem E, Khadiri B, Keshavarz H. Survey the presence of coagulase positive staphylococcus aureus in cottage cheeses produced from sheep milk and sold in Marand county. *Journal of Food Science & Technology* (2008-8787). 2014;12(46).
164. Madahi H, Rostami F, Rahimi E, Dehkordi FS. Prevalence of enterotoxigenic *Staphylococcus aureus* isolated from chicken nugget in Iran. *Jundishapur journal of microbiology*. 2014;7(8). Doi: 10.5812/jjm.10237.
165. Khoshkholgh MR, Isazadeh K. Evaluation of Microbial quality of food samples collected in the East region of Gilan. *Journal of Food Microbiology*. 2015;2(1):27-37.
166. Shabani S, Sadeghi Mahoonak A, Jalali H. Microbial contamination of pastry cream supplied in Gorgan. *Medical Laboratory Journal*. 2014;8(2):62-6.
167. Saadat YR, Fooladi AAI, Shapouri R, Hosseini MM, Khiabani ZD. Prevalence of enterotoxigenic *Staphylococcus aureus* in organic milk and cheese in Tabriz, Iran. *Iranian journal of microbiology*. 2014;6(5):345.
168. Fadaei A. Bacteriological quality of raw cow milk in Shahrekord, Iran. *Veterinary World*. 2014;7(4):240-3.
169. Al-musawi MHJ, Al-Rubaye MTS, Hosseini M. Culture and Molecular Detection of *Staphylococcus aureus* in Dairy Products of Ahwaz. *Al-Mustansiriyah Journal for Pharmaceutical Sciences*. 2014;14(2):103-7.
170. Farajvand N, Alimohammadi M. Prevalence of *Staphylococcus aureus* in Four Famous Brand of Doogh Produced in Iran. *Iranian Journal of Health and Environment*. 2014;7(1):85-94.
171. Shojaei M, KARIMI DH, Javadi A. Distribution of genes encoding biofilm production in *S. aureus* isolated from raw milk in Kurdistan. 2014.
172. Sepidarkish M, Ghane M. Isolation, identification and the presence of enterotoxin A gene in *Staphylococcus aureus* from meat products. 2014.

173. Shانهbandi D, Baradaran B, Sadigh-Eteghad S, Zarredar H. Occurrence of methicillin resistant and enterotoxigenic *Staphylococcus aureus* in traditional cheeses in the North West of Iran. *International Scholarly Research Notices*. 2014;2014. Doi:10.1155/2014/129580
174. Rahimi E. Enterotoxigenicity of *Staphylococcus aureus* isolated from traditional and commercial dairy products marketed in Iran. *Brazilian Journal of Microbiology*. 2013;44(2):393-9.
175. Rahimi E, Alian F. Presence of enterotoxigenic *Staphylococcus aureus* in cow, camel, sheep, goat, and buffalo bulk tank milk. *Veterinarski arhiv*. 2013;83(1):23-30.
176. Mirzaei H, Farhoudi H, Tavassoli H, Farajli M, Monadi A. Presence and antimicrobial susceptibility of methicillin-resistant *Staphylococcus aureus* in raw and pasteurized milk and ice cream in Tabriz by culture and PCR techniques. *African Journal of Microbiology Research*. 2012;6(32):6224-9. Doi: 10.5897/AJMR12.1701
177. Tavakoli H, Farhang K, AA KZ HE. Bacteriological quality of ready to eat food in four military restaurants. *Journal Mil Med*. 2012;13(4):207-12.
178. Shahraz F, Dadkhah H, Khaksar R, Mahmoudzadeh M, Hosseini H, Kamran M, et al. Analysis of antibiotic resistance patterns and detection of *mecA* gene in *Staphylococcus aureus* isolated from packaged hamburger. *Meat science*. 2012;90(3):759-63. Doi:10.1016/j.meatsci.2011.11.009
179. Alian F, Rahimi E, Shakerian A, Momtaz H, Riahi M, Momeni M. Antimicrobial resistance of *Staphylococcus aureus* isolated from bovine, sheep and goat raw milk. *Global Veterinaria*. 2012;8(2):111-4.
180. Sharafati Chaleshtori R, Sharafati Chaleshtori F, Karimi A. Antibiotic resistance pattern of *staphylococcus* strains isolated from orange and apple juices in Shahre-kord, Iran. *Pakistan Journal of Medical Sciences*. 2010;26(3):615-8.
181. Tavakoli HR, Riazipour M. Microbial quality of cooked meat foods in Tehran University's Restaurants. *Pakistan Journal of Medical Sciences*. 2008;24(4):595-9.
182. Basti AA, Misaghi A, Salehi TZ, Kamkar A. Bacterial pathogens in fresh, smoked and salted Iranian fish. *Food control*. 2006;17(3):183-8. Doi:10.1016/j.foodcont.2004.10.001
183. Raeisi M, Afshari A, Shirzad-Aski H, Seifi S, Hashemi M, Khoshbakht R, et al. The Occurrence of Serotypes and Virulence Genes of *Listeria monocytogenes* in Various Food Products. *Journal of food quality and hazards control*. 2023. Doi:10.18502/jfqhc.10.2.12673
184. Kiyandpour Berjoe R, Momtaz H, Lotfollahi L, Bamzaheh Z. Detection and antibiotic resistance pattern of *Listeria monocytogenes* strains isolated from curd and cheese. *Journal of Food Microbiology*. 2022;9(4):18-29. Doi:10.30495/jfm.2022.1943942.1741
185. Farahbakhsh S, Kariminik A. Detection of *Listeria Monocytogenes* in non-Pasteurized Milk in Kerman City by Phenotypic and Molecular Techniques. *Journal of Quality and Durability of Agricultural Products and Food Stuffs*. 2021;1(1):30-9. Doi:10.30495/qafj.2021.685560
186. Dehnavi M, Khanjari A, Rezaei E. Prevalence of *Listeria monocytogenes* in traditional cheeses obtained from food sale centers of Tehran, Iran. *Journal of Food Safety and Hygiene*. 2021;7(3):135-40. Doi:10.18502/jfsh.v7i3.9133
187. Faridi Z, Dallal MMS, Koochy-Kamaly P, Javadi NHS, Hadian Z, Fard RMN. Prevalence and Antimicrobial Resistance of *Listeria monocytogenes* in Raw Milk in Tehran, Iran. Doi:10.22037/afb.v8i3.34379
188. Mashak Z, Banisharif F, Banisharif G, Reza Pourian M, Eskandari S, Seif A, et al. Prevalence of *Listeria* species and serotyping of *Listeria monocytogenes* bacteria isolated from seafood samples. *Egyptian Journal of Veterinary Sciences*. 2021;52(1):1-9. Doi:10.21608/EJVS.2020.17893.1105
189. Farhoumand P, Hassanzadazar H, Soltanpour MS, Aminzare M, Abbasi Z. Prevalence, genotyping and antibiotic resistance of *Listeria monocytogenes* and *Escherichia coli* in fresh beef and chicken meats marketed in Zanjan, Iran. *Iranian Journal of Microbiology*. 2020;12(6):537. Doi: 10.18502/ijm.v12i6.5028
190. Soleimani M, Khalili Sadrabad E, Hamidian N, Heydari A, Akrami Mohajeri F. Prevalence and Antibiotic Resistance of *Listeria Monocytogenes* in chicken meat retailers in Yazd, Iran. *Journal of Environmental Health and Sustainable Development*. 2019;4(4):895-902. Doi:10.18502/jehsd.v4i4.2022
191. Akrami-Mohajeri F, Derakhshan Z, Ferrante M, Hamidiyan N, Soleymani M, Conti GO, et al. The prevalence and antimicrobial resistance of *Listeria* spp in raw milk and traditional dairy products delivered in Yazd, central Iran (2016). *Food and Chemical Toxicology*. 2018;114:141-4. Doi:10.1016/j.fct.2018.02.006
192. Lotfollahi L, Chaharbalesh A, Rezaee MA, Hasani A. Prevalence, antimicrobial susceptibility and multiplex PCR-serotyping of *Listeria monocytogenes* isolated from humans, foods and livestock in Iran. *Microbial pathogenesis*. 2017;107:425-9. Doi:10.1016/j.micpath.2017.04.029.
193. Zeinali T, Jamshidi A, Bassami M, Rad M. Isolation and identification of *Listeria* spp. in chicken carcasses marketed in northeast of Iran. *International Food Research Journal*. 2017;24(2):881.
194. Pournajaf A, Rajabnia R, Sedighi M, Kassani A, Moqarabzadeh V, Lotfollahi L, et al. Prevalence, and virulence determination of *Listeria monocytogenes* strains isolated from clinical and non-clinical samples by multiplex polymerase chain reaction. *Revista da Sociedade Brasileira de Medicina Tropi-*

- cal. 2016;49(5):624-7.
195. Abdollahzadeh E, Ojagh SM, Hosseini H, Ghaemi EA, Iradjian G, Heidarloo MN. Antimicrobial resistance of *Listeria monocytogenes* isolated from seafood and humans in Iran. *Microbial pathogenesis*. 2016;100:70-4. Doi:10.1016/j.micpath.2016.09.012.
 196. Rahimi E, Jahanmard M, SAFARI S, Ansari M, TORIKI Z. PREVALENCE AND ANTIMICROBIAL RESISTANCE OF *LISTERIA SPECIES* ISOLATED FROM FILLETED *ARGYROSOMUS HOLOLEPIDOTUS*, *SCOMBEROMORUS COMMERSON* AND *ALBURNUS SPP.* 2016.
 197. Abbasinejad B, Neyriz-Nagadehi M, Taher Talatappeh N. Prevalence and antimicrobial susceptibility of *Listeria monocytogenes* in Koozeh cheeses of Urmia retails. *Food Hygiene*. 2015;5(1 (17)):27-34.
 198. Maktabi S, Pourmehdi M, Zarei M, Moalemi R. Occurrence and antibiotic resistance of *Listeria monocytogenes* in retail minced beef distributed in Ahvaz, South-West of Iran. *Journal of food quality and hazards control*. 2015;2(3):101-6.
 199. Jamali H, Paydar M, Ismail S, Looi CY, Wong WF, Radmehr B, et al. Prevalence, antimicrobial susceptibility and virulotyping of *Listeria species* and *Listeria monocytogenes* isolated from open-air fish markets. *BMC microbiology*. 2015;15(1):144. Doi:10.1186/s12866-015-0476-7.
 200. Haghi F, Zeighami H, Naderi G, Samei A, Roudashti S, Bahari S, et al. Detection of major food-borne pathogens in raw milk samples from dairy bovine and ovine herds in Iran. *Small Ruminant Research*. 2015;131:136-40. Doi:10.1016/j.smallrumres.2015.08.005.
 201. Abdimoghdam Z, Shamloo E, Atefi M. Frequency of *Listeria species* in raw milk and traditional dairy products in Isfahan, Iran. *Iranian Journal of Nutrition Sciences & Food Technology*. 2015;10(3):101-7.
 202. Mansouri-Najand L, Kianpour M, Sami M, Jajarmi M, editors. Prevalence of *Listeria monocytogenes* in raw milk in Kerman, Iran. *Veterinary Research Forum*; 2015: Faculty of Veterinary Medicine, Urmia University, Urmia, Iran.
 203. Rahimi E, Shakerian A. Prevalence of *Listeria Species* in Ready-to-Eat Food in Shahrekord Restaurants. *Medical Laboratory Journal*. 2014;8(2):83-7.
 204. Haj Hosseini A, Sharifan A, Tabatabaee A. Isolation of *Listeria monocytogenes* from meat and dairy products. *Journal of Medical Microbiology and Infectious Diseases*. 2014;2(4):159-62.
 205. Rahimi E, Momtaz H, Behzadnia A, Baghbadorani ZT. Incidence of *Listeria species* in bovine, ovine, caprine, camel and water buffalo milk using cultural method and the PCR assay. *Asian Pacific Journal of Tropical Disease*. 2014;4(1):50-3. Doi:10.1016/S2222-1808(14)60313-3.
 206. Rahimi E. Prevalence and antimicrobial resistance of *Listeria species* isolated from smoked and salted fish. *Iran J Med Microbiol: Volume*. 2014;8(3).
 207. Jamileh N, SIASIE TE, Baharvand R. Isolation of *Listeria monocytogenes* from meat products and study on *prfA* gene in bacteria isolated and comparing with clinical and standard samples. 2013.
 208. Khamesipour F, Khodadoustan Shahraki A, Moumeni M, Khadivi Boroujeni R, Yadegari M. Prevalence of *Listeria monocytogenes* in the crayfish (*Astacus leptodactylus*) by polymerase chain reaction in Iran. *International Journal of Biosciences*. 2013;3(10):160-9. Doi:10.12692/ijb/3.10.160-169.
 209. Sohrabi R, Rashedi M, Dorcheh MP, Shahbazi AM. Prevalence of *Listeria species* in raw milk in Esfahan Province, Iran. *African Journal of Microbiology Research*. 2013;7(19):2057-60. Doi: 10.5897/AJMR12.1463.
 210. Jamali H, Paydar M, Looi CY, Wong WF. Prevalence of *Listeria species* and *Listeria monocytogenes* serotypes in ready mayonnaise salads and salad vegetables in Iran. *African Journal of Microbiology Research*. 2013;7(19):1903-6. Doi: 10.5897/AJMR2013.5658.
 211. Fallah AA, Saei-Dehkordi SS, Mahzounieh M. Occurrence and antibiotic resistance profiles of *Listeria monocytogenes* isolated from seafood products and market and processing environments in Iran. *Food control*. 2013;34(2):630-6. Doi:10.1016/j.foodcont.2013.06.015.
 212. Jamali H, Radmehr B, Thong KL. Prevalence, characterization, and antimicrobial resistance of *Listeria species* and *Listeria monocytogenes* isolates from raw milk in farm bulk tanks. *Food Control*. 2013;34(1):121-5. Doi:10.1016/j.foodcont.2013.04.023.
 213. Akya A, Najafi F, Moradi J, Mohebi Z, Adabagher S. Prevalence of food contamination with *Listeria spp.* in Kerman-shah, Islamic Republic of Iran. *EMHJ-Eastern Mediterranean Health Journal*, 19 (5), 474-477, 2013. 2013.
 214. Momtaz H, Yadollahi S. Molecular characterization of *Listeria monocytogenes* isolated from fresh seafood samples in Iran. *Diagnostic Pathology*. 2013;8(1):1-6. Doi:10.1186/1746-1596-8-149
 215. Dorcheh MP, Sohrabi R, Salajegheh M. Prevalence of *Listeria species* in retail quail products from Isfahan, Iran. *Journal of Veterinary Medicine and Animal Health*. 2013;5(1):16-9. Doi: 10.5897/JVMAH12.019
 216. Khalili Boroujeni F, Moshtaghi H, Bonyadian M. Study on contamination of sheep meat in Shahrekord area with *Listeria ivanovii* and determination its antibiotic resistance pattern.

Iranian Journal of Medical Microbiology. 2013;7(1):15-21.

217. Rahimi E, Yazdi F, Farzinezhadizadeh H. Prevalence and antimicrobial resistance of *Listeria* species isolated from different types of raw meat in Iran. *Journal of food protection*. 2012;75(12):2223-7. Doi:10.4315/0362-028X.JFP-11-565
218. Fallah AA, Saei-Dehkordi SS, Rahnama M, Tahmasby H, Mahzounieh M. Prevalence and antimicrobial resistance patterns of *Listeria* species isolated from poultry products marketed in Iran. *Food Control*. 2012;28(2):327-32. Doi:10.1016/j.foodcont.2012.05.014
219. Rahimi E, Shakerian A, Raissy M. Prevalence of *Listeria* species in fresh and frozen fish and shrimp in Iran. *Annals of Microbiology*. 2012;62(1):37-40. Doi:10.1007/s13213-011-0222-9
220. Modaresi R, Mardani K, Tukmechi A, Ownagh A. Prevalence of *Listeria* spp. in fish obtained from Urmia fish markets. *African Journal of Microbiology Research*. 2011;5(30):5398-401. Doi: 10.5897/AJMR11.888
221. Rahimi E, Ameri M, Momtaz H. Prevalence and antimicrobial resistance of *Listeria* species isolated from milk and dairy products in Iran. *Food Control*. 2010;21(11):1448-52. Doi:10.1016/j.foodcont.2010.03.014
222. Jami S, Jamshidi A, Khanzadi S. The presence of *Listeria monocytogenes* in raw milk samples in Mashhad, Iran. *Iranian Journal of Veterinary Research*. 2010;11(4):363-7.
223. Mahmoodi MM. Occurrence of *Listeria monocytogenes* in raw milk and dairy products in Noorabad, Iran. *Journal of Animal and Veterinary Advances*. 2010;9(1):16-9.
224. Jalali M, Abedi D. Prevalence of *Listeria* species in food products in Isfahan, Iran. *International journal of food microbiology*. 2008;122(3):336-40. Doi:10.1016/j.ijfoodmicro.2007.11.082
- Get rights and content
225. Rahimi E, Momtaz H, Hemmatzadeh F. The prevalence of *Escherichia coli* O157: H7, *Listeria monocytogenes* and *Campylobacter* spp. on bovine carcasses in Isfahan, Iran. *Iranian Journal of Veterinary Research*. 2008;9(4):365-70. Doi: 10.22099/IJVR.2008.2620
226. Shahbazi N, Javadi B, Afshari A, Jamshidi A, Ghavidel M. Identification of *Coxiella Burnetti* and *Mycobacterium* SPP through Touch-down PCR Examination in Unpasteurized Camel Milk in North-East of Iran. *Journal of Nutrition, Fasting & Health*. 2022;10(4). Doi: 10.22038/JNFH.2022.64095.1382.
227. Mekarizadeh K, Ownagh A, Tajik H, editors. Molecular detection of *Coxiella burnetii* in Kope cheese and cattle milk in West Azerbaijan, Iran. *Veterinary Research Forum*; 2023: Faculty of Veterinary Medicine, Urmia University, Urmia, Iran. Doi:10.30466/vrf.2022.544750.3323
228. Mobarez AM, Mostafavi E, Khalili M, Esmaeili S. Identification of *Coxiella burnetii* in raw milk of livestock animal in Iran. *International Journal of Microbiology*. 2021;2021. Doi:10.1155/2021/6632036
229. Kazemeini H, Partovi R, Nazaktabar A, Shokri H. Detection of *Coxiella burnetii* in raw milk samples collected from dairy farms in Mazandaran province, north of Iran. *Journal of Food Safety and Hygiene*. 2022. Doi:10.18502/jfsh.v7i3.9137
230. Nourozghahi NA, RAHIMI E, SHAKERIAN A. Prevalence of *Coxiella burnetii* in bovine bulk milk by Nested-PCR in Gilan province. 2020.
231. Khademi P, Ownagh A, Ataei B, Kazemnia A, Enferadi A, Khalili M, et al. Prevalence of *C. burnetii* DNA in sheep and goats milk in the northwest of Iran. *International Journal of Food Microbiology*. 2020;331:108716. Doi:10.1016/j.ijfoodmicro.2020.108716
232. Esmaeili S, Mobarez AM, Khalili M, Mostafavi E. High prevalence and risk factors of *Coxiella burnetii* in milk of dairy animals with a history of abortion in Iran. *Comparative immunology, microbiology and infectious diseases*. 2019;63:127-30. Doi:10.1016/j.cimid.2019.01.015
233. Khademi P, Ownagh A, Mardani K, Khalili M. Prevalence of *Coxiella burnetii* in milk collected from buffalo (water buffalo) and cattle dairy farms in Northwest of Iran. *Comparative immunology, microbiology and infectious diseases*. 2019;67:101368. Doi:10.1016/j.cimid.2019.101368
234. Reisi M, Rahimi E, Momeni M. Prevalence of *Coxiella burnetii* in Traditional and Industrial Butter and Cream Using Nested Polymerase Chain Reaction in Shahrekord, Iran. *Avicenna Journal of Clinical Microbiology and Infection*. 2019;6(2):61-5. Doi: 10.34172/ajcmi.2019.12
235. Norouzian H, Diali HG, Azadpour M, Afrough P, Shakib P, Mosavi SM, et al. PCR detection of *Coxiella burnetii* in milk samples of ruminants, Iran. *Journal of Medical Bacteriology*. 2018;7(1-2):31-5.
236. Rahmdel S, Sadat Moezzi M, Azimzadeh N, Hosseinzadeh S. PCR Detection of *Coxiella Burnetii* in bovine bulk tank milk samples in Shiraz, southern Iran. *International Journal of Nutrition Sciences*. 2018;3(4):198-201. Doi: 10.30476/IJNS.2019.82644.1022
237. Mosleh N, Moslehisad M, Moosakhani F. Detection of *Coxiella burnetii* by Real-Time PCR in Raw Milk and Traditional Cheese Distributed in Tehran Province. *Journal of Pharmaceutical & Health Sciences*. 2018;6(2):149-55.
238. Lorestani S, Jaydari A, Maleki S, Khademi P. Genomic detection of *Coxiella burnetii* in sheep milk samples by Nest-

- ed-PCR method in Khorramabad, Iran. 2016.
239. Kargar M, Rashidi A, Doosti A, Najafi A, Ghorbani-Dalini S. The sensitivity of the PCR method for detection of *Coxiella burnetii* in the milk samples. *Zahedan Journal of Research in Medical Sciences*. 2015;17(6). Doi:10.17795/zjrms988.
240. Ahmadizadeh C, Moosakhani F, Jamshidian M. Detection and Identification of *Coxiella burnetii* in Milk Cattles of Tehran Province. *Advances in Bioresearch*. 2015;6(4).
241. Khalili M, Diali HG, Mirza HN, Mosavi SM. Detection of *Coxiella burnetii* by PCR in bulk tank milk samples from dairy caprine herds in southeast of Iran. *Asian Pacific Journal of Tropical Disease*. 2015;5(2):119-22. Doi:10.1016/S2222-1808(14)60638-1.
242. Khademi MP. Genomic detection of *Coxiella burnetii* in cattle milk samples by Nested-PCR method, Iran. *Iran J Med Microbiol*: Volume. 2015;9(2).
243. Khanzadi S, Jamshidi A, Razmyar J, Borji S. Identification of *Coxiella burnetii* by touch-down PCR assay in unpasteurized milk and dairy products in North-East of Iran. *Iranian Journal of Veterinary Medicine*. 2014;8(1):15-9.
244. Borji S, Jamshidi A, Khanzadi S, Razmyar J. Detection of *Coxiella burnetii* and sequencing the IS1111 gene fragment in bulk tank milk of dairy herds. *Iranian Journal of Veterinary Science and Technology*. 2014;6(2):21-8. Doi:10.22067/VETERINARY.V6I2.26330.
245. Amin J, Peyman K, Samira L, Shahram M. DEVELOPMENT OF NESTED-PCR METHOD FOR DETECTING COXIELLA BURNETII IN GOAT MILK SAMPLES. *Iranian Journal of Public Health*. 2014;43(2):171.
246. Ghalyanchi Langeroudi A, Babkhani N, Zolfaghari MR, Majidzadeh Arbadili K, Morovvati A, Soleimani M. Detection of *Coxiella brunetii* in bulk tank milk samples from dairy bovine farms using nested-PCR in Qom, Iran, 2011. *Iranian Journal of Veterinary Medicine*. 2013;7(3):207-11.
247. Kargar M, Rashidi A, Doosti A, Ghorbani-Dalini S, Najafi A. Prevalence of *Coxiella burnetii* in bovine bulk milk samples in southern Iran. *Comparative clinical pathology*. 2013;22(3):331-4. Doi:10.1007/s00580-012-1406-9.
248. Rahimi E, Ameri M, Karim G, Doosti A. Prevalence of *Coxiella burnetii* in bulk milk samples from dairy bovine, ovine, caprine, and camel herds in Iran as determined by polymerase chain reaction. *Foodborne pathogens and disease*. 2011;8(2):307-10. Doi:10.1089/fpd.2010.0684.
249. Rahimi E. *Coxiella burnetii* in goat bulk milk samples in Iran. *African Journal of Microbiology Research*. 2010;4(21):2324-6.
250. Rahimi E, Doosti A, Ameri M, Kabiri E, Sharifian B. Detection of *Coxiella burnetii* by nested PCR in bulk milk samples from dairy bovine, ovine, and caprine herds in Iran. *Zoonoses and public health*. 2010;57(7-8):e38-e41. Doi:10.1111/j.1863-2378.2009.01289.x.
251. Toriki Baghbadorani S, Rahimi E, Shakerian A. Investigation of Virulence and Antibiotic-Resistance of *Bacillus cereus* Isolated from Various Spices. *Canadian Journal of Infectious Diseases and Medical Microbiology*. 2023;2023. Doi:10.1155/2023/8390778.
252. Zeighami H, Nejad-dost G, Parsadanians A, Daneshamouz S, Haghi F. Frequency of hemolysin BL and non-hemolytic enterotoxin complex genes of *Bacillus cereus* in raw and cooked meat samples in Zanjan, Iran. *Toxicology reports*. 2020;7:89-92. Doi:10.1016/j.toxrep.2019.12.006.
253. Heydarzadeh M, Javadi A, Rad MG. Detection of emetic toxin (ces) gene and antimicrobial susceptibility of *Bacillus cereus* isolates from Iranian traditional dairy products. 2020. Doi:10.21203/rs.3.rs-61769/v1.
254. Deilami Khiabani Z, Noori E, Rahnama M, Shapouri R, Bigdelo Y, Ghamari D, et al. Detection of NHE Complex Genes in *Bacillus cereus* Isolated from Rice Samples from Zanjan, Iran by Multiplex PCR. *Pathobiology Research*. 2012;14(4):89-97.
255. Heydarzadeh M, Javadi A. Isolation and enumeration of *Bacillus cereus* in raw milk distributed in Tabriz, Iran and detection of ces gene among the isolates. *Food Hygiene*. 2018;8(2(30)):35-42.
256. Ghourchian S, Douraghi M, Baghani A, Soltan Dallal M. *Bacillus cereus* Assessment in Dried Vegetables Distributed in Tehran, Iran. *Journal of food quality and hazards control*. 2018;5(1):29-32. Doi: 10.29252/jfqhc.5.1.29.
257. Hassanzadazar H, Taami B, Abbasi Z, Aminzare M. Microbial Contamination of Cream filled Pastries supplied in Confectioneries of Zanjan, Iran. *Journal of Nutrition, Fasting and Health*. 2018;6(Issue):30-4. Doi:10.22038/jnfh.2018.33470.1119.
258. Soleimani M, Hosseini H, Pilevar Z, Mehdizadeh M, Carlin F. Prevalence, molecular identification and characterization of *Bacillus cereus* isolated from beef burgers. *Journal of Food Safety*. 2018;38(1):e12414. Doi:10.1111/jfs.12414.
259. Ranjbar R, Shahreza MHS. Prevalence, antibiotic-resistance properties and enterotoxin gene profile of *Bacillus cereus* strains isolated from milk-based baby foods. *Tropical Journal of Pharmaceutical Research*. 2017;16(8):1931-7. Doi:10.4314/tjpr.v16i8.25.
260. Soltan Dallal MM, Nezamabadi S, Mardaneh J, Rajabi Z, Sirdani A. Detection of toxigenic *Bacillus cereus* strains in powdered infant formula (PIF) milk by PCR assay. *Tehran University Medical Journal TUMS Publications*. 2017;75(3):179-86.

261. Rasoulpour T, Mahdavi S. Study of frequency of NHE complex genes in *Bacillus cereus* isolated from milk in Tabriz city in spring, 2017. 2020.
262. Molayi Ks, deilami Kz, Ghasemian A, SHAPOURI R, Taghinejad J, Eslami M, et al. Detection of hbla and bal genes in *Bacillus cereus* isolates from cheese samples using the polymerase chain reaction. 2016.
263. Keshtkar M, Momtaz H, Ghavamizadeh M, Rahimi E. Detection of Enterotoxin Genes Profile of *Bacillus cereus* in Pasteurized and Sterile Milk, Baby Food and Dairy Products. *International Medical Journal*. 2016;23(2).
264. Mashak Z, Langroodi AM, Ehsani A, Ilkhanipoor A, Fathabad AE. Microbiological quality of ready-to-eat foods of Tehran province. *African Journal of Food Science*. 2015;9(5):257-61. Doi: 10.5897/AJFS2015.1260.
265. Teymori R, Ghazanfarirad N, Dehghan K, Asadzadeh J, Hajgholizadeh G, Bahmani M. A survey of bacterial and mold contamination of imported rice into West Azerbaijan Province, northwest of Iran. *Asian Pacific Journal of Tropical Disease*. 2014;4:S833-S5. Doi:10.1016/S2222-1808(14)60737-4.
266. Rahimi E, Abdos F, Momtaz H, Torki Baghbadorani Z, Jalali M. *Bacillus cereus* in infant foods: prevalence study and distribution of enterotoxigenic virulence factors in Isfahan Province, Iran. *The Scientific World Journal*. 2013;2013. Doi:10.1155/2013/292571.
267. Hosseini H, Hippe B, Denner E, Kollegger E, Haslberger A. Isolation, identification and monitoring of contaminant bacteria in Iranian Kefir type drink by 16S rDNA sequencing. *Food Control*. 2012;25(2):784-8. Doi:10.1016/j.foodcont.2011.12.017.
268. Rahimifard N, Fatholahzadeh B, Noory Z, Saadati S, Zavar M, Pirouz B, et al. *Bacillus cereus* contamination in infant formula: a study in food and drug control laboratory. *Tehran University Medical Journal TUMS Publications*. 2007;65(8):64-8.
269. Yazdi SS, Mobasser B, Bonab PT, Yazdi SS, Mirbagheri Z, Dallal MMS. Prevalence and Characteristics of *Yersinia enterocolitica* and *Yersinia Pseudotuberculosis* from Raw Milk Supplied in Tehran. *Journal of Nutrition and Food Security*. 2022. Doi:10.18502/jnfs.v8i1.11776.
270. Najjar Asiabani F, Bazaee M, Rais Sadat SA. Study of *Yersinia enterocolitica* contamination in red meat supplied In Shiraz, Iran. *Journal of Zoonotic Diseases*. 2021;5(3):28-33. Doi:10.22034/jzd.2021.13491.
271. Movafagh F, Zeinali T, Jamshidi A. Contamination Rate of Bovine Raw Milk with *Yersinia enterocolitica* Biotypes in Mashhad, Iran. *Journal of Human Environment and Health Promotion*. 2021;7(1):30-4. Doi: 10.29252/jhehp.7.1.5.
272. Tavassoli M, Jamshidi A, Ranjbar G, Torabi MR, Moghddam AA. Prevalence, Biotyping, and Antimicrobial Resistance of *Yersinia enterocolitica* Isolated from Traditional Iranian Cheeses-Evaluation of *Yersinia enterocolitica* in Traditional Cheeses. *City*. 2021;1(1B):5.
273. Tavassoli M, Jamshidi A, Movafagh F, Afshari A. Virulence Characteristics of *Yersinia enterocolitica* Isolated from Dairy Products in the Northeast of Iran. *Journal of Human, Environment and Health Promotion*. 2019;5(2):72-8. Doi:10.29252/jhehp.5.2.5.
274. Sirghani K, Zeinali T, Jamshidi A. Detection of *Yersinia enterocolitica* in retail chicken meat, Mashhad, Iran. *Journal of pathogens*. 2018;2018. Doi:10.1155/2018/1286216.
275. Alavi S, Rahimi E, Tajbakhsh E, Branch S. Prevalence of *Yersinia enterocolitica* in raw small ruminant milk in Shahrekord, Iran. *Bulgarian J Vet Med*. 2018;21(3):364-70. Doi: 10.15547/bjvm.1088.
276. Hadad A, Tajbakhsh E, Reysi M. Identification of virulence genes in serotype O: 3 *Yersinia enterocolitica* isolated from turkey meat by PCR method in Shahrekord. *Journal of Food Microbiology*. 2018;5(2):1-10.
277. Dabiri H, Aghamohammad S, Goudarzi H, Noori M, Heydari MA, Ghoreyshiamiri SM. Prevalence and antibiotic susceptibility of *Campylobacter* species isolated from chicken and beef meat. *International Journal of Enteric Pathogens*. 2016;2(2):6-17087. Doi:10.3168/jds.2014-8853.
278. Jamali H, Paydar M, Radmehr B, Ismail S. Prevalence, characterization, and antimicrobial resistance of *Yersinia* species and *Yersinia enterocolitica* isolated from raw milk in farm bulk tanks. *Journal of dairy science*. 2015;98(2):798-803. Doi:10.3168/jds.2014-8853.
279. Rahimi E, Sepehri S, Dehkordi FS, Shaygan S, Momtaz H. Prevalence of *Yersinia* species in traditional and commercial dairy products in Isfahan Province, Iran. *Jundishapur Journal of Microbiology*. 2014;7(4). Doi: 10.5812/jjm.9249
280. Saberianpour S, Tajbakhsh E, Khamesipour F, Branch S. Prevalence of virulence genes and biotyping of *Yersinia enterocolitica* isolated from chicken meat in Shahrekord, Iran. *Vidyabharati International Interdisciplinary Research Journal*. 2014;3:71-6.
281. Mehdi SDM, Hamid EK, Kazem SYM, Ali TM, Hamid C. Determination Of *Yersinia* Spp. And *Salmonella Paratyphi B* Isolated From Possibly Contaminated Cream Samples In The City Of Tehran. *Payavard Salamat*. 2014;8(1).
282. Momtaz H, Rahimian MD, Dehkordi FS. Identification and characterization of *Yersinia enterocolitica* isolated from raw chicken meat based on molecular and biological techniques. *Journal of Applied Poultry Research*. 2013;22(1):137-45. Doi:10.3382/japr.2012-00549.

283. Hanifian S, Khani S. Prevalence of virulent *Yersinia enterocolitica* in bulk raw milk and retail cheese in northern-west of Iran. *International Journal of Food Microbiology*. 2012;155(1-2):89-92. Doi:10.1016/j.ijfoodmicro.2012.01.012.
284. Mahdavi S, Farshchian MR, Amini K, Rad MAG, Ebadi AR. Survey of *Yersinia enterocolitica* contamination in distributed broiler meats in Tabriz City, Iran. *African Journal of Microbiology Research*. 2012;6(12):3019-23. Doi: 10.5897/AJMR12.094.
285. MK SY, Soltan-Dallal M, Bakhtiari R. Incidence and antibiotic susceptibilities of *Yersinia enterocolitica* and other *Yersinia* species recovered from meat and chicken in Tehran, Iran. *African Journal of Microbiology Research*. 2011;5(18):2649-53. Doi: 10.5897/AJMR11.248.
286. Mirmoeini SS, Sari AA, Goudarztalejerdi A, Alamoti MP, Staji H. Prevalence of *Campylobacter* spp. Among Broiler Carcasses at Industrial Slaughterhouses in Hamedan, Iran. *Archives of Hygiene Sciences*. 2023;12(2). Doi: 10.34172/AHS.12.2.1.393.
287. Mousavinafchi SB, Rahimi E, Shakerian A. *Campylobacter* spp. isolated from poultry in Iran: Antibiotic resistance profiles, virulence genes, and molecular mechanisms. *Food Science & Nutrition*. 2023;11(2):1142-53. Doi:10.1002/fsn3.3152.
288. Emami MS, Shakerian A, Chaleshtori RS, Rahimi E. Distribution of Virulence Genes in *Campylobacter* spp. Isolated from Agaricus Mushrooms in Iran. *BioMed research international*. 2023;2023. Doi:10.1155/2023/1872655.
289. Hadiyan M, Momtaz H, Shakerian A. Prevalence, antimicrobial resistance, virulence gene profile and molecular typing of *Campylobacter* species isolated from poultry meat samples. *Veterinary Medicine and Science*. 2022;8(6):2482-93. Doi:10.1002/vms3.944.
290. Sadeghi A, Ganji L, Fani F, Pouladfar G, Eslami P, Doregirae F, et al. Prevalence, species diversity, and antimicrobial susceptibility of *Campylobacter* strains in patients with diarrhea and poultry meat samples: one-year prospective study. *Iranian Journal of Microbiology*. 2022;14(3):362. Doi: 10.18502/ijm.v14i3.9775.
291. Shahreza MS, Dehkordi NG, Nassar MF, Al-Saedi R. Genotyping of *Campylobacter jejuni* isolates from raw meat of animal species. *Academic Journal of Health Sciences: Medicina balear*. 2022;47(4):52-7. Doi: 10.3306/AJHS.2022.37.04.52.
292. Jahromi RR, Moradi F, Erfanian S, Pourahmadi M. Evaluation of the Contamination of Poultry Carcasses with *Campylobacter jejuni* and *Campylobacter coli* in Southern Iran: A Molecular Study. *Jundishapur Journal of Health Sciences*. 2021;13(3). Doi:10.5812/jjhs.116991.
293. Shafiei A, Rahimi E, Shakerian A. Prevalence, Virulence, and Antimicrobial Resistance of *Campylobacter* Species Isolated From Carcasses of Camels Slaughtered in Slaughterhouses of Chaharmahal and Bakhtiari Province, 2018-2019. *Epidemiology and Health System Journal*. 2021;8(3):115-21. Doi:10.34172/ijer.2021.22.
294. Farhang Zargar M. Identification of *Campylobacter* species in Jahrom slaughterhouse chickens in 2016-17. *Journal of Jahrom University of Medical Sciences*. 2019;17(3):1-6. Doi:10.52547/JMJ.17.3.1.
295. Maktabi S, Ghorbanpoor M, Hossaini M, Motavalibashi A, editors. Detection of multi-antibiotic resistant *Campylobacter coli* and *Campylobacter jejuni* in beef, mutton, chicken and water buffalo meat in Ahvaz, Iran. *Veterinary Research Forum*; 2019: Faculty of Veterinary Medicine, Urmia University, Urmia, Iran. Doi: 10.30466/vrf.2019.34310.
296. Fani F, Aminshahidi M, Firoozian N, Rafaatpour N. Prevalence, antimicrobial resistance, and virulence-associated genes of *Campylobacter* isolates from raw chicken meat in Shiraz, Iran. *Iranian journal of veterinary research*. 2019;20(4):283.
297. Modirrousta S, Shapouri R, Rezasoltani S, Molaabasazadeh H. Prevalence of *Campylobacter* spp. and their Common Serotypes in 330 Cases of Red-meat, Chicken-meat and Egg-shell in Zanjan City, Iran. 2016. Doi: 10.7508/iem.2016.01.003.
298. Hosseinzadeh S, Mardani K, Aliakbarlu J, Ghorbanzadehghan M. Occurrence of *Campylobacter* in chicken wings marketed in the northwest of Iran. *International Food Research Journal*. 2015;22(1).
299. Zendeabad B, Khayatizadeh J, Alipour A. Prevalence, seasonality and antibiotic susceptibility of *Campylobacter* spp. isolates of retail broiler meat in Iran. *Food Control*. 2015;53:41-5. Doi:10.1016/j.foodcont.2015.01.008.
300. Dabiri H, Aghamohammad S, Goudarzi H, Noori M, Ahmadi Hedayati M, Ghoreyshiamiri SM. Prevalence and antibiotic susceptibility of *Campylobacter* species isolated from chicken and beef meat. *Int J Enteric Pathog*. 2014;2(2):1-4.
301. Rahimi E, Sepehri S, Momtaz H. Prevalence of *Campylobacter* species in milk and dairy products in Iran. *Revue de Médecine Vétérinaire*. 2013;164(5):283-8.
302. Kazemeini H, Valizade Y, Parsaei P, Nozarpour N, Rahimi E. Prevalence of *Campylobacter* species in raw bovine milk in Isfahan, Iran. *Middle-East J Sci Res*. 2011;5:664-6.
303. Rahimi E, Ameri M. Antimicrobial resistance patterns of *Campylobacter* spp. isolated from raw chicken, turkey, quail, partridge, and ostrich meat in Iran. *Food Control*. 2011;22(8):1165-70. Doi:10.1016/j.foodcont.2011.01.010.
304. Rahimi E, Kazemeini HR, Safaei S, Allahbakhshi K, Momeni M, Riahi M. Detection and identification of *Campylobacter* spp. from retail raw chicken, turkey, sheep and goat meat

- in Ahvaz, Iran. African journal of microbiology research. 2010;4(15):1620-3.
305. Rahimi E, Ameri M, Kazemeini HR. Prevalence and antimicrobial resistance of *Campylobacter* species isolated from raw camel, beef, lamb, and goat meat in Iran. Food-borne pathogens and disease. 2010;7(4):443-7. Doi:10.1089/fpd.2009.0421.
306. Rahimi E, Tajbakhsh E. Prevalence of *Campylobacter* species in poultry meat in the Esfahan city, Iran. Bulg J Vet Med. 2008;11(4):257-62.
307. Asadi S, Rahimi E, Shakerian A. *Helicobacter pylori* Strains Isolated from Raw Poultry Meat in the Shahrekord Region, Iran: Frequency and Molecular Characteristics. Genes. 2023;14(5):1006. Doi:10.3390/genes14051006.
308. Mashak Z, Jafariaskari S, Alavi I, Shahreza MS, Dehkordi FS. Phenotypic and genotypic assessment of antibiotic resistance and genotyping of *vacA*, *cagA*, *iceA*, *oipA*, *cagE*, and *babA2* alleles of *Helicobacter pylori* bacteria isolated from raw meat. Infection and drug resistance. 2020;13:257. Doi:10.2147/IDR.S233612.
309. Ranjbar R, Yadollahi Farsani F, Safarpour Dehkordi F. Antimicrobial resistance and genotyping of *vacA*, *cagA*, and *iceA* alleles of the *Helicobacter pylori* strains isolated from traditional dairy products. Journal of Food Safety. 2019;39(2):e12594. Doi:10.1111/jfs.12594.
310. Gilani A, Razavilar V, Rokni N, Rahimi E, editors. *VacA* and *cagA* genotypes of *Helicobacter pylori* isolated from raw meat in Isfahan province, Iran. Veterinary Research Forum; 2017: Faculty of Veterinary Medicine, Urmia University, Urmia, Iran.
311. Talimkhani A, Mashak Z. Prevalence and genotyping of *Helicobacter pylori* isolated from meat, Milk and vegetable in Iran. Jundishapur Journal of Microbiology. 2017;10(11). Doi: 10.5812/jjm.14240.
312. Hemmatinezhad B, Momtaz H, Rahimi E. *VacA*, *cagA*, *iceA* and *oipA* genotypes status and antimicrobial resistance properties of *Helicobacter pylori* isolated from various types of ready to eat foods. Annals of clinical microbiology and antimicrobials. 2016;15(1):1-9. Doi:10.1186/s12941-015-0115-z.
313. Saeidi E, Sheikhshahrokh A. *VacA* genotype status of *Helicobacter pylori* isolated from foods with animal origin. BioMed research international. 2016;2016. Doi:10.1155/2016/8701067.
314. Gilani A, Razavilar V, Rokni N, Rahimi E. *VacA* and *cagA* genotypes status and antimicrobial resistance properties of *Helicobacter pylori* strains isolated from meat products in Isfahan province, Iran. Iranian journal of veterinary research. 2017;18(2):97.
315. Ranjbar R, Khamesipour F, Jonaidi-Jafari N, Rahimi E. *Helicobacter pylori* isolated from Iranian drinking water: *vacA*, *cagA*, *iceA*, *oipA* and *babA2* genotype status and antimicrobial resistance properties. FEBS Open Bio. 2016;6(5):433-41. Doi:10.1002/2211-5463.12054.
316. Ghorbani F, Gheisari E, Dehkordi FS. Genotyping of *vacA* alleles of *Helicobacter pylori* strains recovered from some Iranian food items. Tropical Journal of Pharmaceutical Research. 2016;15(8):1631-6. Doi:10.4314/tjpr.v15i8.5.
317. Talaei R, Souod N, Momtaz H, Dabiri H. Milk of livestock as a possible transmission route of *Helicobacter pylori* infection. Gastroenterology and hepatology from bed to bench. 2015;8(Suppl1):S30.
318. Esmaeilgoudarzi D, Tameshkel FS, Ajdarkosh H, Arsalani M, Sohani MH, Behnod V. Prevalence of *Helicobacter pylori* in Iranian milk and dairy products using culture and *ureC* based-PCR techniques. Biomedical and Pharmacology Journal. 2015;8(1):179-83. Doi:10.13005/bpj/597.
319. Atapoor S, Dehkordi FS, Rahimi E. Detection of *Helicobacter pylori* in various types of vegetables and salads. Jundishapur Journal of Microbiology. 2014;7(5). Doi: 10.5812/jjm.10013.
320. Yahaghi E, Khamesipour F, Mashayekhi F, Safarpour Dehkordi F, Sakhaei MH, Masoudimanesh M, et al. *Helicobacter pylori* in vegetables and salads: genotyping and antimicrobial resistance properties. BioMed Research International. 2014;2014. Doi:10.1155/2014/757941.
321. Rahimi E, Kheirabadi EK. Detection of *Helicobacter pylori* in bovine, buffalo, camel, ovine, and caprine milk in Iran. Food-borne Pathogens and Disease. 2012;9(5):453-6. Doi:10.1089/fpd.2011.1060.
322. Ansarian Barezi A, Shakerian A, Rahimi E, Esfandiari Z. Examining the Extent of Contamination, Antibiotic Resistance, and Genetic Diversity of *Clostridioides (Clostridium) difficile* Strains in Meat and Feces of Some Native Birds of Iran. BioMed Research International. 2023;2023. Doi:10.1155/2023/3524091.
323. Ghorbani Filabadi P, Rahimi E, Shakerian A, Esfandiari Z. Prevalence, antibiotic resistance, and genetic diversities of *Clostridium difficile* in meat nuggets from various sources in Isfahan, Iran. Journal of Food Quality. 2022;2022. Doi:10.1155/2022/9919464.
324. Hassani S, Mahmoudi R, Kabudari A, Ghajarbeygi P, Peymani A, Moosavi S, et al. The Prevalence of *Clostridium difficile* and *Clostridium perfringens* in Minced and Ground Beef in Iran. Journal of Chemical Health Risks. 2022;12(4):575-80.
325. Bacheno H, Ahmadi M, Fazeli F, Ariaii P. Antibiotic Resistance Of *Clostridioides Difficile* Isolated From Bovine, Vone And Caprine Raw Meat. Journal of Pharmaceutical Negative Results. 2022;9721-6. Doi: 10.47750/pnr.2022.13.S09.1136.

326. Hassani S, Pakbin B, Brück WM, Mahmoudi R, Mousavi S. Prevalence, antibiotic resistance, toxin-typing and genotyping of *Clostridium perfringens* in raw beef meats obtained from Qazvin city, Iran. *Antibiotics*. 2022;11(3):340. Doi:10.3390/antibiotics11030340.
327. Ram M, Tavassoli M, Ranjbar G, Afshari A. The Microbial and Chemical Quality of Ready-to-Eat Olivier Salad in Mashhad, Iran. *Journal of Nutrition, Fasting and Health*. 2019;7(4):175-81. Doi: 10.22038/jnfh.2019.40572.1199.
328. Hosseinzadeh S, Bahadori M, Poormontaseri M, Dehghani M, Fazeli M, Nazifi S. Molecular characterization of *Clostridium perfringens* isolated from cattle and sheep carcasses and its antibiotic resistance patterns in Shiraz slaughterhouse, southern Iran. *Veterinarski arhiv*. 2018;88(5):581-91. Doi:10.24099/vet.arhiv.0009.
329. Shahdadnejad N, Mohammadabadi M, Shamsadini M. Typing of *clostridium perfringens* isolated from broiler chickens using multiplex PCR. *Genetics in the 3rd millennium*. 2016;14(4):4368-74.
330. Doosti A, Pasand M, Mokhtari-Farsani A, Ahmadi R, Chelgerdi M. Prevalence of *Clostridium perfringens* type a isolates in different tissues of broiler chickens. *Bulgarian Journal of Veterinary Medicine*. 2017;20(1). Doi: 10.15547/bjvm.919.
331. Hajimohammadi B, Dehghani A, Javadzadeh M, Zandi H, Eslami G. *Clostridium botulinum* spores and fungal contamination in honeys of Iran. *Tolooebehdasht*. 2017;15(6):1-9.
332. Shakerian A, Rahimi E, Mesbah J, Mousavi M. Molecular Detection of *Clostridium Perfringens* in Some Raw Animal Food Origin Products in Shahrekord, 2014. *Journal of Payavard Salamat*. 2017;11(4):391-9.
333. Afshari A, Jamshidi A, Razmyar J, Rad M, editors. Genotyping of *Clostridium perfringens* isolated from broiler meat in northeastern of Iran. *Veterinary Research Forum*; 2015: Faculty of Veterinary Medicine, Urmia University, Urmia, Iran.
334. Afshari A, Jamshidi A, Razmyar J, Rad M. Molecular typing of *Clostridium perfringens* isolated from minced meat. *Iranian Journal of Veterinary Science and Technology*. 2015;7(1):32-9. Doi:10.22067/VETERINARY.V7I1.39040.
335. Poormontaseri M, Hosseinzadeh S, Shekarforoush S. Characterization of *Clostridium botulinum* spores and its toxin in honey. *Iranian Journal of Veterinary Research*. 2014;15(1):36-9.
336. Sadeghi Sarvestani MV, Hosseinzadeh S, Poormontaseri M, Fazeli M. Monitoring the Various Types of *Clostridium botulinum* in Four Kinds of Food Stuffs Using Multiplex PCR. *Journal of Fasa University of Medical Sciences*. 2014;3(4):380-6.
337. Shahcheraghi F, Nobari S, Asl HM, Aslani MM. Identification of botulinum toxin type in clinical samples and foods in Iran. *Archives of Iranian medicine*. 2013;16(11):0-.
338. Hosseini H, Tavakoli HR, Meshgi MA, Khaksar R, Hosseini M, Khakpour M. Survey of *Clostridium botulinum* toxins in Iranian traditional food products. *Comparative clinical pathology*. 2010;19(3):247-50. Doi:10.1007/s00580-009-0865-0.
339. Majzobi MM, Karami P, Khodavirdipour A, Alikhani MY. Brucellosis in humans with the approach of brucella species contamination in unpasteurized milk and dairy products from Hamadan, Iran. *Iranian Journal of Medical Microbiology*. 2022;16(4):282-7.
340. Asgari MH, Ahmadi E. Contamination of bovine milk with *Brucella* spp.: a current public health menace in Kurdistan province of Iran. *Avicenna Journal of Clinical Microbiology and Infection*. 2021;8(1):17-22. Doi: 10.34172/ajcmi.2021.04.
341. Majzobi MM, Karami P, Khodavirdipour A, yousef Alikhani M. The contamination rate of *Brucella* spp. in unpasteurized dairy products used in urban and rural areas of Hamadan County, west of Iran. 2020. Doi:10.21203/rs.2.22703/v1.
342. Alamian S, Dadar M. *Brucella abortus* contamination of camel milk in two Iranian regions. *Preventive veterinary medicine*. 2019;169:104708. Doi:10.1016/j.prevetmed.2019.104708.
343. Mirnejad R, Jazi FM, Mostafaei S, Sedighi M. Molecular investigation of virulence factors of *Brucella melitensis* and *Brucella abortus* strains isolated from clinical and non-clinical samples. *Microbial pathogenesis*. 2017;109:8-14. Doi:10.1016/j.micpath.2017.05.019.
344. Shirazi Z, Khalili M, Sadeghi B, Sharifi H, Ashrafganjooyi S. Detection of *Brucella* spp. in the sheep and goats milks from southeastern Iran using culture and PCR. *Journal of Medical Microbiology and Infectious Diseases*. 2017;5(3):40-2.
345. Ashrafganjooyi S, Saedadeli N, Alamian S, Khalili M, Shirazi Z. Isolation and biotyping of *Brucella* spp. from sheep and goats raw milk in southeastern Iran. *Trop Biomed*. 2017;34(3):507-11.
346. Majid Y, Somayeh N, Parisa S, Behrooz A, Reza FN, Javad R, et al. Prevalence of *Brucella melitensis* and *Brucella abortus* in raw milk and dairy product by real time PCR technique. *The Ulutas Medical Journal*. 2016;2(1):7-11. Doi:10.5455/umj.20151004054235.
347. Khalili M, Aflatoonian MR, Aliabadi FS, Abshenas J. *Brucella* contamination in raw milk by polymerase chain reaction. *Tehran University Medical Journal TUMS Publications*. 2016;74(7):517-21.
348. Shakerian A, Deo P, Rahimi E, Shahjavan A-R, Khamesipour F. Molecular detection of *Brucella melitensis* in sheep and goat milk in Iran. *Tropical Journal of Pharmaceutical Research*. 2016;15(5):913-8. Doi:10.4314/tjpr.v15i5.3.

349. Izadi A, Moslemi E, Tabatabaei Panah A, Kheiri Manjili H. *Brucella* spp. detection in dairy products using nested and hemi nested PCR techniques. *Ann Biol Res.* 2014;5(1):124-31.
350. Gholizadeh SS, Zali M, Hashempour A, AHMADI EM. Investigation of brucellosis in cattle and sheep in Urmia-Iran. *Yüzüncü yıl Üniversitesi Veteriner Fakültesi Dergisi.* 2013;24(3):133-4.
351. Karimi Alavigh A, Sharifzadeh A. Study on the Frequency of *Vibrio* species in fish from Isfahan. *Journal of Food Microbiology.* 2021;8(4):47-55.
352. Shohreh P, Azizkhani M, Tooryan F. Food Poisoning Bacteria from Commercially Important Fish Species Collected from Markets in Mazandaran Province. *Fisheries Science and Technology.* 2019;8(1):1-6.
353. Asgarpoor D, Haghi F, Zeighami H. Detection and molecular characterization of *Vibrio parahaemolyticus* in shrimp samples. *The Open Biotechnology Journal.* 2018;12(1).
354. Mashak Z, Khalajzadeh A, Koozdar V. Study the bacterial and fungal quality and physicochemical properties of cold smoked salted fishes of Caspian sea, Iran. *Biosciences Biotechnology Research Asia.* 2016;13(3):1811-20. Doi: 10.13005/bbra/2334.
355. Raissy M, Rahimi E, Azargun R, Moumeni M, Sohrabi H. Molecular detection of *Vibrio* spp. in fish and shrimp from the Persian Gulf. *Journal of Food Biosciences and Technology.* 2015;5(2):49-52.
356. Hosseini S, Safarpour Dehkordi F, Rahimi E, Shakerian A. Prevalence study of *Vibrio* species and frequency of the virulence genes of *Vibrio parahaemolyticus* isolated from fresh and salted shrimps in Genaveh seaport. *Food Hygiene.* 2014;4(2 (14)):17-26.
357. Khamesipour F, Noshadi E, Moradi M, Raissy M. Detection of *Vibrio* spp. in shrimp from aquaculture sites in Iran using polymerase chain reaction (PCR). *Aquaculture, Aquarium, Conservation & Legislation.* 2014;7(1):1-7.
358. Dehkordi FS, Hosseini S, Rahimi E, Momeni M, Yahaghi E, Darian EK. Study the distribution of virulence genes of *Vibrio parahaemolyticus* isolated from fish, lobster and crab caught from Persian Gulf. *Iran J Med Microbiol: Volume.* 2014;8(2).
359. Raissy M, Khamesipour F, Rahimi E, Khodadoostan A. Occurrence of *Vibrio* spp., *Aeromonas hydrophila*, *Escherichia coli* and *Campylobacter* spp. in crayfish (*Astacus leptodactylus*) from Iran. *Iranian Journal of Fisheries Sciences.* 2014;13(4):944-54.
360. Momtaz H, Dehkordi FS, Rahimi E, Asgarifar A. Detection of *Escherichia coli*, *Salmonella* species, and *Vibrio cholerae* in tap water and bottled drinking water in Isfahan, Iran. *BMC public health.* 2013;13(1):1-7. Doi:10.1186/1471-2458-13-556.
361. Raissy M, Moumeni M, Ansari M, Rahimi E. Occurrence of *Vibrio* spp. in lobster and crab from the Persian Gulf. *Journal of Food Safety.* 2012;32(2):198-203. Doi:10.1111/j.1745-4565.2012.00368.x.
362. Rahimi E, Ameri M, Doosti A, Gholampour AR. Occurrence of toxigenic *Vibrio parahaemolyticus* strains in shrimp in Iran. *Foodborne pathogens and disease.* 2010;7(9):1107-11. Doi:10.1089/fpd.2010.0554.
363. Hosseini H, Cheraghali AM, Yalfani R, Razavilar V. Incidence of *Vibrio* spp. in shrimp caught off the south coast of Iran. *Food control.* 2004;15(3):187-90. Doi:10.1016/S0956-7135(03)00045-8.
364. Pakbin B, Didban A, Brück WM, Alizadeh M. Phylogenetic analysis and antibiotic resistance of *Shigella sonnei* isolates. *FEMS microbiology letters.* 2022;369(1):fnac042. Doi:10.1093/femsle/fnac042.
365. Pakbin B, Amani Z, Allahyari S, Mousavi S, Mahmoudi R, Brück WM, et al. Genetic diversity and antibiotic resistance of *Shigella* spp. isolates from food products. *Food Science & Nutrition.* 2021;9(11):6362-71. Doi:10.1002/fsn3.2603.
366. Pakbin B, Didban A, Monfared YK, Mahmoudi R, Peymani A, Modabber MR. Antibiotic susceptibility and genetic relatedness of *Shigella* species isolated from food and human stool samples in Qazvin, Iran. *BMC research notes.* 2021;14(1):1-6. Doi:10.1186/s13104-021-05554-3.
367. Tehrani ST, Harzandi N, Jabalameli L. Molecular detection of *Shigella* spp. contamination in ready-to-eat salad samples in west of Tehran. *International Journal of Enteric Pathogens.* 2018;6(2):41-4. Doi: 10.15171/ijep.2018.11.
368. Bahador N, Ajampour I. Phenotypic and genotypic characterization of isolated *Shigella* spp. from food based on IpaH gene and evaluation of their antibiotic patterns. *Journal of Food Microbiology.* 2019;5(4):55-65.
369. Yam BAZ, Khomeiri M, Mahounak AS, Jafari SM. Hygienic quality of camel milk and fermented camel milk (chal) in Golestan Province, Iran. *Journal of Microbiology Research.* 2014;4(2):98-103. Doi: 10.5923/j.microbiology.20140402.09.

COPYRIGHTS

©2024 The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited. No permission is required from the authors or the publishers.

**How to cite this article**

Hashemi M, Asadi Touranlou F, Adibi SH, Afshari A, Shakeri G. Foodborne Bacteria in Iran: A 23-year Systematic Review of high-risk foods. *Iran J Vet Sci Technol.* 2024; 16(3): 1-47.
DOI: <https://doi.org/10.22067/ijvst.2024.87069.1357>
https://ijvst.um.ac.ir/article_45491.html