

## Investigation of *Chlamydia abortus* infection in aborted fetuses using molecular and pathological studies in East Azerbaijan province, northwest Iran

**Authors:** Monireh Khordadmehr<sup>a,b\*</sup>, Hassan Sadri<sup>c,b</sup>, Jafar Shirazi<sup>d,b</sup>, Saeed Babazadeh<sup>d</sup>, Farinaz Jigari-Asl<sup>a</sup>, Katayoon Nofouzi<sup>a,b</sup>, Yaser Jafari-Khataylou<sup>a,b</sup>, Faeghesadat Mousavi<sup>1a</sup>, Abolfazl Hajibemani<sup>c,b</sup>

<sup>a</sup> Department of Pathobiology, Faculty of Veterinary Medicine, University of Tabriz, 516665-1647, Tabriz, Iran

<sup>b</sup> [Abortion Research Group](#), Faculty of Veterinary Medicine, University of Tabriz, 516665-1647, Tabriz, Iran

<sup>c</sup> Department of Clinical Sciences, Faculty of Veterinary Medicine, University of Tabriz, 516665-1647, Tabriz, Iran

<sup>d</sup> Veterinary Organization, East Azerbaijan Province, Tabriz, Iran

\*Corresponding author: Dr. Monireh Khordadmehr

Postal address: Faculty of Veterinary Medicine, University of Tabriz, 516665-1647, Tabriz, Iran

University/organization email address: [khordadmehr@tabrizu.ac.ir](mailto:khordadmehr@tabrizu.ac.ir)

ORCID ID: <http://orcid.org/0000-0002-4472-3847>

Tel. number: [09142274973](tel:09142274973)

### Abstract

*Chlamydia abortus* is an important cause of in-utero infections in sheep and goats, resulting in abortion, stillbirth, and the birth of weak offspring. The disease in sheep caused by *Chlamydia abortus* (*C. abortus*) has been known as “ovine enzootic abortion”. Notably, it is recognized as a zoonotic disease, which can lead to abortion in humans. Therefore, the present study was carried out to recognize *Chlamydia* infection in aborted fetuses of domestic small ruminants, including sheep and goats in East Azerbaijan province, northwest Iran. For this purpose, a total of 62 aborted fetuses were obtained from sheep and goat flocks. At necropsy, the fetus was usually well preserved with few gross lesions. The tissue samples were collected for

histopathology and molecular studies. The conventional PCR method using specific primers was performed for detection of the *Chlamydia* genome. Additionally, the formalin-fixed tissue samples were routinely processed for histopathological studies. The genome of *C. abortus* was detected in 33.87% (95%CI: 0.32 ± 0.11) of the examined fetuses. Histopathological examinations presented multifocal hepatitis, pneumonia, and nonsuppurative meningoencephalitis associated with focal hemorrhage in the muscles. In conclusion, the investigation of the *C. abortus* genome in aborted fetuses with high prevalence rates indicates that this infection can play a notable role in the abortion of sheep and goats in East Azerbaijan. To prevent potential abortions in women who are in close contact with aborting ruminants, effective management and control measures for public health in the region are crucial.

**Keywords:** Sheep, Goat, Abortion, Zoonotic infection, Public health

**Abbreviations:** OEA: Ovine Enzootic Abortion; PCR: polymerase chain reaction; H&E: hematoxylin and eosin; CI: confidence interval.

## Introduction

Abortion is a significant problem in small ruminants, particularly in sheep and goats, and it is responsible for major economic losses in the livestock industry due to the loss of fetuses. The cause of abortion can be infectious which is the most common or non-infectious [1]. Infectious causes can be bacterial, viral, fungal, and parasitic [2, 3]. Bacterial causative agents can be a more serious issue in small ruminants such as *Chlamydia abortus* [4]. One of the four families that make up the order *Chlamydiales* is the *Chlamydiaceae* family. At first, the family was divided into two genera, *Chlamydia* and *Chlamydophila*, but recent genomic and phylogenetic studies have resulted in the unification of all species under the genus *Chlamydia* [5, 6]. Besides the species currently known for the genus *Chlamydia* which includes 21 different species in various range hosts [7-11]. Among these species, some of them such as *C. psittaci* and *C.*

*abortus*, are known to cause zoonotic infections. *C. abortus* is a major cause of chlamydiosis in small ruminants, which can lead to reproductive problems such as abortions, stillbirths, and infertility [12, 13]. *C. abortus* in sheep is also known as Ovine Enzootic Abortion (OEA) or Enzootic Abortion of Ewes (EAE) [12, 13].

*Chlamydia* is transmitted through multiple routes, including aerosol transmission, horizontal, ingestion, and fecal-oral. Fomites and environmental agents can also be important in spreading the infection. These multiple transmission pathways are involved in making this zoonotic infection more serious [14]. Oronasal route has been considered as the primary way of transmission and it can happen from direct contact between animals or with abortion remnants. Dissemination of *C. abortus* to the placenta happens during pregnancy triggers inflammation and placental insufficiency, which leads to abortion or stillbirth [15]. The pathological lesions are detectable after day 90 of gestation and the first lesion can be seen as cytoplasmic chlamydial inclusions produced in the cells of cotyledon [16]. The lesions in the fetus can include primarily focal necrosis in the liver and additionally, small necrotic areas in the lungs, and spleen, and less commonly in the brain and lymph nodes, and sometimes a mild form of semipurulent focal interstitial nephritis in fetal kidneys [16].

The PCR methods are frequently used for detecting *chlamydia* species due to their high specificity and sensitivity, quick results, and ability to handle large volumes [17]. Due to the presence of the organism in different organs such as the abomasum and jejunum, as well as the possibility of the fetus swallowing the infected amniotic fluid, *chlamydia* can be diagnosed using the abomasal content of aborted fetuses [18]. This study was conducted to detect *Chlamydia* infection in the aborted fetuses of sheep and goats flocks in east Azerbaijan province (Iran) using conventional PCR method, due to the high prevalence of infectious abortion in the region. Also, it investigated the main histopathological changes associated with this infection to improve diagnostic accuracy.

## Results

### *Molecular findings*

The molecular study results associated with the age groups are presented in Figure 1 and Table 1. Briefly, the genome of *C. abortus* was detected in 21 out of 62 samples (33.87%, 95%CI: 0.32 ± 0.11) from the examined fetuses. Of note, the higher infection rates were found in fetuses aged 4 to 5 months. However, the four age groups had no significant difference in infection rates.

**Table 1.**

PCR results for detection of *Chlamydia* in the aborted fetuses (N = 62).

Age groups	No. of the positive samples (%)	95% CI
2-3 m 5/62 (8.06%)	0/5 (0%)	0.0±0.0
3-4 m 14/62 (22.58%)	2/14 (14.28%)	0.1428±0.1813
4-5 m 36/62 (58.06%)	17/36 (47.22%)	0.472±0.162
5-6 m 7/62 (11.29%)	2/7 (28.57%)	0.2857±0.3242
Total	21/62 (33.87%)	0.3387±0.114

### *Pathological findings*

There were focal areas of cellular necrosis in the liver and spleen, which were variably surrounded by mononuclear cells. An increase in mononuclear cells was observed throughout the liver and concentrated in portal areas. In the lung, alveolar septa were thickened by mononuclear cells associated with hyperemia of the pulmonary vessels. Additionally, there were mild to moderate hemorrhages in the muscles and spleen. In the brain, mild meningoencephalitis accompanied by hyperemia and focal hemorrhage were found.

## Discussion

In this study, out of the 62 samples, 21 (33.87%) tested positive for *Chlamydia* infection using *Chlamydia* genus-specific primers for conventional PCR. In pathologic studies, the main lesions of *Chlamydia* infection were observed in the examined aborted fetuses. Due to the importance of *Chlamydia* in the discussion of abortion in small ruminants and also zoonotic aspects, various studies have been conducted within Iran to estimate the prevalence of this infection. Based on some studies conducted in Iran from 2015 to 2023, the prevalence of *Chlamydia* infection in individual animals varied between 5.71% and 56.41%. This variation was attributed to the differences in geographical areas, the detection methods used, and the types of samples taken from the animals. For example, a previous study found a seroprevalence of 25.6% using the ELISA test in individual animals across seven provinces in Iran [19]. In contrast, another study reported a higher prevalence of 56.41% in samples of aborted material from sheep and cattle in Shahr-e-Kord and Bagh-e-Malek, using *Chlamydiales* order-specific primers in PCR testing [20]. A recent study used cell culture for *chlamydia* detection, which differs from methods used in other studies. Chlamydial inclusion bodies were seen with an optical microscope in 14.28% of samples [21]. In contrast, other researchers reported a significantly lower prevalence of 5.71% as evaluated by an ELISA assay among 16 non-vaccinated goat flocks in Khuzestan province [22]. In another ELISA assay investigation, 9.7% of the samples were positively reported for sheep and goats in Khorasan Razavi province [23]. Moreover, the prevalence of *Chlamydia* infection in vaginal and ocular swab samples of small ruminants was 24.1% by real-time PCR [24]. Besides, the results of the nested-PCR of the milk samples showed 8.61% of milk samples were infected [25]. The seroprevalence of chlamydiosis in sera samples from sheep and goats with a history of abortion was 4.3% in sheep and 21.7% in goats, respectively, as determined by the indirect ELISA assay in Fars province [26]. Also, it was reported a prevalence of 21.6% in vaginal swab samples of sheep

and goats by PCR method in Kerman province [27]. In studies in which samples were from aborted fetuses, similar variability in prevalence was reported. In this regard, evidence indicates infection rates of 0%, 11%, and 23.5% in aborted fetuses of small ruminants from southeast [28], south [29], and northwest [30] Iran, respectively, as determined by the PCR method. Studies in other countries have also reported different prevalences. In Sulaimani province, Iraq, a low prevalence of 3.33% was detected in aborted fetus samples of sheep [31]. While in northwestern Italy, a higher seroprevalence was observed, with 56.6% of sheep and goat flocks testing positive for *C. abortus* antibodies. Sheep flocks had a higher prevalence of 71.4%, compared to 44.8% in goat flocks [32]. A study conducted in Africa found a much lower prevalence of 2% for *C. abortus* in small ruminants using an indirect ELISA assay [33]. This shows the correlation between prevalence and geographical differences. Emerging evidence shows that the global prevalence of *C. abortus* is 20.1% in sheep and 14.4% in goats, with notable regional differences. The highest prevalence has been reported in South Asia, at 30.6%, while East Asia and Oceania have the lowest rate at 14%. European countries like Romania, Hungary, and Germany reported high prevalences, ranging from 53.3% to 87%. In contrast, lower rates were found in Costa Rica, Australia, and Zimbabwe, ranging between 4.7% and 5.2% [34]. In *Chlamydia* infection, abortion is likely the result of several factors, including tissue destruction by *Chlamydia*, vasculitis, thrombosis, and a fetal inflammatory response, such as the production of TNF- $\alpha$  by fetal macrophages [16]. In the present study, the main pathological lesions included focal necrotic hepatitis, mild interstitial pneumonia, and mild meningoencephalitis associated with focal hemorrhage in the muscles, spleen, and brain. Similar pathological findings previously described by others [16], are in agreement with our results.

## **Conclusion**

In conclusion, the prevalence of *Chlamydia* infection in small ruminants shows significant variation across different regions, affected by other factors such as management practices and environmental conditions. Studies have reported prevalence rates ranging from as low as 2% in Africa to over 70% in some flocks in Italy, with significant regional differences existing globally. PCR and ELISA are common methods used for detecting infections in most studies, each with its benefits. The type of samples collected, like sera, aborted fetuses, and milk, also affects detection. Our study showed that 33.87% of the samples tested positive for *Chlamydia* using genus-specific primers in PCR. This significant prevalence highlights the need for more effective strategies to manage this infection and emphasizes the importance of increasing efforts toward prevention in the region for public health.

## **Materials and Methods**

### ***Study area***

The present study was performed in seven cities in the East-Azerbaijan province in northwest Iran, including Tabriz, Charuymaq, Khoda Afrin, Jolfa, Heris, Bostan Abad, and Mianeh. This study presents findings on *Chlamydia* infection as part of a larger investigation into the infectious and non-infectious causes of abortion in small ruminants in East Azerbaijan province, northwest Iran. For this purpose, from November 2023 to February 2024, a total of 62 aborted fetuses were collected from sheep and goats flocks in the mentioned regions, as the owners had contacted for abortions on their farms. We studied a total of 43 sheep flocks, documenting the history of the sampled herds, which included details such as herd size and abortion rate. The sheep flocks studied were categorized by size: 7 small flocks (1–100 sheep), 19 medium flocks (101–300 sheep), and 17 large flocks (over 300 sheep).

### ***Ethical approval***

All relevant international, national, and institutional guidelines for the care and use of animals, including the protocol approved by the University of Tabriz's Animal Research Ethics Committee (ID: IR.TABRIZU.REC.1403.049), were followed.

### ***Sampling***

All samples of dead fetuses belonged to the herds with the traditional conditions. At first, the age of the aborted fetuses was estimated using the formula  $(X + 17) \times 1/2$ , where X is the size of the fetus in centimeters, which were measured from forehead to tail. Then, a systematic necropsy was performed and the pathological lesions were recorded. Next, 50 mg of the abomasal contents was placed in a 2 mL microtube and stored in a freezer at -70 °C for further molecular studies. Additionally, tissue samples of the various organs including the liver, kidney, lung, muscles, spleen, and brain were collected and transferred to 10% formalin solution for histopathology purposes.

### ***Pathological study***

The tissue samples were kept in a 10% neutral buffered formalin solution for at least 48 hours. After that, they were routinely using a DS2080/H tissue processor (Didsabz, Iran), embedded in paraffin, and cut into 5 µm thick sections. These sections were stained with common hematoxylin and eosin (H&E) and studied under a light microscope (Olympus, CH-30, Japan), with the observed lesions recorded.

### ***Molecular studies (DNA extraction and PCR assay)***

The genomic DNA (gDNA) of the abomasal contents was extracted, using a DNA extraction kit® (Pishgam Sanjesh, Tehran, Iran) based on the manufacturer's instructions. The genome's quality and quantity were analyzed using NanoPhotometer® NP80 (IMPLEN, Germany). All PCR assays were performed in a final volume of 25 µL with Taq DNA Polymerase Master Mix

RED® (Ampliqon, Denmark) and 3 µL DNA/cDNA, using the T100 Thermal Cycler (Bio-Rad, USA). The amplified products were detected through electrophoresis on 1.5% agarose gels stained with a safe DNA stain (SinaClon, Iran). To perform a PCR test, specific primers for *Chlamydia abortus* were used: CHLAMA1 (Forward: 5'-CTCACCATTTGTCTCAGGTGGA-3') and CHLAMA2 (Reverse: 5'-ACCGTAATGGGTAGGAGGGGT-3'), targeting an 823 base pair (bp) sequence. Moreover, *C. abortus* ATCC VR656 was used as positive control. The reaction conditions included 35 cycles and an annealing temperature of 59°C.

### ***Statistical analyses***

The Chi-Square test was used to determine the correlations between infections and age groups (four groups, including 2-3, 3-4, 4-5, and over 5 months old) of the fetuses. Differences were considered significant at  $P < 0.05$ . The analyses were performed with IBM SPSS Statistics v.22 software. Also, the data was assessed using a 95% confidence interval (CI).

### **References**

1. Holler LD. Ruminant abortion diagnostics. *Veterinary Clinics of North America-Food Animal Practice*. 2012;28(3):407. Doi: 10.1016/j.cvfa.2012.07.007.
2. Pal M. Growing role of fungi in mycotic abortion of domestic animal. *Journal of Bacteriology and Mycology*. 2015;2(1):1009.
3. Haif A, Khelifi-Ouchene NA, Khelifi M, Ouchetati I, Zeroual F, Ouchene N. Abortive diseases and their various associated risk factors in small ruminants in Algeria: a systematic review. *Tropical Animal Health and Production*. 2021;53(6):520. Doi: 10.1007/s11250-021-02926-6.

4. Schnydrig P, Vidal S, Brodard I, Frey C, Posthaus H, Perreten V, et al. Bacterial, fungal, parasitological and pathological analyses of abortions in small ruminants from 2012–2016. *Schweizer Archiv fur Tierheilkunde*. 2017;159:647-56. Doi: 10.17236/sat00136.
5. Sachse K, Bavoil PM, Kaltenboeck B, Stephens RS, Kuo C-C, Rosselló-Móra R, et al. Emendation of the family Chlamydiaceae: proposal of a single genus, Chlamydia, to include all currently recognized species. *Systematic and Applied Microbiology*. 2015;38(2):99-103. Doi: 10.1016/j.syapm.2014.12.004.
6. Vorimore F, Hölzer M, Liebler-Tenorio E, Barf L-M, Delannoy S, Vittecoq M, et al. Evidence for the existence of a new genus Chlamydiifrater gen. nov. inside the family Chlamydiaceae with two new species isolated from flamingo (*Phoenicopterus roseus*): *Chlamydiifrater phoenicopteri* sp. nov. and *Chlamydiifrater volucris* sp. nov. *Systematic and Applied Microbiology*. 2021;44(4):126200. Doi: 10.1016/j.syapm.2021.126200.
7. Laroucau K, Vorimore F, Aaziz R, Solmonson L, Hsia R, Bavoil P, et al. *Chlamydia buteonis*, a new *Chlamydia* species isolated from a red-shouldered hawk. *Systematic and Applied Microbiology*. 2019;42(5):125997. Doi: 10.1016/j.syapm.2019.06.002.
8. Chaiwattananrungruengpaisan S, Thongdee M, Anuntakarun S, Payungporn S, Arya N, Punchukrang A, et al. A new species of *Chlamydia* isolated from Siamese crocodiles (*Crocodylus siamensis*). *PLoS One*. 2021;16(5):e0252081. Doi: 10.1371/journal.pone.0252081.
9. Laroucau K, Ortega N, Vorimore F, Aaziz R, Mitura A, Szymanska-Czerwinska M, et al. Detection of a novel *Chlamydia* species in captive spur-thighed tortoises (*Testudo graeca*) in southeastern Spain and proposal of *Candidatus Chlamydia testudinis*. *Systematic and Applied Microbiology*. 2020;43(2):126071. Doi: 10.1016/j.syapm.2020.126071.

10. Mitura A, Niemczuk K, Zaręba K, Zając M, Laroucau K, Szymańska-Czerwińska M. Free-living and captive turtles and tortoises as carriers of new *Chlamydia* spp. *PLoS One*. 2017;12(9):e0185407. Doi: 10.1371/journal.pone.0185407.
11. Bellinati L, Pesaro S, Marcer F, Danesi P, Natale A, Ceglie L. Detection of a novel chlamydia species in invasive turtles. *Animals*. 2022;12(6):784. Doi: 10.3390/ani12060784.
12. Merdja SE, Khaled H, Aaziz R, Vorimore F, Bertin C, Dahmani A, et al. Detection and genotyping of *Chlamydia* species responsible for reproductive disorders in Algerian small ruminants. *Tropical Animal Health and Production*. 2015;47:437-43. Doi: 10.1007/s11250-014-0743-x.
13. Rodolakis A, Laroucau K. Chlamydiaceae and chlamydial infections in sheep or goats. *Veterinary Microbiology*. 2015;181(1-2):107-18. Doi: 10.1016/j.vetmic.2015.07.010.
14. Nietfeld JC. Chlamydial infections in small ruminants. *Veterinary Clinics of North America: Food Animal Practice*. 2001;17(2):301-14. Doi: 10.1016/s0749-0720(15)30030-x.
15. Borel N, Sachse K. Zoonotic transmission of *Chlamydia* spp.: known for 140 years, but still underestimated. *Zoonoses: Infections Affecting Humans and Animals*. 2023; 1-28. Doi:10.1007/978-3-030-85877-3\_53-1.
16. Buxton D, Barlow R, Finlayson J, Anderson I, Mackellar A. Observations on the pathogenesis of *Chlamydia psittaci* infection of pregnant sheep. *Journal of Comparative Pathology*. 1990;102(2):222-37.
17. Pantchev A, Sting R, Bauerfeind R, Tyczka J, Sachse K. Detection of all *Chlamydoxila* and *Chlamydia* spp. of veterinary interest using species-specific real-time PCR assays. *Comparative Immunology, Microbiology and Infectious Diseases*. 2010;33(6):473-84. Doi: 10.1016/j.cimid.2009.08.002.
18. Menzies PI. Abortion in sheep: diagnosis and control. *Current Therapy In Large Animal Theriogenology*: Elsevier; 2007. p. 667-80. Doi:10.1016/B978-072169323-1.50093-3

19. Esmaeili H, Bolourchi M, Mokhber-Dezfouli MR. Seroprevalence of Chlamydia abortus infection in sheep and goats in Iran. *Iranian Journal of Veterinary Medicine*. 2015;9(2):73-77.
20. Barati S, Moori-Bakhtiari N, Najafabadi MG, Momtaz H, Shokuhizadeh L. The role of zoonotic chlamydial agents in ruminants abortion. *Iranian Journal of Microbiology*. 2017;9(5):288.
21. Esmaeili H, Hamed M, Madani S. Isolation of Chlamydia spp. from Ewes and Does in Iran. *Archives of Razi Institute*. 2017;72(4):249-53. Doi: 10.22092/ari.2017.113297.
22. Borujeni MP, Bakhtiari NM, Hajikolaei M, Mousavi M. Chlamydia abortus infection in goats in the southwest of Iran. *Revue de Médecine Vétérinaire*. 2019;170:9-14.
23. Iraninezhad Z, Azizzadeh M, Taghavi Razavizadeh A, Mehrzad J, Rashtibaf M. Seroepidemiological feature of Chlamydia abortus in sheep and goat population located in northeastern Iran. *Veterinary Research Forum*. 2020;11(4):423-426. Doi: 10.30466/vrf.2019.101946.2429.
24. Esmaeili H, Bolourchi M, Mokhber-Dezfouli MR, Teimourpour A. Detection of Chlamydia abortus and risk factors for infection in small ruminants in Iran. *Small Ruminant Research*. 2021;197:106339. Doi:10.1016/j.smallrumres.2021.106339.
25. Taheri F, Ownagh A, Mardani K. Molecular identification and phylogenetic analysis of Chlamydia abortus isolated from sheep and goats. *Iranian Journal of Veterinary Science and Technology*. 2021;13(1):55-63. Doi: 10.22067/ijvst.2021.68614.1016.
26. Abnaroodheleh F, Emadi A, Dadar M. Seroprevalence of brucellosis and chlamydiosis in sheep and goats with history of abortion in Iran. *Small Ruminant Research*. 2021;202:106459. Doi: 10.1016/j.smallrumres.2021.106459.
27. Afrisham S, Golchin M, Mohammadi E, Eskandarzadeh N, Shamshirgaran MA. Prevalence of Chlamydia abortus infection in aborted sheep and goats in Kerman province,

southeast of Iran. Iranian Journal of Veterinary Science and Technology. 2023;15(3):42-7. Doi: 10.22067/ijvst.2023.82794.1263.

28. Hosein Abadi E, Saadati D, Najimi M, Hassanpour M. A Study on *Mycoplasma agalactiae* and *Chlamydia abortus* in aborted ovine fetuses in Sistan and Baluchestan region, Iran. Archives of Razi Institute. 2019;74(3):295-301. Doi: 10.22092/ari.2018.120393.1193.

29. Heidari S, Derakhshandeh A, Firouzi R, Ansari-Lari M, Masoudian M, Eraghi V. Molecular detection of *Chlamydia abortus*, *Coxiella burnetii*, and *Mycoplasma agalactiae* in small ruminants' aborted fetuses in southern Iran. Tropical Animal Health and Production. 2018;50:779-85. Doi: 10.1007/s11250-017-1494-2.

30. Hamed M, Esmaili H, Madani SA, Tajik P. The Frequency of Abortion Caused by *Chlamydia abortus* in Aborted Fetuses of Sheep and Goats in Iran. Journal of Medical Bacteriology. 2020;9(1-2):1-8.

31. Arif ED, Saeed NM, Rachid SK. Isolation and Identification of from Aborted Ewes in Sulaimani Province, Northern Iraq. Polish journal of microbiology. 2020;69(1):65-71. Doi: 10.33073/pjm-2020-009.


32. Nogarol C, Marchino M, Scala S, Belvedere M, Renna G, Vitale N, et al. Seroprevalence and Risk Factors Associated with *Chlamydia abortus* Infection in Sheep and Goats in North-Western Italy. Animals. 2024;14(2):291. Doi: 10.3390/ani14020291.

33. Tarusikirwa DF, Blacklaws B, Trotter CL. Seroprevalence and assessment of public awareness of *Brucella* spp., *Toxoplasma gondii* and *Chlamydia abortus* in small ruminants from selected smallholder commercial farms of Zimbabwe. Plos one. 2023;18(6):e0287902. Doi: 10.1371/journal.pone.0287902.


34. Castro-Flores R, Gaxiola SM, Diaz-Aparicio E, Rodriguez-Gaxiola MA, Montero A, Enriquez-Verdugo I, et al. Global, Regional, and National Prevalence of Chlamydiosis in Small

**Figure legends:**

**Figure 1.** Molecular findings of the present study for detection of the *Chlamydia* genome in aborted fetuses. The PCR products with an 83 bp band were detected through electrophoresis on agarose gels stained with a safe DNA stain. L: ladder 100 bp; S1, S3, S5: the samples with negative results; S2 and S4: positive samples with an 823 bp band.



**Figure 2.** A: liver: there are focal areas of cellular necrosis, surrounded by mononuclear cells (arrows). B: lung: The alveolar septa are thickened by mononuclear cells, which are associated with hyperemia of the pulmonary vessels. C: focal hemorrhage (arrow) in the cortex of the kidney. C: muscles: there are focal hemorrhages. D: spleen: there is lymphatic depletion with moderate hemorrhage E: brain: there are inflammatory cells in the meninges associated with notable hyperemia (arrowhead). F: brain: there is a focal hemorrhage (arrow). H&E.



## بررسی عفونت کلامیدیا در جنین های سقط شده با استفاده از مطالعات مولکولی و پاتولوژیک در استان آذربایجان شرقی، شمال غرب ایران

کلامیدوفیلا یکی از علل مهم عفونت های داخل رحمی در گوسفند و بز است که منجر به سقط جنین، مرده زایی و تولد فرزندانی ضعیف می شود. این بیماری در گوسفند به عنوان "سقط جنین انزوتیک گوسفند" شناخته شده است. قابل ذکر است که این بیماری به عنوان یک بیماری مشترک بین انسان و دام شناخته شده است که می تواند منجر به سقط جنین در انسان شود. بنابراین، مطالعه حاضر به منظور شناسایی عفونت کلامیدیا در جنین های سقط شده نشخوارکنندگان کوچک اهلی از جمله گوسفند و بز در استان آذربایجان شرقی، در شمال غرب ایران انجام شد. برای این منظور در مجموع از 62 جنین سقط شده از گله های گوسفند و بز نمونه برداری انجام شد. در کالبدگشایی، جنین معمولاً با وضعیت خوبی همراه با ضایعات کمی حفظ شده بود. نمونه های بافت برای مطالعات هیستوپاتولوژی و مولکولی جمع آوری شد. روش PCR معمولی با استفاده از پرایمرهای اختصاصی برای تشخیص ژنوم کلامیدیا انجام شد. علاوه بر این، نمونه های بافت تثبیت شده با فرمالین به طور معمول برای مطالعات هیستوپاتولوژیک آماده سازی شدند. ژنوم کلامیدیا در 33.87٪ (95٪ CI: 0.11 ± 0.32) از جنین های مورد بررسی تشخیص داده شد. معاینات هیستوپاتولوژیک هپاتیت چند کانونی، پنومونی و مننگوآنسفالیت غیر چرکی همراه با خونریزی کانونی در عضلات را نشان داد. در نتیجه، بررسی ژنوم کلامیدیا در جنین های سقط شده با شیوع بالا نشان می دهد که این عفونت می تواند نقش قابل توجهی در سقط جنین گوسفند و بز در آذربایجان شرقی داشته باشد. برای جلوگیری از سقط جنین های احتمالی در زنانی که در تماس نزدیک با نشخوارکنندگان سقط جنین هستند، اقدامات مدیریتی و کنترلی موثر برای سلامت عمومی در منطقه بسیار مهم است.

واژگان کلیدی: گوسفند، بز، سقط جنین، عفونت زئونوتیک، بهداشت عمومی.