



Effects of Adding α -tocopherol to Brahman Bull Chilled Semen on Sperm Quality, Lipid Peroxidation, Membrane Integrity, and DNA Integrity

Nur Ducha,^a Dyah Hariani,^a Widowati Budijastuti,^a Trinil Susilawati,^b Aulanni'am,^c
Sri Wahyuningsih^b

^a Biology Department, Faculty of Mathematics and Science, State University of Surabaya, Surabaya, East Java, Indonesia.

^b Reproduction Laboratory, Faculty of Animal Husbandry, Brawijaya University, Malang, East Java, Indonesia.

^c Biochemistry Laboratory, Faculty of Veterinary Medicine, Brawijaya University, Malang, East Java, Indonesia.

ABSTRACT

During storage at low temperatures, the spermatozoa quality changes due to cold shock and free radicals. Diluent supplementation with antioxidants is an effort to maintain the quality of spermatozoa during storage. This study aimed to evaluate the α -tocopherol effect in CEP extender on sperm quality, membrane integrity, and lipid peroxidation during storage at 4°C-5°C. This was a laboratory experiment that compared the use of 2 mM of α -tocopherol in CEP with no addition of α -tocopherol (as control) in five bulls. Semen was collected from Brahman bulls, diluted in CEP with and without α -tocopherol, and stored at 4°C-5°C. Sperm motility and viability were investigated by a light microscope at a $\times 400$ magnification using Eosin-Nigrosin staining. Moreover, membrane integrity was evaluated by lipid peroxidation using the MDA assay and hypoosmotic swelling test. The sperm motility, viability, and membrane integrity were higher in CEP with α -tocopherol. Lipid peroxidation was significantly different between the treatment and control groups. The α -tocopherol supplementation in the diluent CEP could maintain the spermatozoa quality during storage at 4°C-5°C.

Keywords

α -tocopherol, Brahman bull, Cauda epididymal plasma, Frozen, Spermatozoa

Number of Figures: 0
Number of Tables: 5
Number of References: 42
Number of Pages: 10

Abbreviations

AI: Artificial insemination

BSA: Bovine serum albumin

CEP: Cauda Epididymal Plasma

EDTA: Ethylenediaminetetraacetic acid

MDA: Malondialdehyde

PCR: Polymerase chain reaction

ROS: Reactive oxygen species

SNI: Indonesian National Standard

TBA: Thiobarbituric acid

Introduction

Some procedure in sperm preservation (storage) is performed at 4°C-5°C and freezing conditions, which affects sperm quality [1, 2]. Therefore, fresh semen must comply with a standard quality to be frozen, such as 70%-80% of motility and 80%-90% of viability [3]. The storage of semen at a temperature of 4°C-5°C can maintain the quality of spermatozoa for up to 2-3 days [4]. Moreover, this method does not depend on the availability of liquid nitrogen as a preservation agent. However, by modifying the diluent media, the storage time can be extended. Several studies showed an extension of storage time by using INRA96 diluent which could maintain the quality of bovine spermatozoa for up to 4 days [5], in commercial diluent Megapor S.L. Spain was able to maintain the quality of boar spermatozoa for up to 10 days [6].

Spermatozoa preservation at low temperatures cannot prevent the presence of free radicals. It reduces the acrosome integrity, motility, plasmalemma function, and viability, and induces chromatin damage [7, 8]. The freezing of spermatozoa preservation has been also reported to decrease spermatozoa motility and viability. The spermatozoa head region surrounding the plasma membrane is most susceptible to glutony. Spermatozoa heads have been reported to swell more than 50% [9]. Moreover, it triggers spermatozoa of RNA expression before and after freezing on its preservation [10]. Fresh semen commonly contains not only spermatozoa cells but also other cells, such as leukocytes and epithelial cells, before storage. These cells and immature spermatozoa are the sources of ROS [11].

Lipid peroxidation is caused by free radical and spermatozoa damage [12]. Lipids, proteins, and DNA oxidation are usually induced by high ROS levels in cells [2, 13]. There are high lipid membrane phospholipids in spermatozoa that lead to high ROS generation [14-16]. Spermatozoa with damaged and abnormal morphology produce higher ROS than spermatozoa with normal morphology [17]. Dead spermatozoa are the main source of ROS during storage at low temperatures [18, 19].

Antioxidants are required to reduce lipid peroxidation in the extender. Vitamin E (α -tocopherol) is one of the non-enzymatic antioxidants that protects polyunsaturated fatty acids, cell components, or cell membranes from oxidation by free radicals [20].

The addition of α -tocopherol in the semen diluent affects the spermatozoa quality at low-temperature conditions. It is able to maintain motility and protects spermatozoa from damage [21]. A study reported that the addition of α -tocopherol caused a significant change in the diluents of Madura bull spermato-

zoa [22]. Furthermore, α -tocopherol can maintain the integrity, motility, and viability of the spermatozoa membrane on the Simental bull after storage at 4°C-5°C. High-quality semen has a motility of $\geq 70\%$ and a viability of $\geq 75\%$. The use of antioxidants in diluents is useful for maintaining spermatozoa quality [23].

The CEP diluent was first developed by Verberckmoes et al. [24] who used it to store bovine semen at a temperature of 4°C-5°C with the constituent components mimicking the physical and chemical conditions in the cauda epididymal plasma of cattle [25]. They modified the concentration of egg yolk and replaced the antibiotic gentamicin with penicillin-streptomycin, and managed to maintain the quality of spermatozoa until day 8. The CEP diluent is also able to maintain the quality of sexed spermatozoa during storage at low temperatures [26]. This study aimed to evaluate the sperm quality, membrane integrity, and DNA integrity of Brahman bull that was stored in CEP with or without α -tocopherol at 4°C-5°C.

Results

As shown in Table 1, the mean percentage of sperm motility was not significantly different on days 0-3 between the control and treatment groups. However, a significant difference was found in spermatozoa motility in CEP with α -tocopherol compared to without α -tocopherol on days 4-7. Based on our study, sperm motility was better in CEP with α -tocopherol than without α -tocopherol. There were no significant differences ($p > 0.05$) in the viability percentage of sperms from day 0 to 4. However, on days 4-7, the percentage of sperm viability was higher ($p < 0.05$) in diluents with α -tocopherol than without α -tocopherol (Table 1).

Membrane Integrity

The motility and viability of spermatozoa were assessed along with spermatozoa function during storage based on their membrane integrity. The evaluations of membrane integrity percentage are illustrated in Table 2.

There was no significant difference in membrane integrity at the beginning of storage (days 0-4). On the other hand, there was a significant difference in membrane integrity in each treatment after day 5. The best results were in the CEP diluent using α -tocopherol on day 7.

Lipid Peroxidation

The MDA levels were measured to determine the existence of radicals inside and outside the cell. The MDA results can be seen in Table 3.

Table 1.

Effects of α -tocopherol addition to cauda epididymal plasma extender on sperm quality (Motility % and Viability %) during seven days (D) of storage at refrigerator temperature (4°C -5°C)

Treatments	% Motility in days							
	D0	D1	D2	D3	D4	D5	D6	D7
CEP without α -tocopherol	58.83 ^a ± 0.84	55.83 ^a ± 0.83	52.50 ^a ± 1.43	48.33 ^a ± 1.66	44.17 ^b ± 0.96	41.67 ^a ± 1.67	38.33 ^b ± 0.85	35.00 ^b ± 0.00
CEP + α -tocopherol	56.67 ^a ± 0.83	55.00 ^a ± 0.00	50.00 ^{ab} ± 0.00	50.00 ^a ± 0.00	49.17 ^a ± 0.83	46.67 ^a ± 1.66	45.83 ^a ± 0.83	45.00 ^a ± 0.00
Treatments	% Viability in days							
	D0	D1	D2	D3	D4	D5	D6	D7
CEP without α -tocopherol	80.76 ^a ± 1.06	70.94 ^{ab} ± 0.76	73.00 ^{ab} ± 0.59	69.54 ^a ± 0.68	62.61 ^{ab} ± 1.53	49.46 ^b ± 2.68	47.17 ^b ± 1.78	45.50 ^b ± 1.00
CEP + α -tocopherol	79.2 ^a ± 0.20	71.45 ^a ± 0.82	69.77 ^b ± 0.77	68.37 ^{ab} ± 0.43	64.13 ^a ± 0.31	60.58 ^a ± 1.31	58.44 ^a ± 0.47	56.10 ^a ± 0.11

CEP: Cauda epididymal plasma

^{a-b} uppercase letters in the same column indicate significant differences based on Student's T-test ($\alpha = 5\%$)

Table 2.

Effects of α -tocopherol addition to cauda epididymal plasma extender on membrane integrity % during seven days (D) of storage at refrigerator temperature (4°C-5°C)

Treatments	Membrane integrity (%)							
	D0	D1	D2	D3	D4	D5	D6	D7
CEP without α -tocopherol (control)	79.59 ^a ± 1.06	72.61 ^a ± 0.72	71.76 ^a ± 0.65	68.93 ^{ab} ± 0.35	60.86 ^{ab} ± 1.60	48.55 ^b ± 2.57	46.41 ^b ± 1.99	43.98 ^b ± 1.33
CEP + α -tocopherol	78.23 ^a ± 0.32	70.53 ^a ± 0.99	68.84 ^{ab} ± 0.95	67.69 ^a ± 0.35	65.92 ^a ± 0.63	65.66 ^a ± 1.94	60.91 ^a ± 1.02	56.29 ^a ± 0.33

CEP: Cauda epididymal plasma

^{a-b} uppercase letters in the same column indicate significant differences based on Student's T-test ($\alpha = 5\%$)

Table 3.

Effects of α -tocopherol addition to cauda epididymal plasma extender on malondialdehyde values during four days (D) of storage at refrigerator temperature (4°C-5°C)

Treatments	MDA value			
	D1	D3	D5	D7
CEP without α -tocopherol (control)	0.222 ± 0.191 ^a	0.306 ± 0.179 ^a	0.390 ± 0.055 ^a	0.613 ± 0.234 ^a
CEP + α -tocopherol	0.116 ± 0.007 ^b	0.168 ± 0.168 ^b	0.175 ± 0.018 ^b	0.182 ± 0.008 ^b

CEP: Cauda epididymal plasma; MDA: malondialdehyde

^{a-b} uppercase letters in the same column indicate significant differences based on Student's T-test ($\alpha=5\%$)

There were significant differences in MDA levels between treatments. A significant difference was found on days 1-7 of storage. The lowest MDA levels were shown in the α -tocopherol treatment, while the highest MDA levels were observed in the control group (CEP without α -tocopherol).

DNA Integrity

DNA integrity during storage may change due to the presence of free radicals. DNA integrity was assessed by sequencing a gene that controls spermatozoa motility (NAD1-1) (Table 4).

Table 4.
Data of NAD-1 sequences

	10	20	30	40	50	60	70	80
Control Day-1	GTTCTAGGCT	ACATACAAC	CCGAAAAGGC	CCAAATGTTCG	TAGGCCCAT	TGGCCTACTT	CAACCTATCG	CCGATGCAAT
Control Day-3	GTTCTAGGCT	ACATACAAC	CCGAAAAGGC	CCAAATGTTCG	TAGGCCCAT	TGGCCTACTT	CAACCTATCG	CCGATGCAAT
Control Day-5	GTTCTAGGCT	ACATACAAC	CCGAAAAGGC	CCAAATGTTCG	TAGGCCCAT	TGGCCTACTT	CAACCTATCG	CCGATGCAAT
Control Day-7	GTTCTAGGCT	ACATACAAC	CCGAAAAGGC	CCAAATGTTCG	TAGGCCCAT	TGGCCTACTT	CAACCTATCG	CCGATGCAAT
α-tocopherol Day-1	GTTCTAGGCT	ACATACAAC	CCGAAAAGGC	CCAAATGTTCG	TAGGCCCAT	TGGCCTACTT	CAACCTATCG	CCGATGCAAT
α-tocopherol Day-3	GTTCTAGGCT	ACATACAAC	CCGAAAAGGC	CCAAATGTTCG	TAGGCCCAT	TGGCCTACTT	CAACCTATCG	CCGATGCAAT
α-tocopherol Day-5	GTTCTAGGCT	ACATACAAC	CCGAAAAGGC	CCAAATGTTCG	TAGGCCCAT	TGGCCTACTT	CAACCTATCG	CCGATGCAAT
α-tocopherol Day-7	GTTCTAGGCT	ACATACAAC	CCGAAAAGGC	CCAAATGTTCG	TAGGCCCAT	TGGCCTACTT	CAACCTATCG	CCGATGCAAT
	90	100	110	120	130	140	150	160
Control Day-1	CAAACTTTTC	ATTAAGAAG	CAC TACGACC	CGCCACATCT	TCAACCTCAA	TATTCATCCT	AGCACCCATT	TTAGCCCTAG
Control Day-3	CAAACTTTTC	ATTAAGAAG	CAC TACGACC	CGCCACATCT	TCAACCTCAA	TATTCATCCT	AGCACCCATT	TTAGCCCTAG
Control Day-5	CAAACTTTTC	ATTAAGAAG	CAC TACGACC	CGCCACATCT	TCAACCTCAA	TATTCATCCT	AGCACCCATT	TTAGCCCTAG
Control Day-7	CAAACTTTTC	ATTAAGAAG	CAC TACGACC	CGCCACATCT	TCAACCTCAA	TATTCATCCT	AGCACCCATT	TTAGCCCTAG
α-tocopherol Day-1	CAAACTTTTC	ATTAAGAAG	CAC TACGACC	CGCCACATCT	TCAACCTCAA	TATTCATCCT	AGCACCCATT	TTAGCCCTAG
α-tocopherol Day-3	CAAACTTTTC	ATTAAGAAG	CAC TACGACC	CGCCACATCT	TCAACCTCAA	TATTCATCCT	AGCACCCATT	TTAGCCCTAG
α-tocopherol Day-5	CAAACTTTTC	ATTAAGAAG	CAC TACGACC	CGCCACATCT	TCAACCTCAA	TATTCATCCT	AGCACCCATT	TTAGCCCTAG
α-tocopherol Day-7	CAAACTTTTC	ATTAAGAAG	CAC TACGACC	CGCCACATCT	TCAACCTCAA	TATTCATCCT	AGCACCCATT	TTAGCCCTAG
	170	180	190	200	210	220	230	240
Control Day-1	GCTTAGCCTT	AACCATGTGA	ATCCCCCTCC	CAATACCCTA	CCCTCTTATT	AACATAAAATC	TAGGAATCCT	ATTTATACTA
Control Day-3	GCTTAGCCTT	AACCATGTGA	ATCCCCCTCC	CAATACCCTA	CCCTCTTATT	AACATAAAATC	TAGGAATCCT	ATTTATACTA
Control Day-5	GCTTAGCCTT	AACCATGTGA	ATCCCCCTCC	CAATACCCTA	CCCTCTTATT	AACATAAAATC	TAGGAATCCT	ATTTATACTA
Control Day-7	GCTTAGCCTT	AACCATGTGA	ATCCCCCTCC	CAATACCCTA	CCCTCTTATT	AACATAAAATC	TAGGAATCCT	ATTTATACTA
α-tocopherol Day-1	GCTTAGCCTT	AACCATGTGA	ATCCCCCTCC	CAATACCCTA	CCCTCTTATT	AACATAAAATC	TAGGAATCCT	ATTTATACTA
α-tocopherol Day-3	GCTTAGCCTT	AACCATGTGA	ATCCCCCTCC	CAATACCCTA	CCCTCTTATT	AACATAAAATC	TAGGAATCCT	ATTTATACTA
α-tocopherol Day-5	GCTTAGCCTT	AACCATGTGA	ATCCCCCTCC	CAATACCCTA	CCCTCTTATT	AACATAAAATC	TAGGAATCCT	ATTTATACTA
α-tocopherol Day-7	GCTTAGCCTT	AACCATGTGA	ATCCCCCTCC	CAATACCCTA	CCCTCTTATT	AACATAAAATC	TAGGAATCCT	ATTTATACTA
	250	260	270	280	290	300	310	320
Control Day-1	GCCATATCAA	GCCTAGCCGT	ATACTCTATC	CTCTGATCAG	GTTGAGCCTC	CAATTCAAAA	TACGCACTAA	TCCGAGCCCT
Control Day-3	GCCATATCAA	GCCTAGCCGT	ATACTCTATC	CTCTGATCAG	GTTGAGCCTC	CAATTCAAAA	TACGCACTAA	TCCGAGCCCT
Control Day-5	GCCATATCAA	GCCTAGCCGT	ATACTCTATC	CTCTGATCAG	GTTGAGCCTC	CAATTCAAAA	TACGCACTAA	TCCGAGCCCT
Control Day-7	GCCATATCAA	GCCTAGCCGT	ATACTCTATC	CTCTGATCAG	GTTGAGCCTC	CAATTCAAAA	TACGCACTAA	TCCGAGCCCT
α-tocopherol Day-1	GCCATATCAA	GCCTAGCCGT	ATACTCTATC	CTCTGATCAG	GTTGAGCCTC	CAATTCAAAA	TACGCACTAA	TCCGAGCCCT
α-tocopherol Day-3	GCCATATCAA	GCCTAGCCGT	ATACTCTATC	CTCTGATCAG	GTTGAGCCTC	CAATTCAAAA	TACGCACTAA	TCCGAGCCCT
α-tocopherol Day-5	GCCATATCAA	GCCTAGCCGT	ATACTCTATC	CTCTGATCAG	GTTGAGCCTC	CAATTCAAAA	TACGCACTAA	TCCGAGCCCT
α-tocopherol Day-7	GCCATATCAA	GCCTAGCCGT	ATACTCTATC	CTCTGATCAG	GTTGAGCCTC	CAATTCAAAA	TACGCACTAA	TCCGAGCCCT
	330	340	350	360	370	380	390	400
Control Day-1	ACGAGCAGTA	GCACAAACAA	TCTCATACGA	AGTAACACTA	GCAATTATCC	TATTATCAGT	GTCCTAATA	AGTGGGTCCT
Control Day-3	ACGAGCAGTA	GCACAAACAA	TCTCATACGA	AGTAACACTA	GCAATTATCC	TATTATCAGT	GTCCTAATA	AGTGGGTCCT
Control Day-5	ACGAGCAGTA	GCACAAACAA	TCTCATACGA	AGTAACACTA	GCAATTATCC	TATTATCAGT	GTCCTAATA	AGTGGGTCCT
Control Day-7	ACGAGCAGTA	GCACAAACAA	TCTCATACGA	AGTAACACTA	GCAATTATCC	TATTATCAGT	GTCCTAATA	AGTGGGTCCT
α-tocopherol Day-1	ACGAGCAGTA	GCACAAACAA	TCTCATACGA	AGTAACACTA	GCAATTATCC	TATTATCAGT	GTCCTAATA	AGTGGGTCCT
α-tocopherol Day-3	ACGAGCAGTA	GCACAAACAA	TCTCATACGA	AGTAACACTA	GCAATTATCC	TATTATCAGT	GTCCTAATA	AGTGGGTCCT
α-tocopherol Day-5	ACGAGCAGTA	GCACAAACAA	TCTCATACGA	AGTAACACTA	GCAATTATCC	TATTATCAGT	GTCCTAATA	AGTGGGTCCT
α-tocopherol Day-7	ACGAGCAGTA	GCACAAACAA	TCTCATACGA	AGTAACACTA	GCAATTATCC	TATTATCAGT	GTCCTAATA	AGTGGGTCCT

Table 4 - Cont'd

Control Day-1	TTACCCCTCT
Control Day-3	TTACCCCTCT
Control Day-5	TTACCCCTCT
Control Day-7	TTACCCCTCT
α -tocopherol Day-1	TTACCCCTCT
α -tocopherol Day-3	TTACCCCTCT
α -tocopherol Day-5	TTACCCCTCT
α -tocopherol Day-7	TTACCCCTCT

Note: Alignment results for sample

Discussion

At the beginning of storage, no significant differences were found between the control and treatment groups. It demonstrated that short storage time did not affect the spermatozoa quality. This result also indicated that the dilution process of semen in CEP with and without α -tocopherol was a success. Therefore, it did not cause a significant change in spermatozoa quality. It was also supported by the spermatozoa quality such as motility and viability become slowly.

In the present study, we revealed that the CEP media was very suitable for the storage of bull spermatozoa, although the semen was categorized as rejected semen. The CEP media could maintain the spermatozoa quality from the rejected semen during the storage process at 4°C-5°C. These findings are also in line with Verbeckmoes [24] and Duchá [25, 27, 28]. The CEP media provides chemical and physical conditions to store bulls' spermatozoa. The basic diluent of CEP was developed by imitating the physical and chemical conditions of bovine epididymal plasma which has an osmolarity in the range of 290-300 mosm, contains ionic components Na, K, Ca, Cl, Mg, P, bicarbonate, energy source molecules (i.e., fructose and sorbitol), molecules for pH balance (i.e., tris and citric acid), as well as antibiotics and egg yolk [24, 27]. Another research [24] showed that CEP was able to maintain the quality of bovine spermatozoa for up to 6 days at a temperature of 4°C-5°C with the addition of 20% egg yolk. Furthermore, the replacement of gentamicin with penicillin-streptomycin showed that CEP could maintain the quality of bovine spermatozoa for up to 8 days [25].

The addition of α -tocopherol to the CEP diluent strengthens the ability to maintain the quality of Brahman bull spermatozoa during storage at 4°C-5°C. It was similar to the SNI for AI until the seventh day of storing Brahman bull semen. The CEP media without α -tocopherol keeps the quality of bull spermatozoa from the rejected semen until the fifth day of storage. The increase in free radicals during the storage of spermatozoa at low temperatures may be due to

the production of free radicals from dead or abnormal spermatozoa. The addition of antioxidants into the diluent media tends to minimize the production of free radicals [19].

Sperm stored at a low temperature can be damaged by free radicals. α -tocopherol is known as an antioxidant that reduces free radicals and can improve the quality of spermatozoa. We found that the addition of α -tocopherol into the CEP extender did not protect the spermatozoa of the Brahman bull at the beginning of storage. We observed no significant difference between the control and treatment groups. Furthermore, free radicals in the diluent can be overcome by antioxidants in spermatozoa cells during five days of storage. Therefore, the extent of damage to cells was low. α -tocopherol plays the role of an antioxidant and protects the cells or other compounds against free radicals by donating one hydrogen atom from the OH group to free radical compounds, resulting in tocopheroxyl radical compounds that are more stable and non-damaging. It can stop the propagation process when lipid peroxidation occurs in the membrane of spermatozoa [29]. The concentration of α -tocopherol can vary considering the percentage of lipids in the environment, affecting the solubility of vitamin E and the morphology of spermatozoa. Therefore, the type and composition of the diluent media and the type of animal will influence the dose of α -tocopherol [30].

There were significant differences in membrane integrity, motility, and viability on the sixth day of storage. These conditions were caused by free radicals in the diluent. Antioxidants were unable to reduce free radicals in the cell, especially in CEP diluents without adding α -tocopherol. The decrease in spermatozoa quality, such as motility, viability, and membrane integrity was faster in diluents without α -tocopherol compared to diluents containing α -tocopherol. Brahman bull spermatozoa stored in CEP diluents without α -tocopherol showed a low quality of sperm on the fifth day of storage. It was due to the presence of free radicals in semen. This finding is similar to buffalo sperm which has a higher quality with an extender containing α -tocopherol or vitamin C [31]. These results indicate that α -tocopherol provides the best protection for the plasma membrane of spermatozoa. α -tocopherol is known as an antioxidant that prevents lipid peroxidation during storage in bovine spermatozoa [32].

The integrity of spermatozoa DNA during storage was assessed based on the profile of the genes encoding spermatozoa motility (NAD1-1). Spermatozoa motility was a major parameter in determining spermatozoa quality at the center of AI. The motility parameter is determined by the ability of spermatozoa to fertil-

ize eggs. These results revealed no changes in genes during various treatments until day seven of storage. It indicated that the presence of free radicals during storage for up to seven days did not damage the cell and DNA structure. Gene sequences did not show any changes in either the control or treatment groups, before or after storage. The process of storing the liquid at 4°C-5°C was simpler than the freezing method and fewer free radicals existed in the liquid method than in the frozen technique. The freezing process can trigger damage to the integrity of the spermatozoa membrane, compared to storage at 4°C-5°C as happened in the freezing process of human spermatozoa. Freezing storage triggers damage in chromatin integrity by oxidative stress [33]. The storage of human spermatozoa at freezing temperatures damages DNA integrity [34]. DNA integrity did not damage after storage at 4°C-5°C, while DNA integrity was damaged after freezing [35].

Assessment of both sperm motility and sperm viability by light microscope was an important limitation of the present study. However, computer-aided sperm analysis technology (CASA) is more accurate for such assessments. Moreover, other concentrations of α -tocopherol or combinations of other antioxidants with α -tocopherol may be required to obtain better results on bull spermatozoa stored at 4°C-5°C. Therefore, further research is needed in this important field of research concerning spermatozoa preservations. It is necessary to study the effect of adding α -tocopherol to the CEP diluent in which each diluent has compositional characteristics, especially lipids that can affect α -tocopherol solubility. Consequently, it can influence the effectiveness of α -tocopherol in counteracting free radicals.

Conclusion

The sperm motility, viability, and membrane integrity were higher in CEP with the supplementation of α -tocopherol. Our present study showed that lipid peroxidation was significantly different between the treatment and control groups. The CEP with α -tocopherol supplementation could maintain the spermatozoa quality during storage at 4°C-5°C.

Materials & Methods

Preparation of Extender with α -tocopherol Addition

The CEP extender in the present study was prepared according to Verbeckmoes et al. [18, 27] with antibiotic, egg concentration, and a different method (water jacket method) [20]. The following compounds were included in the CEP extender: BSA 2.0 gr/l, CaCl₂(H₂O)₂ 3.0 mmol/l, citrate acid mmol/l, fructose 55 mmol/l, KH₂PO₄ 20.0 mmol/l, KCl 7.0 mmol/l, NaH₂PO₄ 8.0

mmol/l, NaHCO₃ 11.9 mmol/l, NaCl 15 mmol/l, MgCl₂(H₂O)₆ 3.0 mmol/l, penicillin 1000 IU, streptomycin 1 gr, sorbitol 1.0 gr/l, and tris 133.7 mmol/l. Next, 2 mM of α -tocopherol (Sigma, USA) was added to the CEP extender while the control group was not supplemented with α -tocopherol.

Bull Semen Collection and Preparation

Fresh semen of the bull was obtained from AI Center in Singosari, Malang, Indonesia by an artificial vagina. Bulls' semen was obtained twice a week to gain optimum semen quality. The fresh semen was observed to evaluate motility and viability before being diluted. The low quality of semen, according to the SNI, was continued for the process of dilution and freezing. If the quality was below the SNI, semen was rejected. According to the SNI, low-quality semen does not meet the requirements of the freezing process. For example, the motility of spermatozoa is less than 70% and the viability percentage is less than 75%, and it is called rejected semen. Fresh semen was diluted 25 times in a CEP extender with 2 mM α -tocopherol and without antioxidants as the control group. Spermatozoa were stored at refrigerator temperature (4°C-5°C) in darkness conditions at 25×10⁶/ml concentration.

In the present study, we used fresh semen with low quality. We evaluated the level of spermatozoa degradation from fresh low-quality semen (rejected for the freezing process) during storage at low temperatures on CEP diluents with α -tocopherol as an antioxidant, compared to those without α -tocopherol.

Sperm Motility

Spermatozoa motility was measured by observing semen using a light microscope to determine the percentage of progressive motility. Spermatozoa were taken by a stick glass on days 0 and 8 in the CEP-2 extender with and without egg yolk. Next, it was placed on an object glass and covered with cover glass. The observation was conducted under a light microscope with ×200 magnification. Evaluation of the motility-based method of Garner and Hafez [36] was performed by two people to determine sperm motility.

Sperm Viability

Sperm viability was assessed using Eosin-Nigrosin staining to gain the permanent slides. Nigrosin provides a dark background to recognize viable cells. Non-viable sperms had red or dark pink heads and viable sperms had white or faintly-pink heads.

Membrane Integrity

Membrane integrity was determined using the hypoosmotic swelling test. It was performed by incubating 100 μ l semen in the control and treatment groups with 1 ml of 125 mOsm/l hypoosmotic (0.31 g sodium citrate and 0.565 g fructose in 50 ml H₂O) at 37°C for 30 min. Afterwards, 0.2 ml of the mixture was spread using a coverslip on a warm slide after incubation. The observation was performed using 200 sperms under light microscopy at a magnification of ×400. Sperms with swollen or coiled tails were recorded [37].

Lipid Peroxidation

Sperm oxidative levels were determined using the MDA assay-based TBA reaction. A total of 1 ml in each spermatozoa treatment (five technical replications) was incubated in 0.7 ml of tris buffer (100 mM, pH 7.4), 10 mM of iron sulfate, and 100 mM sodium ascorbate at 37°C for 60 min. This reaction was stopped by adding 1 ml of 10% trichloroacetic acid on ice for 15 min. Samples were immediately centrifuged at 7800 rpm for 15 min and the supernatant was removed. A total of 2 ml supernatant was added.

ed with 1 ml of 1% TBA into endogenous peroxidation. Next, the specimen was boiled for 10 min and kept in a cool condition. The absorbance levels were read by spectrometry at 530 nm [38, 39].

DNA Integrity

DNA extraction

DNA extraction was conducted according to the method of Silva et al. [40]. Each semen treatment was divided into three aliquots. DNA was extracted with phenol-chloroform according to Hanson and Ballantyne [41]. A total of 100 µl of semen aliquots were centrifuged at 6000 rpm for 5 min. Pellet was resuspended using 1 ml Tris EDTA solution (1 mM EDTA, 100 mM NaCl, 100 mM Tris-HCl, pH 8.0) and centrifuged at 6000 rpm for 5 min.

A volume of 500 µl of lysis buffer containing 25 mM EDTA, 0.5% sodium dodecyl sulfate, 100 mM NaCl, and 10 mM Tris-HCl with a pH of 8.0 was added to the pellet, followed by 22 µl of 0.1 M dithiothreitol and 25 µl proteinase K (QIAGEN GmbH). Then, the sample was incubated at 55°C for 3 h, with a vortex in each hour. A total of 500 µl phenol was added, balanced with Tris at pH 7.8, and followed by a vortex. Centrifugation was conducted at 10000 rpm for 3 min. The supernatant was transferred to another tube, along with 300 µl phenol and 300 µl chloroform, followed by vortex and centrifugation at 10000 rpm for 3 min. The supernatant was placed into a new tube and 700 µl chloroform was added. The mixture was centrifuged and gently vortexed again. The supernatant was placed into a new tube containing 95% ethanol. Samples were incubated at -20°C for 4 h. Each sample was centrifuged at 10000 rpm for 10 min, and the supernatant was removed. Then, the DNA pellet was dried and resuspended with 50 µl 1X TE buffer (100 mM Tris-HCl, pH 7.5, 0.25 M EDTA), and stored at -20°C until further analysis.

DNA concentration and purity

Genomic DNA concentration and purity were assessed by measuring optical density in a Genesys Spectrophotometer. DNA absorbance was read at 260 nm to determine DNA concentration. DNA purity was evaluated by determining the A260/280 ratio and comparing it with a reference value of 1.8 [42].

DNA visualization on agarose gel

The existence and quality of DNA genomes were analyzed with a 0.8% agarose gel. Five microliters of sperm pellets were stained with 0.3 µl of Blue-Green (LGC Biotecnologia, Cotia, SP, Brazil), and electrophoresis was performed on the agarose gel. The results were visualized under an ultraviolet transilluminator (Vilber Lourmat, Paris, France), and the image was digitalized (C7070; Olympus, Tokyo, Japan).

DNA amplification

Genomic DNA amplification with PCR used ND1 gene primers. The ND1 gene is 925 bp and the PE Biosystems ABI 3700 DNA Analyzer 96 capillary electrophoresis system could not accommodate the 925 bp sequence.

Table 5 presents the primers for segments 1 and 2 of the ND1 sequence.

The first segment of the ND1 sequence used the antisense primer (3'-5'), and the second segment used the sense primer (5'-3') to align the two segments. Primers were selected specifically for mtDNA to distinguish between mitochondrial and nuclear DNA.

Data Analysis

Data including average motility, viability, and membrane integrity were analyzed by analysis of variance to determine whether the treatment caused an effect. Moreover, the Student's T-test ($\alpha = 5\%$) was utilized to assess the differences between treatments.

Fundings

This investigation was supported by a research grant from the Indonesian government.

Authors' Contributions

ND: Conceptualization, Supervision, Formal analysis, Writing original draft; DH : Writing original draft, Review and Editing. Data Curation; WB : Methodolgy, Formal Analysis, Review and Editing; TS : Data Curation, Formal Analysis, Writing Original Draft; AA : Formal Analysis, Writing-Review and Editing, Methodology; SW : Methodology, Data Curation, Writing-Review and Editing.

Acknowledgements

The authors give thanks to Brawijaya University, State Surabaya University, and Artificial Insemination (AI) Center in Singosari, for the contributions of this research.

Conflict of interest

The authors declared no conflict of interest for this research.

References

- Awad MM. Effect of some permeating cryoprotectants on CASA motility results in cryopreserved bull spermatozoa. *Anim Reprod Sci.* 2011;123(3-4):157-162. Doi: 10.1016/j.anireprosci.2011.01.003.
- Kardirvel G, Kumar S and Kumaresan A. Lipid peroxidation, mitochondrial membrane potential and DNA integrity of spermatozoa relation to intracellular reactive oxygen

Table 5.
Primers in the mtDNA sequencing procedure

Segment	Sense/antisense primer	Primer sequence
1	457 bp (3131-3588)	BOVNADH1S (sense)
		BOVNADH1A (antisense)
2	488 bp (3567-4055)	BOVNADH1S (sense)
		BOVNADH1A (antisense)

- species in liquid and frozen-thawed buffalo semen. *Anim Reprod Sci.* 2009;114(1-3):125-134. Doi: 10.1016/j.anireprosci.2008.10.002.
3. Ax RL, Dally M, Didion BA, Lenz RW, Love CC and Varner DD et al. Semen Evaluation: Hafez/Reproduction in Farm Animals. In Hafez B, Hafez ESE. *Reproduction in Farm Animals*. 7th eds. Baltimore, Maryland, USA: Lippincott Williams & Wilkins; 82 – 96; 2008. Doi:10.1002/9781119265306.ch25.
 4. Vishwanath R, Shannon P. Storage of bovine semen in liquid and frozen state. *Anim. Reprod. Sci.*, 2000;62 (2000): 23-53. Doi: 10.1016/s0378-4320(00)00153-6.
 5. Murphy EM, Meara CO, Eivers B, Lonergan P and Fair S. Optimizing storage temperature of liquid bovine semen diluted in INRA96. *J. Dairy Sci.* 2018;101:5549–5558. Doi: 10.3168/jds.2017-14205.
 6. Gaczarzewicz D, Udala J, Piasecka M, Blaszczyk B, Stankiewicz T. *Turkish Journal of Biology.* 2015;39: 582-594. Doi:10.3906/biy-1412-76.
 7. Ansari MS, Rakha BA, Malik MF, Andrabi SMH, Ullah N and Iqbal R et al. Effect of cysteine addit ion to the freezing extender on the progressive motility, viability, plasma membrane, and DNA integrity of nili-Ravi buffalo (*Bubalus bubalis*) bull spermatozoa. *J Appl Anim Res.* 2016;44(1):36–41. Doi: 10.1080/09712119.2014.987292.
 8. Guthrie HD, Welch GR, Theisen DD and Woods LC. Effects of hypothermic storage on intracellular calcium, reactive oxygen species formation, mitochondrial function, motility, and plasma membrane integrity in striped bass (*Morone saxatilis*) sperm. *Theriogenology.* 2011;75(5):951–961. Doi:10.1016/j.theriogenology.2010.10.037.
 9. Khalil WA, El-Harairy MA, Zaidan AEB and Hassan MAE. Elsaeed OM. Evaluation of bull spermatozoa during and after cryopreservation: Structural and ultrastructural insights. *International Journal of Veterinary Science and Medicine.* 2017;6(1): S49-S56. Doi: 10.1016/j.ijvsm.2017.11.001.
 10. Dai DH, Qazi IH, Ran MX, Liang K, Zang Y and Zhang M et al. Exploration of miRNA and mRNA Profiles in Fresh and Frozen-Thawed Boar Sperm by Transcriptome and Small RNA Sequencing. *Int J Mol Sci.* 2019;20(802):1-19. Doi:10.3390/ijms20040802.
 11. Cocuzza M, Sikka SC, Athayde KS and Agarwal A. Chemical relevance of oxidative stress and sperm chromatin damage in male infertility an evidence based analysis. *Int Braz J Urol.* 2007;33(5):603–621. Doi: 10.1590/s1677-55382007000500002.
 12. Sanocka D, and Kurpisz M. Reactive Oxygen Species and Sperm Cells. *Reprod Biol Endocrinol.* 2004;2:12. Doi:10.1186/1477-7827-2-12.
 13. Hezavehei M, Sharafi M, Kouchesfahani HM, Henkel R, Agarwal A, and Esmaeili V et al. Sperm cryopreservation. A review on curent molecular cryobiology and advanced approaches. *Reprod Biomed Online.* 2018;37(3):327-339. Doi:10.1016/j.rbmo.2018.05.012.
 14. Marnett LJ. Oxy radicals, lipid peroxidation and and DNA damage. *Toxicology.* 2002;181–182:219–222. Doi: 10.1016/s0300-483x(02)00448-1.
 15. Kim S, Lee YJ and Kim YJ. Changes in sperm membrane and ROS following cryopreservation of liquid boar semen stored at 15°C. *Anim Reprod Sci.* 2011;124(1–2):118–124. Doi: 10.1016/j.anireprosci.2011.01.014.
 16. Prapaiwan N, Tharasanit T, Punjachairpornpol S, Yamtang D, Roongsithichai A and Moonarmart W et al. Low - Density lipoprotein improves cryopreserved epididymal spermatozoa. *Asian-Australas J Anim Sci.* 2016;29(5):646-651. Doi: 10.5713/ajas.15.0572.
 17. Ball BA, Vo AT, Baumber J. Generation of reactive oxygen species by equine spermatozoa. *AJVR,* 2001;62(4). Doi:10.2460/ajvr.2001.62.508.
 18. Alomar M, Alzoabi M, Zarkawi M. Kinetics of hydrogen peroxide generated from live and dead ram spermatozoa and the effects of catalase and oxidase substrates addition. *Czech J. Anim. Sci.,* 2016; 61(1): 1–7. Doi:10.17221/8662-CJAS.
 19. Alomar M. Formation of Hydrogen Peroxide by Chilled Goat Spermatozoa and the Effects of Dead Spermatozoa on Motility Characteristics. *Iranian Journal of Applied Animal Science.* 2019;1(9): 79-85. Available at: https://ijas.rasht.iau.ir/article_663557.html.
 20. Motemani M, Chamani M, Sharafi M and Masoudi R. Alpha-tocopherol improves frozen-thawed sperm quality by reducing hydrogen peroxide during cryopreservation of bull semen. *Span J Agric Res.* 2017;15(1):1-7. Doi:10.5424/sjar/2017151-9761.
 21. Kalthur G, Raj S, Thiyagarajan A, Kumar SK, Kumar P and Adiga SK. Vitamin E supplementation in semen-freezing medium improves the motility and protects sperm from freeze-thaw-induced DNA damage. *Fertil Steril.* 2011;95(3):1149-1151. Doi: 10.1016/j.fertnstert.2010.10.005.
 22. Ratnani H, Ihsan MN, Ciptadi G and Suyadi S. Effect of α -tocopherol supplementation in the Extender on the Sperm Quality of Maduran Bull Before and After Quick Freezing. *Int J Adv Res.* 2017;5(7):1378-1389. Doi: 10.21474/IJAR01/4849.
 23. Azura S, Ratnani H, Soepranianondo K, Susilowati S, Hariadi M and Samik A. Effect of α - tocopherol supplementation in diluents on motility, viability, and plasma membrane integrity of Simmental bull spermatozoa after cooling. *Ovozoa [serial on the Internet].* [cited 2020 February 4] 2020;9(1):1-6. Available from: <https://e-journal.unair.ac.id/OVZ/index>.

24. Verberckmoes S, Van Soom A, Dewulf J, De Pauw I and de Kruif A. Storage of fresh bovine semen in diluent based on the ionic composition of cauda epididymal plasma. *Reprod Domest Anim.* 2004;39(6):1-7. Doi: 10.1111/j.1439-0531.2004.00521.x.
25. Ducha N, Susilawati T, Aulanni'am and Wahyuningsih S. Motilitas dan Viabilitas Spermatozoa Sapi Limousin Selama Penyimpanan pada Refrigerator dalam Pengencer CEP-2 dengan Suplementasi Kuning Telur. *Indonesian Journal of Veterinary Sciences.* 2013;7(1):2002-2005. Doi: 10.21157/j.ked.hewan.v7i1.555.
26. Firdaus F and Ratnawati D. Effectiveness of cauda epididymal plasma-2 and lecithin based diluents to minimize abnormality of sexing albumin spermatozoa during cold storage. *Veterinary World, EISSN: 2231-0916; 2021.* Doi:10.14202/vetworld.2021.2543-2548.
27. Ducha N, Hariani D and Budijastuti W. The relationship of physical and chemical quality before and after storage at temperature 4-5°C. *J Phys Conf Ser.* 2018;953(1):012208. Doi: 10.1088/1742-6596/953/1/012208.
28. Ducha N, Susilawati T, Aulanni'am, Wahyuningsih S and Pangestu M. Ultrastructure and fertilizing ability of Limousin bull sperm after storage in CEP-2 extender with and without egg yolk. *Pak J Biol Sci.* 2012; 15(20). Doi: 10.3923/pjbs.2012.979.985.
29. Bansal AK, dan Bilaspuri G, 2009. Antioxidant Effect of Vitamin E on Motility, Viability and Lipid Peroxidation of Cattle Spermatozoa Under Oxidative Stress. *Jurnal Anim Sci Papand Rep.* 27(1):11-13. Available at: https://www.researchgate.net/publication/285767630_Antioxidant_effect_of_vitamin_E_on_motility_viability_and_lipid_peroxidation_of_cattle_spermatozoa_under_oxidative_stress#fullTextFileContent.
30. Azzi A. Molecular mechanism of alpha-tocopherol action. *Free. Radic. Biol. Med.* 43:16-21. Doi:10.1016/j.freeradbiomed.2007.03.013; 2007.
31. Andrabi SMH, Ansari MS, Ullah N and Afzal M. Effect of non-enzymatic antioxidants in extender on post-thaw quality of buffalo (*Bubalus bubalis*) bull spermatozoa. *Pakistan Vet J.* 2008;28:159-162. Available at: https://www.researchgate.net/publication/26579329_Effect_of_non-enzymatic_antioxidants_in_extender_on_post-thaw_quality_of_buffalo_Bubalus_bubalis_bull_spermatozoa#fullTextFileContent.
32. Golden C, Rosenkrans CF, and Johnson Z. Effects of ascorbic acid and alpha-tocopherol on cryopreserved boar sperm. *Ark Anim Sci Dep Rep.* 2002;499:120-123. Available at: https://www.researchgate.net/publication/303173077_Effects_of_ascorbic_acid_and_alpha-tocopherol_on_cryopreserved_boar_sperm.
33. Fraser L, Strzezek J and Kordan W. Effect of Freezing on Sperm Nuclear DNA. *Reprod Domest Anim.* 2011;46(Suppl 2):14-7. DOI: 10.1111/j.1439-0531.2011.01815.x.
- [34]. Lusignan MF, Li X, Herrero B, Delbes G and Chan PTK. Effects of different cryopreservation methods on DNA integrity and sperm chromatin quality in men. *Andrology.* 2018;6(6): 829-835. Doi: 10.1111/andr.12529.
35. Nandre R, Derashi H and Joshi C. Evaluation of Buffalo Bull Spermatozoa DNA Damage Using Single Cell Gel Electrophoresis. *Int J Life Sci Pharma Res.* 2011;(1):38-43. Available at: https://www.researchgate.net/publication/216021238_Evaluation_of_buffalo_bull_spermatozoal_DNA_damage_using_single_cell_gel_electrophoresis#fullTextFileContent.
36. Garner DL, and Hafez ESE. Spermatozoa and Seminal Plasma: Hafez/Reproduction in Farm Animals. In: ESE Hafez, Hafez B. *Reproduction in farm Animal.* 7th eds. Baltimore, Maryland, USA: Lippincott & Williams. Baltimore. pp. 96 – 109; 2008. Doi: 10.1002/9781119265306.ch7.
37. Fonseca JF, Torres CAA, Maffili VV, Borges AM, Santos ADF and Rodrigues MT, et al. The hypoosmotic swelling test in fresh goat spermatozoa. *Anim Reprod.* 2005;2(2):139-144. Available at: https://www.researchgate.net/publication/267683023_The_hypoosmotic_swelling_test_in_fresh_goat_spermatozoa#fullTextFileContent.
38. Twigg J, Irvine DS and Aitken RJ. Oxidative damage to DNA in human spermatozoa does not preclude pronucleus formation at intracytoplasmic sperm injection. *Hum Reprod.* 1998;13(7):1864;1871. Doi: 10.1093/humrep/13.7.1864.
39. Breininger E, Beorlegui NB, O'Flaherty C and Beconi MT. Alpha-tocopherol improves biochemical and dynamic parameters in cryopreserved boar semen. *Theriogenology.* 2005;63(8):2126-2135. Doi:10.1016/j.theriogenology.2004.08.016.
40. Silva EC, Pellinca MA, Acosta AC, Silva DMF, Filho GMA and Guerra MMP. Comparative study of DNA extraction methodologies from goat sperm and its effects on polymerase chain reaction analysis. *Genet Mol Res.* 2014;13(3):6070-6078. Doi: 10.4238/2014.August.7.21.
41. Hanson EK and Ballantyne. An Ultra-High Discrimination Y Chromosome Short Tandem Repeat Multiplex DNA Typing System. *PlosOne.* 2007; 2007(8):1-14. Doi:10.1371/journal.pone.0000688.
42. Santella RM. Approaches to DNA/RNA extraction and whole genome amplification. *Cancer Epidemiol Biomarkers Prev.* 2006;15(9):1585-1587. Doi:10.1158/1055-9965.EPI-06-0631.

COPYRIGHTS

©2023 The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited. No permission is required from the authors or the publishers.

**How to cite this article**

Ducha N, Hariani D, Budijastuti W, Susilawati T, Aulanni'am A, Wahyuningsih S. Effects of α -tocopherol addition to Brahman bull chilled semen on sperm quality, lipid peroxidation, membrane integrity and DNA integrity. Iran J Vet Sci Technol.2023; 15(1): 31-40.
DOI: <https://doi.org/10.22067/ijvst.2023.74571.1108>
URL: https://ijvst.um.ac.ir/article_43434.html