

Nephrolithiasis in two Arabian horses

Mohammadreza Aslani ^{1*}, Amir Hossein Askari ²

¹ Department of Clinical Sciences, School of Veterinary Medicine, Ferdowsi University of Mashhad, Mashhad, Iran.

² Private Veterinarian, Yazd, Iran

Received: May 20, 2009

Accepted: August 11, 2009

Abstract

Nephrolithiasis was incidentally diagnosed in 2 Arabian horses, a young foal and an adult mare, from Maybod, Yazd province, Iran. The foal had bilateral nephrolithiasis, as multiple sand like stones in renal papillary area. Large nephroliths were found in the left kidney of mare. Prominent composition of those nephroliths was magnesium ammonium phosphate. Risk factors and other aspects of nephrolithiasis in horses are discussed.

Key words: Arabian horse, nephrolith, water hardness, kidney

Corresponding author: Mohammad Reza Aslani
Email: mraslani@um.ac.ir
Tel: +98 511 8788944
Fax: +98 511 8763852

Introduction

Nephrolithiasis is uncommon in horses comparing to bladder and urethral calculi. Of 68 horses reported with urolithiasis, 59.7% had calculi in bladder and 24% had urethral calculi while 12.6% had renal calculi, and 3.7% had ureteral calculi (Lavery et al., 1992). Equine uroliths are most commonly found in the bladder (60%) or urethra (24%). The pathogenesis of nephrolith formation is uncertain and the importance of nephroliths as causes of distal urinary tract obstruction are unknown (Macbeth, 2008). Nephroliths may develop around a nidus associated with a variety of renal diseases, including polycystic kidney disease, pyelonephritis, papillary necrosis, and neoplasia. Indeed, data on upper urinary tract stones in horses are insufficient to know whether they develop spontaneously (in the absence of damage) as in humans or whether they differ significantly in mineral composition from cystic calculi. On the other hand, although nephrolithiasis are quite painful in humans, horses with nephroliths often remain asymptomatic until bilateral obstructive disease leads to development of acute or chronic renal failure (Schott, 1998). However, a history of chronic weight loss and colic in a horse with renal failure indicates the possible presence of renal calculi (Radostits et al., 2007). In the present report, for the first time in Iran, two cases of nephrolithiasis from Maybod, Yazd province are described.

A 7-year-old Arabian mare and a 7-month-old foal of same race were necropsied during investigation on an outbreak of carbamate poisoning in a small horse boarding stable in Maybod, Yazd province, Iran. Nephroliths were incidentally found in both of cases. The left kidney of adult horse was larger than normal and contained 6 calculi with different sizes. The stones were hard with irregular spongy appearance and milky in color (Fig 1). The stones weighed 750 gr.



Figure 1. One of the large nephroliths found in a 7-year-old Arabian horse.

Case report

There were numerous small sand-like and white calculi in papillary area of both kidneys in the foal. Nephroliths of the adult horse were analyzed. The result of that analysis is showed in Table 1. The composition of the nephroliths was 20% calcium oxalate, 30% calcium phosphate and 50% magnesium ammonium phosphate.

The horses were fed a mixture of wheat straw, alfalfa hay and barley grain. Drinking water provided from a well and the owner complained for its high salinity. Drinking water of horses was analyzed for total hardness (TH), total dissolved solids (TDS), some elements and pH value. The results are indicated in Table 2. There was a history of several months of low performance for adult horse.

Discussion

Although cystic calculi are the most commonly recognized form of equine uroliths, a number of renal or ureteral calculi have been reported, describing nephrolithiasis and ureterolithiasis in horses (Ehnen et al., 1990; Lavery et al., 1992; Macbeth 2008). Nephroliths may develop around a nidus

associated with a variety of renal diseases, including congenital renal dysplasias such as polycystic kidney disease, pyelonephritis, papillary necrosis and neoplasia (Schott 1998). Renal papillary necrosis is a common phenomenon secondary to concurrent dehydration and non-steroidal anti-inflammatory therapy in horses (Dusterdieck 2003). Indeed, tissue damage is believed to be the most important factor for development of urolithiasis in horses. Tissue damage accompanied by necrotic tissue, desquamated cell and leukocytes can act as a nidus, then minerals deposited around the nidus followed by slow and continuous crystal growth, the second phase of calculus formation.

Equine urine is highly alkaline, favoring crystallization of most urolith components, especially calcium carbonate (Dusterdieck 2003).

Regardless of location, almost equine calculi are predominantly composed of calcium carbonate, usually in calcite or vaterite forms (Diaz-Espineira et al., 1997). However, predominant composition of calculi found in the adult Arab horse of this report was struvite (magnesium ammonium phosphate), which is less common component of equine urolithiasis (Schott 1998). High physical activities and insufficient replacement of water losses, as well as high ambient temperature increase the incidence and prevalence of urolithiasis (Siener and Hesse 2003). On the other hand a number of studies on human, beings have suggested that water hardness and magnesium levels in drinking water can contribute in greater incidence of urolithiasis (Bellizzi et al., 1999; Medina-Escobedo et al., 2002; Sahinduran et al., 2007). According to the water analysis in Maybod region, drinking water showed high hardness. Although the effect of water hardness on urolithiasis is not known in equine, but it can be attributed to imbalances of minerals intakes that may be a predisposing factor in the development of uroliths.

Table 1. Chemical Values of water sample.

Parameter	Water sample	Drinking water
TDS	2800 mg/l	1000 mg/kg
TH	1045 mg/l	100 mg/dl
Ca	320 mg/l	75 mg/dl
Mg	70 mg/l	50 mg/dl
Na	1012 mg/l	175
K	7.8 mg/l	12
Cl	2612 mg/l	200 mg/l
Sulfate	122 mg/l	200 mg/l
HCO ₃ ⁻	620 mg/l	-
Ammonium	0.25 mg/l	0.0002 mg/l
Phosphate	0.4 mg/l	0.1 mg/l
pH	8.2	7

Although any breed predisposition for nephrolithiasis has not been described, it has been speculated that racehorses are at a greater risk, because of the common use of nonsteroidal anti-inflammatory drugs and then associated risk of papillary necrosis.

Table 2. Constituents of analysed nephroliths

Element	%
Phosphate	30
Magnesium	7
Calcium	47
Oxalate	12
Ammonium	4

Horses with nephrolithiasis often remain asymptomatic until bilateral obstructive disease results in acute or chronic renal failure. Nonspecific signs of uremia such as poor performance, inappetance, and weight loss are commonly observed (Dusterdieck 2003). Diagnosis of upper urinary uroliths is usually made during rectal or ultrasonographic examination. Rectal palpation may reveal an enlarged kidney or ureter and in some instances, the calculus can be palpated on the enlarged ureter. In ultrasonography calculi are detected as hyperechoic structures with strong anechoic shadow. Small stones lesser than 1 cm in diameter can be missed despite of complete examination, but other findings such

as dilatation of renal pelvic and hydronephrosis supports the diagnosis (Schott 1998; Dusterdieck 2003). However, upper urinary tract stones may be an incidental finding at necropsy (Ehnen et al., 1990; Schott 1998). For decreasing the possibility of urolith formation, the owners should be cautioned to provide safe drinking water and balanced ration for horses and to avoid overdose non-steroidal anti-inflammatory drugs.

References

- Bellizi, V., L. De Incola, R. Minutolo, D. Russo, B. Cianciaruso, and M. Andrucci. 1999. Effects of water hardness on urinary risk factors for kidney stones in patient with idiopathic nephrolithiasis. *Nephron*. 81:66-70.
- Diaz-Espineira, M., J. Nellanato, and M. A. De La Fuente. 1997. Infrared and atomic spectrometry analysis of the mineral composition of a series of equine sabulous material and urinary caculi. *Res. Vet. Sci*. 63:93-95.
- Dusterdieck, K. F. 2003. Obstructive disease of the urinary tract. . Pages 832-834 *in* Current Therapy in Equine Medicine. N. E. Robinson, ed. Saunders, St-Louis.
- Ehnen, S. J., T. J. Divers, D. Gillette, and V. B. Reef. 1990. Obstructive nephrolithiasis and ureterolithiasis associated with chronic renal failure in horses: eight cases (1981-1987). *J. Am. Vet. Med. Assoc*. 197:249-253.
- Laverty, S., J. R. Pascoe, G. V. Ling, J. P. Lavoie, and A. L. Ruby. 1992. Urolithiasis in 68 horses. *Vet. Surg*. 21:56-62.
- Macbeth, B. J. 2008. Obstructive urolithiasis, unilateral hydronephrosis and probable nephrolithiasis in a 12-year old Clydesdale gelding. . *Can. Vet. J*. 49:287-290.
- Medina-Escodedo, M., M. Zaidi, E. Real-de-Leon, and S. Orozco-Rivadeneira. 2002. Prevalence and risk factors of urinary lithiasis in Yucatan, Mexico. *Salud Pulica de Mexico*. 44:1-5.
- Newton, S. A., M. A. Cheeseman, and G. B. Edwards. 1999. Bilateral renal and ureteral calculi in a 10-year-old gelding. *Vet Rec*. 144:383-385.
- Radostits, O. M., C. C. Gay, K. W. Hinchcliff, and P. D. Constable, eds. 2007. *Veterinary Medicine*. 10 ed. Saunders & Elsevier, London.
- Sahinduran, S., T. Buyukoglu, M. S. Gulay, and F. Tasci. 2007. Increased water hardness and magnesium levels may increase occurrence of urolithiasis in cows from the Brder region (Turkey). *Vet. Res. Commun*. 31:665-671.
- Schott, H. C. 1998. Obstructive disease of the urinary tract. Pages 880-890 *in* Equine Internal Medicine. S. M. Reed, Bayly, W. M., ed. Saunders, Philadelphia.
- Siener, R., Hesse, A. 2003. Fluid intake and epidemiology of urolithiasis. *Europ. J. Clin. Nutr*. 57:547-551.

سنگ کلیوی در دو اسب نژاد عرب

محمد رضا اصلانی^۱، امیر حسین عسگری^۲

^۱ گروه علوم در مانگامی، دانشکده دامپزشکی دانشگاه فردوسی مشهد، مشهد، ایران

^۲ دامپزشک بخش خصوصی، یزد، ایران

دریافت مقاله: ۸۸/۲/۳۰ پذیرش نهایی: ۸۸/۵/۲۰

چکیده

سنگ کلیوی به طور تصادفی در دو اسب نژاد عرب، یک کره اسب جوان و یک مادبان درمیید استان یزد در ایران تشخیص داده شد. کره اسب دارای سنگ کلیوی دوطرفه به صورت سنگ ریزه های متعدد در ناحیه لگنچه کلیه بود. سنگ های کلیوی بزرگ در کلیه چپ مادبان یافت شدند. ترکیب غالب این سنگها فسفات آمونیوم منیزیم بود. عوامل خطر ساز و سایر جنبه های ایجاد سنگ کلیوی در اسبها در این مقاله مورد بحث قرار می گیرند.

واژه های کلیدی: اسب عرب، سنگ کلیه، سختی آب، کلیه

Iranian Journal of Veterinary Science and Technology

Instruction for Authors

Scope of Journal

Iranian journal of Veterinary Science and Technology (IJVST) is a peer-reviewed multi-disciplinary journal, publishes important new advances in research areas and case histories covering all aspects of veterinary medicine and its relevant subjects including: anatomy, bacteriology, biochemistry, biotechnology, food hygiene, immunology, molecular biology, parasitology, pathology, pharmacology, physiology and virology. Contributions related to clinical sciences including large and small animals, poultry, equine species and aquaculture are welcomed. Articles can comprise basic sciences, as well as applied veterinary findings and experimental studies and their impact on diagnosis, treatment and prevention of diseases. This journal is currently published twice a year.

General information

1. Submitted manuscripts should not be previously published elsewhere and should not be under consideration by any other journal.

2. The submitted manuscript should be accompanied by a written statement signed by the corresponding author on behalf of all the authors that its publication has been approved by all coauthors, stating that the whole manuscript or a part of it has not and will not be published in or submitted to elsewhere.

3. The corresponding author should provide all co-authors with information regarding the manuscript, and obtain their approval before submitting any revisions.

4. Full-length articles, Short Communications, Case reports, Reviews and Commentaries that address scope of the journal will be considered for publication.

5. Ethics: Authors must state that the protocol for the research project has been approved by the Ethics Committee of the institution within which the work was undertaken. Authors are responsible for animal welfare and all statements made in their work.

Manuscript Preparation

Manuscripts should be written in English, with Abstract in both English and Persian (where applicable), must be typewritten in the format of the word processing program, double-spaced, in 12-point Times New Roman font on A4 paper format. Authors are requested to reserve margins of 2.5 cm all around the paper. Manuscript should also have line numbers. Original photos, tables and figures should be included together with the manuscripts.

All pages of the manuscripts should be enumerated from the title page.

In review articles, one or more writers attempt to sum up the current state of the research on a particular topic. Review articles presented by the main people working in a specific field should provide an update on recent advances in a particular field. Authors wishing to submit review articles should contact the Editor with an outline of the proposed paper prior to submission, to allow consideration by the scientific editors.

Short communications and Case reports should include a full Title Page, Abstract, Keywords, Introduction, Case Presentation, Results and Discussion, Acknowledgement, References, Tables, Figures, and Legends -limited to 2000 words excluding the references and two tables and/or figures.

Original articles should contain a full Title Page, Abstract, Keywords, Introduction, Materials and Methods, Results, Discussion, Acknowledgement, References, Tables, Figures, and Legends.

Full **Title Page** should include title (title of the article should be concise but informative), author(s) (include the complete name, academic degree, department affiliation, and institution), running head (≤ 50 characters, including spaces), name and address of the authors to whom correspondence and reprint requests should be addressed.

Abstract should contain the purpose of the study, findings (including its statistical significance) and the conclusion made on the basis of the findings should be clearly stated in the abstract-limited to 250 words. Authors who are not native Persian speakers may submit their manuscript with an abstract in English only. Do not use abbreviations and reference citation in abstracts.

Keywords: For indexing purposes, each submitted manuscript should include three to five keywords, following the abstract and preferably chosen from the Medical Subject Headings (MESH). Keywords should express the precise content of the manuscript.

Introduction should be as concise as possible, and clearly explain the main objective and hypothesis of the investigation.

Materials and Methods should be described in sufficient details to allow other workers to reproduce the results. Specify any statistics computer programs used. The methods of data collection and use of statistical analysis will be checked by the referees and if necessary, a statistician. Drugs and therapeutic agents, reagents, softwares and equipments should be given in the format: name (trade name, manufacturer name, city, country), e.g. Statview 5 (SAS Institute, Inc., Cary, NC, USA).

Results indicate the results of an original research in a clear and logical sequence. Do not repeat data that are already covered in tables and illustrations. In manuscripts describing more than one animal, all animals should be assigned a case number.

Discussion should emphasize the new and important aspects of the study and the conclusions that follow from them. It should describe the implication of the findings and their limitations, relates the observations to other relevant studies, and links the conclusions with the goals of the study. Recommendations, when appropriate, may be included.

Acknowledgements: Personal acknowledgement, sources of financial support, contributions to accomplish the submitted manuscript and everything that does not justify authorship should be mentioned in this section, if required.

References must be up-to-dated and limited to those that are necessary. Lists of references should be given in date order in the text, but alphabetically in the reference list. Listed below, are sample references to a book, a chapter in a book, journal articles, proceedings, thesis, and online document respectively, in the correct format.

Book

Halliwell, B. and J. Gutteridge. 1999. Free Radicals in Biology and Medicine. 3 ed. Oxford University Press, New York.

Chapter in a Book

Courtney, C. H. 2001. Chemotherapy of parasitic diseases. Pages 947-1017 in Veterinary Pharmacology and Therapeutics. 8th. ed. R. H. Adams, ed. Iowa State University Press, Iowa.

Journal Article

Sultana, R., D. Boyd-Kimball, J. Cai, W. M. Pierce, J. B. Klein, M. Merchant, and D. A. Butterfield. 2007. Proteomics analysis of the Alzheimer's disease hippocampal proteome. *J Alzheimers Dis* 11(2):153-164.

Deepak, M. and S. S. Handa. 2000. Antiinflammatory activity and chemical composition of extracts of *Verbena officinalis*. *Phytother Res* 14(6):463-465.

Conference proceedings

Seifi, H. A., M. Mohri, and J. Kalamati Zade. 2002. The study on correlation of serum calcium and inorganic phosphorus values with urine pH in dry cows. in 22nd World Buiatrics Congress. Hannover , Germany.

Dissertation or Thesis

Afkhami, A. 2007. Protective effects of crocin in neurodegenerative disorders, Role of Proteinase-activated receptor-2. University of Tehran, Tehran.

Please cite references in the text by name and year in parentheses. Some examples:

(Courtney, 2001)
(Halliwell and Gutteridge, 1999)
(Seifi et al., 2002)
(Deepak and Handa, 2000, Sultana et al., 2007)

The list of references should only include works that are cited in the text and that have been published or accepted for publication. Reference list should be alphabetized by the last names of the first author of each work.

Please apply the standard abbreviation of a journal's name according to the ISSN List of Title Word Abbreviations, for more information please see:

www.issn.org/2-22661-LTWA-online.php

Tables should be self-explanatory, clearly arranged, and supplemental to the text. Tables should provide easier understanding and not duplicate information already included in the text or figures. Each table should be typewritten with double spacing on a separate page and numbered in order of citation in the text with Arabic numerals. Each table should have a concise heading that makes it comprehensible without reference to the text of the article. Explain any nonstandard abbreviations in a footnote to the table.

Figures must be submitted in high-resolution (300 dpi or greater) and should be initially saved in a neutral data format such as JPEG, TIFF. Illustrations should be numbered as cited in the sequential order in the text, with a legend on a separate sheet. Color photographs are welcome at no extra charge. The editors and publisher reserve the right to reject illustrations or figures based upon poor quality of submitted materials.

Use of Italics

Gene symbols, Latin terms (i.e. *in vivo*, *in vitro*, *ex vivo*, *in utero*, *in situ*, and etc.) and species scientific names (using the binomial nomenclature), should be typed in italics, while the first letter of the genus name must be capitalized (i.e. *Homo sapiens*).

Copyright

Accepted manuscripts become the permanent property of the IJVST and may not be reproduced, in whole or in part, without the written permission of the Editor.

Submission

Authors should submit their manuscript in electronic format directly through <https://ijvst.um.ac.ir> or as an e-mail attachment to ijvst@um.ac.ir with a transmittal letter signed by the author to whom correspondence should be addressed. The entire article should be submitted as a single file. Please ensure that your e-mail address and full postal address are included on the front page of the manuscript.

For further information, please contact the Editorial Office through:

Dr. Ahmad Reza Movassaghi, DVM, PhD.
Editor-in-Chief
Iranian Journal of Veterinary Science and Technology
Tel: +98 511 8788944
Fax: +98 511 8763852
Email: ijvst@um.ac.ir