



## Study of Toxic Effects of Oxothiazole Derivative as a New Antibacterial Agent

Hamid Beyzaei<sup>a</sup>, Abbas Jamshidian<sup>b</sup>, Mohammad Reza Hajinezhad<sup>c</sup>, Seyed Hadi Hashemi<sup>d</sup>,  
Nasim Amel<sup>e</sup>, Maliheh Abdollahi<sup>f</sup>, Fatemeh Arab-Sahebi<sup>g</sup>, Behzad Ghasemi<sup>f</sup>

<sup>a</sup>Department of Chemistry, Faculty of Science, University of Zabol, Zabol, Iran. <sup>b</sup>Department of Pathobiology, Faculty of Veterinary Medicine, University of Zabol, Zabol, Iran. <sup>c</sup>Department of Basic Sciences, Faculty of Veterinary Medicine, University of Zabol, Zabol, Iran. <sup>d</sup>Department of Clinical Sciences, Faculty of Veterinary Medicine, Zabol University, Zabol, Iran. <sup>e</sup>Student Research Committee, Torbat Heydariyeh University of Medical Sciences, Torbat Heydariyeh, Iran. <sup>f</sup>Torbat Jam Faculty of Medical Sciences, Torbat Jam, Iran. <sup>g</sup>Student Research Committee, Sabzevar University of Medical Sciences, Sabzevar, Iran.

### ABSTRACT

The spread of antibiotic-resistant bacteria in many humans and animals has driven researches to identify and design novel antibacterial agents. *In vitro* inhibitory activity of (2E)-2-(4,5-dihydro-4-oxothiazol-2-yl)-2-(thiazolidin-2-ylidene) acetonitrile against many bacterial pathogens has been proven in both veterinary and human medicine. In this study, its *in vivo* toxic effects was studied in mice. The median lethal dose (LD50) value of 239.88 mg/kg was estimated using intraperitoneal injection in 8 groups of mice after 48 h treatment. Then, intraperitoneal injections of LD50 of oxothiazole solution into 4 other mice were done to evaluate histopathological changes in their liver and kidney tissues. The histopathological studies were identified as fatty change, hepatitis, necrosis and regeneration in liver, and fibrosis, necrosis, nephritis, hyaline cast and hyperaemia in kidney. In conclusion, the synthesized oxothiazole derivative causes renal and hepatic toxicity in mice at medium concentrations. The change of thiazole substituents and complexation may reduce its toxicity

### Keywords

Toxic effects, Histopathological study, Oxothiazole derivative

### Abbreviations

LD50: Median Lethal Dose

## Introduction

Species of pathogenic bacteria have developed resistance to current antibiotics such as penicillin, amoxicillin, clindamycin, tetracycline, erythromycin and sulfamethoxazole due to overuse of drugs in humans and animals [1]. This health problem increases costs of medical care [2]. In recent years, many researchers were encouraged to discover new antibacterial agents including herbal extracts, metal nanoparticles and heterocyclic derivatives [3].

Thiazoles are a major class of heterocyclic compounds, which have been used in the treatment of diseases such as cancer, hypertension and lipid disorders [4]. Significant antibacterial effects were observed by thiazole derivatives. It was found that they can inhibit the activity of DNA gyrase B and eCKAS III enzymes, and were effective against the replication of DNA and the synthesis of fatty acids, in bacteria [5].

*In vitro* inhibitory activities of thiazoles have been assessed against a variety of pathogenic bacteria including both Gram-positive and Gram-negative strains [6-8]. These compounds inhibit the growth of bacteria by blocking ion channels as well as inhibiting enzymatic and cellular activities [9-12]. However, there are a few *in vivo* studies on body and organ of animals.

Oxothiazole derivative is a recently synthesized heterocyclic compound that its *in vitro* inhibitory activities were evaluated against some Gram-positive and Gram-negative bacteria such as *Staphylococcus aureus*, *Streptococcus agalactiae*, *Listeria monocytogenes*, *Bacillus cereus*, *Escherichia coli* and *Salmonella typhimurium* [13-16]. The present study was performed to assess the function and histology of the liver and kidney in mice exposed to the synthetic oxothiazole.

## Results

The toxic and therapeutic effects of oxothiazole derivative were evaluated in 8 groups of 8 mice each. The 48 h LD50 value of compound was 239.88 mg/kg (Tables 1 and 2, Figure 1). Histopathological changes including fatty change, hepatitis, necrosis and regen-

**Table 1**

Mortality percent in each group after IP injection of oxothiazole.

Dose mg/kg	log	Died	Percent	Lived	Total
160	2.20	0	0	8	8
200	2.30	1	12.5	7	8
220	2.34	2	25	6	8
240	2.38	4	50	4	8
265	2.42	6	75	2	8
300	2.47	8	100	0	8
350	2.54	8	100	0	8

**Table 2**


Different LD doses of oxotiazole

LD10	190.55
LD40	230.14
LD50	239.88
LD80	267.92
LD90	278.61


eration in liver, and fibrosis, necrosis, nephritis, hyaline cast and hyperaemia in kidney were observed as illustrated in Figures 2 and 3.

## Discussion

Thiazole ring is an essential part of numerous biologically active compounds. Thiazole derivatives are recognized to affect cell cycle progression, protein production and DNA replication [17-19]. Sulfathiazole as an antibiotic containing thiazole ring was used in the treatment of gastrointestinal and respiratory infections. Its oral LD50 in mice is 4500 mg/kg. Although toxicity of the drug is low, it is prescribed less frequently in humans due to its side effects such as high fever, severe headache, stomatitis, conjunctivi-



**Figure 1**  
Mortality of mice infected with oxothiazole




**Figure 2**  
Histopathological changes of liver tissue in mice treated with oxothiazole

tis, rhinitis, urethritis and balanitis [20]. Febuxostat is a xanthine oxidase inhibitor containing thiazole ring that lowers uric acid levels in blood. The LD50 of drug administered orally to mice is 300 mg/kg [21]. Sprycel (dasatinib) is a chemotherapy medication used to treat chronic myelogenous leukemia (CML) and acute lymphoblastic leukemia (ALL). It has a relatively high toxicity (oral LD50 = 100 mg/kg) in rat. Its consumption may cause anemia, swelling, pulmonary edema, heart failure, rash and diarrhea [22].

Anticoagulant and antiarrhythmic effects of some thiazole derivatives were studied by Abd El-Galil et al. in 2009 via determination of LD50 values of 292-736 mg/kg. Although oxothiazole is more toxic than these derivatives, the determination of median effective dose (ED50) and therapeutic index (TI) of this heterocycle in future researches is necessary for a better conclusion [23]. Siddiqui et al. in 2010 calculated the toxic effects of 3-[4-(substituted phenyl)-1,3-thiazol-2-ylamino]-4-(substituted phenyl)-4,5-dihydro-1H-1,2,4-triazole-5-thiones as new anticonvulsant agents, their LD50 values (834 and 936 mg/mL) are higher than those in phenytoin (a common anticonvulsant drug with LD50 = 224 mg/mL) [24].

Efficiency of some synthesized imidazo[2,1-b]thiazole derivatives as diuretic agents was proven in comparison with those of hydrochlorothiazide and furosemide, LD50 values were reported in the range of 1000 to 1500 mg/mL [25].

Among 12 thiazolidines synthesized by Wang et al., only three derivatives showed significant anti-inflammatory activities. They can be used to treat



**Figure 3**  
Histopathological changes of kidney tissue in mice treated with oxothiazole

hepatic injuries according to the results obtained by further Con A-induced acute liver injury of BALB/c mice [26]. Thiazole-Zn is a chinese-created systemic fungicide that had no pathological effects on the liver and kidney of female rats when used as oral gavage, while its destructive effects were observed in the thyroid gland [27].


The toxicity of oxothiazole derivative has been preliminarily studied in mice at doses of 350 and 160 mg/kg without determination of LD50. All mice died at dose of 350 mg/kg after 24 h. The liver of mice treated at dose 160 mg/kg was evaluated after 48 h [15]. To extend toxicological studies, some biochemical and physiological effects of this potential antibacterial agent were assessed via determination of LD50 value and evaluation of histopathological changes in the liver and kidney of mice. Renal and hepatic toxicity were observed in mice when medium doses of heterocyclic derivative were injected. New oxothiazole derivatives with low toxicity can be designed in future studies by changing thiazole substituents and complexation.

## Material and methods

### General procedure for the synthesis of oxothiazole compound

Oxothiazole (3) was synthesized according to a previous procedure [28] as follows: a suspension of 1 mmol of each (*E*)-2-cyano-2-(thiazolidin-2-ylidene) ethanethioamide (1), ethyl bromoacetate (2) and sodium bicarbonate ( $\text{NaHCO}_3$ ) in 1 mL DMF as solvent was stirred at room temperature for 8 h. 5 mL of water was added to the reaction mixture. The precipitate was filtered out, and washed with water (1 × 5 mL) and ethanol (1 × 5 mL),



**Figure 4**

Synthesis of oxothiazole derivative 3  
(2E)-2-(4,5-Dihydro-4-oxothiazol-2-yl)-  
2-(thiazolidin-2-ylidene)acetonitrile.

respectively. The pure thiazole 3 was obtained in 75% yield after recrystallization from acetonitrile, and its chemical structure was confirmed through the elemental analysis and spectroscopic data (Figure 4).

m.p. 248-252 °C; IR (KBr)  $\nu$ : 3422 (NH), 2193 (C≡N), 1688 (C=O), 1573 (C=C)  $\text{cm}^{-1}$ ;  $^1\text{H NMR}$  (DMSO- $d_6$ )  $\delta$ : 3.63 (t,  $J$  = 7.8 Hz, 2H,  $\text{SCH}_2$ ), 4.01 (s, 2H,  $\text{COCH}_2$ ), 4.12 (t,  $J$  = 7.8 Hz, 2H,  $\text{NHCH}_2$ ), 10.53 (s, 1H, NH) ppm; MS (EI,  $m/z$ ): 225 ( $M^+$ , 21), 176 (37), 150 (94), 109 (28), 46 (100); Anal. Calcd for  $\text{C}_8\text{H}_7\text{N}_3\text{OS}_2$ : C, 42.65; H, 3.13; N, 18.65. Found: C, 42.74; H, 3.00; N, 18.46.

### In vivo study

68 white male BALB/c mice, weighing 26-34 g, were obtained from the Central Animal House, University of Zabol, Zabol, Iran. Mice were maintained at all stages with free access to standard food and refined water under  $25 \pm 2$  °C temperature, 50% humidity, 12/12 light dark cycle [24].

### Assessment of median lethal dose (LD50)

An up-and-down method was applied for acute toxicity testing. 64 mice were divided into 8 groups. Finally, doses including 160, 200, 220, 240, 265, 300 and 350 mg/kg of oxothiazole in 10% DMSO were selected for intraperitoneal injection (IP). 10% DMSO was also injected as negative control. Mice mortality was recorded in each group during 48 h [24].

### Histopathological examination

4 BALB/c were injected intraperitoneally with the median lethal doses of oxothiazole. The abdominal walls of dead and live mice were excised immediately or after ether-euthanasia. The liver and kidney of mice were removed from their abdominal cavity, and stored in 10% formalin. All solutions were refreshed three times every 24 h. The histological sections were molded in 4- to 6- $\mu\text{m}$ -thick paraffin according to the histopathological principles. The prepared slides were stained with hematoxylin and eosin, and examined under a light microscope [27].

### Acknowledgment

This work was supported by the University of Zabol under Grant number UOZ-GR-9517-15.

### Author Contributions

All authors contributed to the design of study, data analysis and manuscript preparation.

### Conflict of Interest

The authors declare that there is no conflict of interest.

### References

1. Wang D, Zhang L, Zhou X, He Y, Yong C, Shen M, et al. An-

- timicrobial susceptibility, virulence genes, and randomly amplified polymorphic DNA analysis of *Staphylococcus aureus* recovered from bovine mastitis in Ningxia, China. *J Dairy Sci.* 2016;99(12):9560-9.
- Ahumada-Santos YP, Soto-Sotomayor ME, Baez-Flores ME, Diaz-Camacho SP, Lopez-Angulo G, Eslava-Campos CA, et al. Antibacterial synergism of *Echeveria subrigida* (B. L. Rob & Seaton) and commercial antibiotics against multidrug resistant *Escherichia coli* and *Staphylococcus aureus*. *Eur J Int Med.* 2016;8(5):638-44.
- Santos L, Ramos F. Analytical strategies for the detection and quantification of antibiotic residues in aquaculture fishes: A review. *Trends Food Sci Technol.* 2016;52:16-30.
- Bakavoli M, Beyzaei H, Rahimizadeh M, Eshghi H. Regioselective synthesis of 2[(E)(benzo[d]thiazol-2(3H)ylidene)(cyano)methyl]thiazoles. *Heterocycl Commun.* 2011;17:151-4.
- Ghasemi B, Sanjarani G, Sanjarani Z, Majidiani H. Evaluation of anti-bacterial effects of some novel thiazole and imidazole derivatives against some pathogenic bacteria. *Iran J Microb.* 2015;7(5):281-6.
- Beyzaei H, Aryan R, Molashahi H, Zahedi MM, Samzadeh-Kermani A, Ghasemi B. MgO nanoparticle-catalyzed, solvent-free Hantzsch synthesis and antibacterial evaluation of new substituted thiazoles. *J Iran Chem Soc.* 2017;14(5):1023-31.
- Pawar CD, Sarkate AP, Karnik KS, Bahekar SS, Pansare DN, Shelke RN, et al. Synthesis and antimicrobial evaluation of novel ethyl 2-(2-(4-substituted)acetamido)-4-substituted-thiazole-5-carboxylate derivatives. *Bioorg Med Chem Lett.* 2016;26(15):3525-8.
- Qin YJ, Wang PF, Makawana JA, Wang ZC, Wang ZN, Gu Y, et al. Design, synthesis and biological evaluation of metronidazole-thiazole derivatives as antibacterial inhibitors. *Bioorg Med Chem Lett.* 2014;24(22):5279-83.
- Washburn DG, Holt DA, Dodson J, McAtee JJ, Terrell LR, Barton L, et al. The discovery of potent blockers of the canonical transient receptor channels, TRPC3 and TRPC6, based on an anilino-thiazole pharmacophore. *Bioorg Med Chem Lett.* 2013;23:4979-84.
- Kang SY, Song KS, Lee J, Lee SH, Lee J. Synthesis of pyridazine and thiazole analogs as SGLT2 inhibitors. *Bioorg Med Chem Lett.* 2010;18:6069-79.
- Andreani A, Granaiola M, Guardigli M, Leoni A, Locatelli A, Morigi R, et al. Synthesis and chemiluminescent high throughput screening for inhibition of acetylcholinesterase activity by imidazo[2,1-b]thiazole derivatives. *Eur J Med Chem.* 2005;40:1331-4.
- Andreani A, Granaiola M, Leoni A, Locatelli A, Morigi R, Rambaldi M. Effects of new ubiquinone-imidazo[2,1-b]thiazoles on mitochondrial complex I (NADH-ubiquinone reductase) and on mitochondrial permeability transition pore. *Bioorg Med Chem Lett.* 2004;12:5525-32.

13. Ghasemi B, Beyzaei H, Hashemi SH, Majidiani H. Study of antibacterial effect of novel thiazole, imidazole and tetrahydropyrimidine derivatives against *Escherichia coli*. *J Med Bacteriol*. 2015;4(6):7-12.
14. Ghasemi B, Beyzaei H, Hashemi SH. Study of antibacterial effect of novel thiazole, imidazole, and tetrahydropyrimidine derivatives against *Listeria Monocytogens*. *Ann Mil Health Sci Res*. 2015;13:101-5.
15. Ghasemi B, Najimi M, Beyzaei H, Jamshidian A. Antibacterial effects of novel thiazole derivatives and the toxicity of oxothiazole on liver histopathology of mice. *Med Lab J*. 2015;9(4):6-12.
16. Ghasemi B, Beyzaei H, Majidiani H. A comparative study on the antibacterial effects of some newly synthesized thiazole, imidazolidine and tetrahydropyrimidine derivatives against *Bacillus cereus* and *Salmonella typhimurium*. *Pharm Sci*. 2016;22(1):54-9.
17. Chen H, Song J, Wang J, Xu C, Chen C, Gu W. Synthesis and biological evaluation of thiazole derivatives as novel USP7 inhibitors. *Bioorg Med Chem Lett*. 2017;27(4):845-9.
18. Reichelt A, Bailis JM, Bartberger MD, Yao G, Shu H, Kaller MR. Synthesis and structure-activity relationship of trisubstituted thiazoles as Cdc7 kinase inhibitors. *Eur J Med Chem*. 2014;80:364-82.
19. Chen YL, Cherry K, Michael L, Ebbinghaus CF, Gamlath CB, Liston D. Thiazole-diamides as potent  $\gamma$ -secretase inhibitors. *Bioorg Med Chem Lett*. 2007;17(20):5518-22.
20. Prinsen MK, Romijn T, Snoeij NJ. Skin sensitization testing: the relevance of rechallenge and pretreatment with sodium lauryl sulfate in the guinea pig maximization test. *Food Chem Toxicol*. 1997;35(9):923-6.
21. Love BL, Barrons R, Veverka A, Snider KM. Urate-lowering therapy for gout: focus on febuxostat. *Pharmacother*. 2010;30(6):594-608.
22. Keating, GM. Dasatinib: A review in chronic myeloid leukaemia and Ph+ acute lymphoblastic leukaemia. *Drugs*. 2017;77(1):85-9.
23. Abd El-Galil EA, Sabrry NM, Abdalla MM, Abdel-Wahab BF. Synthesis, antiarrhythmic and anticoagulant activities of novel thiazolo derivatives from methyl 2-(thiazol-2-ylcarbonyl)acetate. *Eur J Med Chem*. 2009;44:725-35.
24. Siddiqui N, Ahsan W. Triazole incorporated thiazoles as a new class of anticonvulsants: Design, synthesis and in vivo screening. *Eur J Med Chem*. 2010;45:1536-43.
25. Andreani A, Rambaldi M, Mascellani G, Rugarli P. Synthesis and diuretic activity of imidazo[2,1-b]thiazole acetohydrazones. *Eur J Med Chem*. 1987;22(1):19-22.
26. Wang G, Deng C, Xie C, Ma L, Yang J, Qiu N, et al. Synthesis and biological evaluation of novel dimethyl[1,10-biphenyl]-2,20-dicarboxylate derivatives containing thiazolidine-2,4-dione for the treatment of concanavalin A-induced acute liver injury of BALB/c mice. *Eur J Med Chem*. 2011;46:5941-8.
27. Yang H, Zhang W, Kong Q, Liu H, Sun R, Lin B, et al. Effects of pubertal exposure to thiazole-Zn on thyroid function and development in female rats. *Food Chem Toxicol*. 2013;53:100-4.
28. Bakavoli M, Beyzaei H, Rahimizadeh M, Eshghi H, Takjoo R. Regioselective synthesis of new 2-(E)-cyano(thiazolidin-2-ylidene)thiazoles. *Molecules*. 2009;14:4849-57.

